

REFERENCES

- Abubaker, T. A., Y. Matsui and H. Naka, 2024(a). Identification of the female sex pheromone of *Bastilla arctotaenia* (Lepidoptera: Erebidae). *Biosci. Biotechnol. Biochem.*, **88**: 577-580.
- Abubaker, T. A. S., Y. Matsui and H. Naka, 2024(b). Identification of the female sex pheromone of *Grammodes geometrica* (Lepidoptera: Erebidae: Erebinae). *J. Chem. Ecol.*, **50**: 866-873.
- Acín, P., G. Rosell, A. Guerrero and C. Quero, 2010. Sex pheromone of the Spanish population of the beet armyworm *Spodoptera exigua*. *J. Chem. Ecol.*, **36**: 778-786.
- Adachi, Y., N. D. Do, M. Kinjo, S. Makisako, R. Yamakawa, K. Mori and T. Ando, 2010. Positions and stereochemistry of methyl branches in the novel sex pheromone components produced by a lichen moth, *Lyclene dharma dharma*. *J. Chem. Ecol.*, **36**: 814-823.
- Adati, T. and S. Tatsuki 1999. Identification of female sex pheromone of the legume pod borer, *Maruca vitrata* and antagonistic effects of geometrical isomers. *J. Chem. Ecol.*, **25**: 105-115.
- Adati, T., S. Kato, W. Toriumi, K. Kawazu, W. Susila, K. Sumiarta, G. Susatika, Dadang and I. M. Samudra, 2013. Field trials of synthetic sex pheromone lures for the large cabbage-heart caterpillarmoth, *Crocidolomia pavonana* (Lepidoptera: Crambidae) in Indonesia. *Appl. Entomol. Zool.*, **48**: 113-118.
- Adler, V. E., M. Jacobson, J. F. Edmiston and M. H. Fleming, 1976. (Z,E)-9,12-Tetradecadien-1-ol acetate: a potent attractant for male *Ephesiodes infimella*. *J. Econ. Entomol.*, **69**: 706-708.
- Adler, V. E., 1983. Adult Sesiidae of Maryland collected in traps baited with isomers of 3,13-octadecadien-1-ol acetate fruit orchards. *J. Environ. Sci. Health.*, **A 18**: 611-619.
- Akter, S., MD S. Hossain, R. Ali, R. Regmi, S. J. Park and B. Mainali, 2025. Variations in sex pheromone of the Australian population of fall armyworm: influence of age and mating status. *J. Chem. Ecol.*, **51**: 55.
- Albages, R., R. Canela, M. Eizaguirre, M. Riba and A. Guerrero, 1988. (Z)-9-Tetradecen-1-ol, an attractant for *Athetis Hospes* Freyer. *Colloq. INRA (Mediateurs Chim.: Comportement Syst. Lepidopteres, Appl. Agron.)* **46**: 55-58. [*Chem. Abst.*, 110: 52909n]
- Albert, von R., H. Bogenschütz and E. König, 1984. Investigation on the use of pheromone-baited traps for monitoring the population

- dynamics of *Operophtera brumata* L. (Lepid., Geometridae). *Z. Angew. Entomol.*, **98**: 286-298.
- Alford, D. V., 1978. Observations on the specificity of pheromone-baited traps for *Cydia funebrana* (Treitschke) (Lepidoptera: Tortricidae). *Bull. Entomol. Res.*, **68**: 97-103.
- Alford, A. R., P. J. Silk, M. McClure, C. Gibson and J. Fitzpatrick, 1983. Behavioural effects of secondary components of the sex pheromone of the eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **115**: 1053-1058.
- Alford, A. R. and P. J. Silk, 1984. Behavioral effects of secondary components of sex pheromone of western spruce budworm (*Choristoneura occidentalis*) Free. *J. Chem. Ecol.*, **10**: 265-270.
- AliNiazee, M. T., 1976. Field studies on sex pheromone trapping of the filbert leafroller, *Archips rosanus*, in Oregon. *Ann. Entomol. Soc. Am.*, **69**: 820-824.
- Allison, J. D. and R. T. Cardé, 2006. Heritable variation in the sex pheromone of the almond moth, *Cadra cautella*. *J. Chem. Ecol.*, **32**: 621-641.
- Alm, S. R., R. N. Williams, D. M. Pavuk, J. W. Snow and M. A. Heinlein, 1989. Distribution and seasonal flight activity of male grape root borers (Lepidoptera: Sesiidae) in Ohio. *J. Econ. Entomol.*, **82**: 1604-1608.
- Altamar-Varón, P., D. Pérez-Maldonado, D. Rodríguez-Caicedo, C. Guerrero-Perilla and E. Coy-Barrera, 2016. Chemical composition of the low-polar fraction of the *Copitarsia uncilata* Burgos & Leiva (Lepidoptera: Noctuidae) eversible pheromone gland. *Neotropical Entomol.*, **45**: 734-739.
- Altesor, P., C. Rossini, P. H. G. Zarbin and A. Gonzalez, 2009. Sex pheromone of the bud borer *Epinotia aporema*: chemical identification and male behavioral response. *J. Chem. Ecol.*, **35**: 349-354.
- Anagnostakis, S. L., K. M. Welch, J. W. Snow, K. Scarborough and T. D. Eichlin, 1994. The rediscovery of the clearwing chestnut moth, *Synanthedon castaneae* (Busck) (Lepidoptera: Sesiidae) in Connecticut. *J. N.Y. Entomol. Soc.*, **102**: 111-112.
- Ando, T., S. Yoshida, S. Tatsuki and N. Takahashi, 1975. Lepidopterous sex attractants. *Agric. Biol. Chem.*, **39**: 1163-1165.
- Ando, T., S. Yoshida, S. Tatsuki and N. Takahashi, 1977. Sex attractants for male Lepidoptera. *Agric. Biol. Chem.*, **41**: 1485-1492.

- Ando, T., H. Kuroko, S. Nakagaki, O. Saito, T. Oku and N. Takahashi, 1978. Two-component sex attractants for male moths of the subfamily Tortricinae (Lepidoptera). *Agric. Biol. Chem.*, **42**: 1081-1083.
- Ando, T., T. Koshihara, H. Yamada, M. H. Vu, N. Takahashi and Y. Tamaki, 1979. Electroantennogram activities of sex pheromone analogues and their synergistic effect on field attraction in the diamondback moth. *Appl. Entomol. Zool.*, **14**: 362-364.
- Ando, T., K. Kishino, S. Tatsuki and N. Takahashi, 1980(a). Sex pheromone of the rice green caterpillar: chemical identification of three components and field tests with synthetic pheromones. *Agric. Biol. Chem.*, **44**: 765-775.
- Ando, T., O. Saito, K. Arai and N. Takahashi, 1980(b). (*Z*)- and (*E*)-12-Tetradecenyl acetates: sex pheromone components of oriental corn borer (Lepidoptera: Pyralidae). *Agric. Biol. Chem.*, **44**: 2643-2649.
- Ando, T., H. Kuroko, S. Nakagaki, O. Saito, T. Oku and N. Takahashi, 1981. Multi-component sex attractants in systematic field tests of male Lepidoptera. *Agric. Biol. Chem.*, **45**: 487-495.
- Ando, T., M. H. Vu, S. Yoshida, N. Takahashi, S. Tatsuki, K. Katagiri, A. Yamane, T. Ikeda and S. Yamazaki, 1982. (*5Z,7E*)-5,7-Dodecadien-1-ol: female sex pheromone of the pine moth *Dendrolimus spectabilis* Butler. *Agric. Biol. Chem.*, **46**: 709-715.
- Ando, T., K. Taguchi, M. Uchiyama and T. Horikawa, 1985(a). Female sex pheromone of the tea leafroller, *Caloptilia theivora* Walsingham (Lepidoptera: Gracillariidae). *Agric. Biol. Chem.*, **49**: 233-234.
- Ando, T., K. Taguchi, M. Uchiyama, T. Ujiye and H. Kuroko, 1985(b). (*7Z,11Z*)-7,11-Hexadecadienal: sex attractant of the citrus leafminer moth, *Phyllocnistis citrella* Stainton (Lepidoptera, Phyllocnistidae). *Agric. Biol. Chem.*, **49**: 3633-3635.
- Ando, T., M. Koike, M. Uchiyama and H. Kuroko, 1987. Lepidopterous sex attractants with a conjugated diene system. *Agric. Biol. Chem.*, **51**: 2691-2694.
- Ando, T., H. Ohsawa, T. Ueno, H. Kishi, Y. Okamura and S. Hashimoto, 1993. Hydrocarbons with a homoconjugated polyene system and their monoepoxy derivatives: sex attractants of geometrid and noctuid moths distributed in Japan. *J. Chem. Ecol.*, **19**: 787-798.

- Ando, T., H. Kishi, N. Akashio, X.-R. Qin, N. Saito, H. Abe and S. Hashimoto, 1995(a). Sex attractants of geometrid and noctuid moths: chemical characterization and field test of monoepoxides of 6,9-dienes and related compounds. *J. Chem. Ecol.*, **21**: 299-311.
- Ando, T., R. Ohno, T. Kokuryu, A. Funayoshi and M. Nomura, 1995(b). Identification of female sex pheromone components of rice looper, *Plusia festucae* (L.), (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **21**: 1181-1190.
- Ando, T., K. Ohtani, M. Yamamoto, T. Miyamoto, X.-R. Qin and Witjaksono, 1997. Sex pheromone of Japanese giant looper, *Ascotis selenaria cretacea*: identification and field tests. *J. Chem. Ecol.*, **23**: 2413-2423.
- Ando, T., S. Inomata, R. Shimada, M. Nomura, S. Uehara and G.-Q. Pu, 1998. Sex pheromones of *Thysanoplusia intermixta* and *T. orichalcea*: identification and field tests. *J. Chem. Ecol.*, **24**: 1105-1116.
- Anglade, P., J. Stockel and I. W. G. O. cooperators, 1984. Intraspecific sex-pheromone variability in the European corn borer, *Ostrinia nubilalis* Hbn. (Lepidoptera, Pyralidae). *Agronomie*, **4**: 183-187.
- Anshelevich, L., M. Kehat, E. Dunkelblum and S. Greenberg, 1993. Sex pheromone traps for monitoring the honeydew moth, *Cryptoblabes gnidiella*: effect of pheromone components, pheromone dose, field aging of dispenser, and type of trap on male captures. *Phytoparasitica*, **21**: 189-198.
- Arai, K., T. Ando, A. Sakurai, H. Yamada, T. Koshihara and N. Takahashi, 1982. Identification of the female sex pheromone of the cabbage webworm. *Agric. Biol. Chem.*, **46**: 2395-2397.
- Arakawa, A., F. Mochizuki, Y. Yumoto and K. Fukumoto, 2009. Sex pheromone components of *Toleria romanovi* and insect attractant containing octadecadienol and octadecadienyl acetate. *Jpn. Kokai Tokkyo Koho* JP 2009209104. [*SciFinder*. AN 2009:1131520]
- Aranda-Arguello, R., J. C. Rojas, E. A. Malo, G. López-Guillén and L. Cruz-López, 2022. Male attraction of *Gymnandrosoma aurantianum* (Lepidoptera: Tortricidae), from Guatemala, to its sex pheromone major component is not affected by the addition of secondary components. *Can. Entomol.*, **154**: e51.
- Arn, H., C. Schwarz, H. Limacher and E. Mani, 1974. Sex attractant inhibitors of the codling moth *Laspeyresia pomonella* L. *Experientia*, **30**: 1142-1144.
- Arn, H., E. Städler and S. Rauscher, 1975. The electroantennographic

- detector - a selective and sensitive tool in the gas chromatographic analysis of insect pheromones. *Z. Naturforsch.*, **30c**: 722-725.
- Arn, H., B. Delley, M. Baggiolini and P. J. Charmillot, 1976(a). Communication disruption with sex attractant for control of the plum fruit moth, *Grapholitha funebrana*: a two-year field study. *Entomol. Exp. Appl.*, **19**: 139-147.
- Arn, H., S. Rauscher, H.-R. Buser and W. L. Roelofs, 1976(b). Sex pheromone of *Eupoecilia ambiguella*: *cis*-9-dodecenyl acetate as a major component. *Z. Naturforsch.*, **31c**: 499-503.
- Arn, H., E. Priesner, H. Bogenschütz, H. R. Buser, D. L. Struble, S. Rauscher and S. Voerman, 1979. Sex pheromone of *Tortrix viridana*: (Z)-11-tetradecenyl acetate as the main component. *Z. Naturforsch.*, **34c**: 1281-1284.
- Arn, H., E. Städler, S. Rauscher, H. R. Buser, H. Mustaparta, P. Esbjerg, H. Philipsen, O. Zethner, D. L. Struble and R. Bues, 1980. Multicomponent sex pheromone in *Agrotis segetum*: preliminary analysis and field evaluation. *Z. Naturforsch.*, **35c**: 986-989.
- Arn, H., P. Esbjerg, R. Bues, M. Tóth, G. Szöcs, P. Guerin and S. Rauscher, 1983. Field attraction of *Agrotis segetum* males in four European countries to mixtures containing three homologous acetates. *J. Chem. Ecol.*, **9**: 267-276.
- Arn, H., P. M. Guerin, H. R. Buser, S. Rauscher and E. Mani, 1985. Sex pheromone blend of the codling moth, *Cydia pomonella*: evidence for a behavioral role of dodecan-1-ol. *Experientia*, **41**: 1482-1484.
- Arn, H., M. Tóth and E. Priesner, 1986(a). "List of sex pheromones of Lepidoptera and related attractants" OILB-SROP / IOBC-WPRS, Paris, pp. 123.
- Arn, H., S. Rauscher, H.-R. Buser and P. M. Guerin, 1986(b). Sex pheromone of *Eupoecilia ambiguella* female: analysis and male response to ternary blend. *J. Chem. Ecol.*, **12**: 1417-1429.
- Arn, H., S. Rauscher, P. Guerin and H.-R. Buser, 1988. Sex pheromone blends of three tortricid pests in European vineyards. *Agric. Ecosys. Environ.*, **21**: 111-117.
- Arn, H., M. Tóth and E. Priesner, 1992. "List of sex pheromones of Lepidoptera and related attractants" OILB-SROP / IOBC-WPRS, Secrétariat Général, F-84143 Montfavet, France, ISBN 92-9067-044-4. 179 pp.
- Arn, H., M. Tóth and E. Priesner, 1997. "List of sex pheromones of

Lepidoptera and related attractants, supplement 1992-1996"
Techno. Trans. Mating Disrup., **20** : 257-293.

Asaro, C., M. J. Dalusky and C. W. Berisford, 2001. Quantity and ratio of pheromone components among multiple generations of the nantucket pine tip moth (Lepidoptera: Tortricidae) in Georgia and Virginia. *Environ. Entomol.*, **30**: 1006-1011.

Arsura, E., A. Capizzi, P. Piccardi and P. Spinelli, 1977. (Z)-9-Tetradecen-1-ol and (Z)-9-tetradecenyl acetate: a potent attractant system for male *Sesamia cretica* Led. (Lep., Noctuidae). *Experientia*, **33**: 1423-1424.

Attygalle, A. B., J. Schwarz, O. Vostrowsky and H. J. Bestmann, 1986. Individual variation in the sex pheromone components of the false codling moth, *Cryptophlebia leucotreta* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **41c**: 1077-1081.

Attygalle, A. B., M. Herrig, O. Vostrowsky and H. J. Bestmann, 1987. Technique for injecting intact glands for analysis of sex pheromones of Lepidoptera by capillary gas chromatography: reinvestigation of pheromone complex of *Mamestra brassicae*. *J. Chem. Ecol.*, **13**: 1299-1311.

Attygalle, A. B., J. Schwarz and N. E. Gunawardena, 1988(a). Sex pheromone of brinjal shoot and pod borer *Leucinodis orbonalis* Guenée (Lepidoptera: Pyralidae: Pyraustinae). *Z. Naturforsch.*, **43c**: 790-792.

Attygalle, A. B., C.-H. Wu, J. Schwarz, O. Vostrowsky, I. Hasenfuss and H. J. Bestmann, 1988(b). Sex pheromone of female *Myelois cribrella* Hübner (Lepidoptera: Pyralidae). Chemical identification, electrophysiological evaluation, and field attractancy tests. *J. Chem. Ecol.*, **14**: 485-494.

Attygalle, A. B., G. N. Jham, A. Svatoš, R. T. S. Frighetto and J. Meinwald, 1995. Microscale, random reduction to the characterization of (3E,8Z,11Z)-3,8,11-tetradecatrienyl acetate, a new lepidopteran sex pheromone. *Tetrahedron Lett.*, **36**: 5471-5474.

Attygalle, A. B., G. N. Jham, A. Svatoš, R. T. S. Frighetto, F. A. Ferrara, E. F. Vilela, M. A. Uchoa-Fernandes and J. Meinwald, 1996. (3E,8Z,11Z)-3,8,11-Tetradecatrienyl acetate, major sex pheromone component of the tomato pest *Scrobipalpuloides absoluta* (Lepidoptera: Gelechiidae). *Bioorg. Medic. Chem.*, **4**: 305-314.

Avand-Faghih, A. and B. Frérot, 2008. Identification of the sex pheromone of *Sesamia cretica* Lederer. *J. Chem. Ecol.*, **34**:

103-106.

- Ayre, G. L., W. J. Turnock and D. L. Struble, 1982(a). Spatial and temporal variability in sex attractant trap catches of *Scotogramma trifolii* (Lepidoptera: Noctuidae) in relation to its biology in Manitoba. *Can. Entomol.*, **114**: 145-154.
- Ayre, G. L., W. J. Turnock and D. L. Struble, 1982(b). Spatial and temporal variability in sex attractant trap catches of *Euxoa messoria* and *E. ochrogaster* (Lepidoptera: Noctuidae) in relation to their biology in Manitoba. *Can. Entomol.*, **114**: 993-1001.
- Babu, V. R., S. Satpathy, B. V. S. Reddy, 2021. Identification and characterisation of female released sex pheromone components of jute semilooper, *Anomis sabulifera* Guenee (Lepidoptera: Noctuidae). *J. Environ. Biol.*, **42**: 254-264.
- Bacon, O. G., J. N. Seiber and G. G. Kennedy, 1976. Evaluation of survey trapping techniques for potato tuberworm moths with chemical baited traps. *J. Econ. Entomol.*, **69**: 569-572.
- Bailey, J. B., L. M. McDonough and M. P. Hoffmann, 1986. Western avocado leafroller, *Amorbia cuneana* (Walsingham), (Lepidoptera: Tortricidae). Discovery of populations utilizing different ratios of sex pheromone components. *J. Chem. Ecol.*, **12**: 1239-1245.
- Baker, J. L., A. S. Hill, R. T. Cardé, A. Kurokawa and W. L. Roelofs, 1975. Sex pheromone field trapping of the omnivorous leafroller, *Platynota stultana*. *Environ. Entomol.*, **4**: 90-92.
- Baker, R., J. W. S. Bradshaw and W. Speed, 1982. Methoxymercuration-demercuration and mass spectrometry in the identification of the sex pheromones of *Panolis flammea*, the pine beauty moth. *Experientia*, **38**: 233-234.
- Baker, R., R. H. Herbert, N. N. Neequaye and K. N. Rao, 1985. Sex pheromone of tobacco stem borer, *Scrobipalpa heliopa* (Lower) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **11**: 989-998.
- Baker, T. C., W. Francke, C. Löfstedt, B. S. Hansson, J.-W. Du, P. L. Phelan, R. S. Vetter and R. Youngman, 1989. Isolation, identification and synthesis of sex pheromone components of the carob moth, *Ectomyelois ceratoniae*. *Tetrahedron Lett.*, **30**: 2901-2902.
- Baker, T. C., W. Francke, J. G. Millar, C. Löfstedt, B. Hansson, J.-W. Du, P. L. Phelan, R. S. Vetter, R. Youngman and J. L. Todd, 1991. Identification and bioassay of sex pheromone components of carob moth, *Ectomyelois ceratoniae* (Zeller). *J. Chem. Ecol.*, **17**: 1973-1988.

- Baltensweiler, W., E. Priesner, H. Arn and V. Delucchi, 1978. Evidence for different sexual attractants used by the larch form and the cembran pine form of the grey larch bud moth (*Zeiraphera diniana* Gn.). *Mitt. Schweiz. Entomol. Ges.*, **51**: 133-142.
- Baltensweiler, W. and E. Priesner, 1988. A study of pheromone polymorphism in *Zeiraphera diniana* Gn. (Lep., Tortricidae). 3. Specificity of attraction to synthetic pheromone source by different male response types from two host races. *J. Appl. Entomol.*, **106**: 217-231.
- Barrer, P. M., M. J. Lacey and A. Shani, 1987. Variation in relative quantities of airborne sex pheromone components from individual female *Ephestia cautella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **13**: 639-653.
- Barros-Parada, W., J. Bergmann, T. Curkovic, C. Espinosa, E. Fuentes-Contreras, J. Guajardo, H. Herrera, S. Morales, A. F. O. Queiroz and A. Vidal, 2020. 3,7-Dimethylpentadecane: a novel sex pheromone component from *Leucoptera sinuella* (Lepidoptera: Lyonetiidae). *J. Chem. Ecol.*, **46**: 820-829.
- Barry, M. W., D. G. Nielsen, F. F. Purrington and J. H. Tumlinson, 1978. Attractivity of pheromone blends to male peachtree borer, *Synanthedon exitiosa*. *Environ. Entomol.*, **7**: 1-3.
- Bartell, R. J., T. E. Bellas and C. P. Whittle, 1988. Evidence for biological activity of two further alcohols in the sex pheromone of female *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). *J. Aust. Entomol. Soc.*, **27**: 11-12.
- Bathon, H. and M. Glas, 1983. The geographical distribution of the cereal leaf roller, *Cnephasia pumicana* Zeller (Lepidoptera: Tortricidae), in the Federal Republic of Germany. Preliminary results of a pheromone trap survey. *Nachrichtenbl. Deut. Pflanzenschutzd.*, **35**: 81-86.
- Batista-Pereira, L. G., E. M. Santangelo, K. Stein, C. R. Unelius, A. E. Eiras and A. G. Correa, 2002. Electrophysiological studies and identification of possible sex pheromone components of Brazilian populations of the sugarcane borer, *Diatraea saccharalis*. *Z. Naturforsch.*, **57c**: 753-758.
- Batista-Pereira, L. G., K. Stein, A. F. de Paula, J. A. Moreira, I. Cruz, M. de L. C. Figueiredo, J. Perri Jr. and A. G. Correa, 2006. Isolation, identification, synthesis, and field evaluation of the sex pheromone of the Brazilian population of *Spodoptera frugiperda*. *J. Chem. Ecol.*, **32**(5): 1085-1099.

- Bavaresco, A., S. Nunez, M. S. Garcia, M. Botton and J. Sant'Ana, 2005. Attraction of males of the South American tortricid moth *Argyrotaenia sphaleropa* (Meyrick) (Lepidoptera: Tortricidae) to the components of the synthetic sexual pheromone in persimmon. *Neotrop. Entomol.*, **34**: 619-625.
- Beauvais, F., G. Biwer and P. J. Charmillot, 1977. Tests of sex trapping of *Grapholitha janthinana* Dup. with a mixture of Z-8 dodecenyl acetate containing 28 p. 100 of its isomer E. *Ann. Zool. Ecol. Anim.*, **9**: 457-467.
- Becker, D., T. Kimmel, R. Cyjon, I. Moore, M. Wysoki, H. J. Bestmann, H. Platz, K. Roth and O. Vostrowsky, 1983. (3Z,6Z,9Z)-3,6,9-Nonadecatriene -- a component of the sex pheromonal system of the giant looper, *Boarmia* (*Ascotis*) *selenaria* Schiffermüller (Lepidoptera: Geometridae). *Tetrahedron Lett.*, **24**: 5505-5508.
- Becker, D., R. Cyjon, A. Cossé, I. Moore, T. Kimmel and M. Wysoki, 1990. Identification and enantioselective synthesis of (Z,Z)-6,9-cis-3S,4R-epoxynonadecadiene, the major sex pheromone component of *Boarmia selenaria*. *Tetrahedron Lett.*, **31**: 4923-4926.
- Bédard, C., R. Gries, G. Gries and R. Bennett, 2002. Sex pheromone and diel periodicity of *Cydia strobilella* (Lepidoptera: Tortricidae) pheromonal communication. *Can. Entomol.*, **134**: 847-850.
- Beevor, P. S., D. R. Hall, R. Lester, R. G. Poppi, J. S. Read and B. F. Nesbitt, 1975. Sex pheromones of the armyworm moth, *Spodoptera exempta* (Wlk.). *Experientia*, **31**: 22-23.
- Beevor, P. S., D. R. Hall, B. F. Nesbitt, V. A. Dyck, G. Arida, P. C. Lippold and H. Oloumi-Sadeghi, 1977. Field trials of the synthetic sex pheromones of the stripe rice borer, *Chilo suppressalis* (Walker) (Lepidoptera: Pyralidae), and of related compounds. *Bull. Entomol. Res.*, **67**: 439-447.
- Beevor, P. S., A. Cork, D. R. Hall, B. F. Nesbitt, R. K. Day and J. D. Mumford, 1986. Components of female sex pheromone of cocoa pod borer moth, *Conopomorpha cramerella*. *J. Chem. Ecol.*, **12**: 1-23.
- Beevor, P. S., O. Youm, D. R. Hall and A. Cork, 1999. Identification and field evaluation of components of female sex pheromone of millet stem borer, *Coniesta ignefusalis*. *J. Chem. Ecol.*, **25**(12): 2643-2663.
- Bell, T. W. and J. Meinwald, 1986. Pheromones of two arctiid moths

- (*Cretonotos transiens* and *C. gangis*): chiral components from both sexes and achiral female components. *J. Chem. Ecol.*, **12**: 385-409.
- Bellas, T. E., R. J. Bartell and A. Hill, 1983. Identification of two components of the sex pheromone of the moth, *Epiphyas postvittana* (Lepidoptera, Tortricidae). *J. Chem. Ecol.*, **9**: 503-512.
- Bengtsson, M., St. Rauscher, H. Arn, W.-C. Sun and G. D. Prestwich, 1990. Fluorine-substituted pheromone components affect the behavior of the grape berry moth. *Experientia*, **46**: 1211- 1213.
- Benn, M. H., R. A. Galbreath, V. A. Holt and H. Young, 1981. A sex attractant for *Sperchia intractana* Walker (Lepidoptera: Tortricidae) in New Zealand. *Z. Naturforsch.*, **36 c**: 178-179.
- Benn, M. H., R. A. Galbreath, V. A. Holt, H. Young, G. Down and E. Priesner, 1982. The sex pheromone of the silver Y moth *Chrysodeixis eriosoma* (Doubleday) in New Zealand. *Z. Naturforsch.*, **37 c**: 1130-1135.
- Benz, G. and G. von Salis, 1973. Use of synthetic sex attractant of larch bud moth *Zeiraphera diniana* (Gn.) in monitoring traps under different conditions, and antagonistic action of cis-isomer. *Experientia*, **29**: 729-730.
- Berger, R. S., 1966. Isolation, identification, and synthesis of the sex attractant of the cabbage looper, *Trichoplusia ni*. *Ann. Entomol. Soc. Am.*, **59**: 767-771.
- Berger, R. S. and T. D. Canerday, 1968. Specificity of the cabbage looper sex attractant. *J. Econ. Entomol.*, **61**: 452-454.
- Bergmann, J., L. Reyes-Garcia, C. Ballesteros, Y. Cuevae, M. F. Flores, and T. Curkovic, 2016. Identification of the female sexpheromone of the leafroller *Proeulia triquetra* Obraztsov (Lepidoptera: Tortricidae). *Neotrop. Entomol.*, **45**: 351-356.
- Beroza, M., K. Katagiri, Z. Iwata, H. Ishizuka, S. Suzuki and B. A. Bierl, 1973(a). Disparlure and analogues as attractants for two Japanese lymantriid moths. *Environ. Entomol.*, **2**: 966.
- Beroza, M., A. A. Punjabi and B. A. Bierl, 1973(b). Disparlure and analogues as attractants for *Lymantria obfuscata*. *J. Econ. Entomol.*, **66**: 1215-1216.
- Beroza, M., C. R. Gentry, J. L. Blythe and G. M. Muschik, 1973(c). Isomer content and other factors influencing captures of oriental fruit moth by synthetic pheromone traps. *J. Econ. Entomol.*, **66**:

1307-1311.

Beroza, M., B. A. Bierl and H. R. Moffitt, 1974. Sex pheromones: (E,E)-8,10-dodecadien-1-ol in the codling moth. *Science*, **183**: 89-90.

Bestmann, H. J., O. Vostrowsky, K.-H. Koschatzky, H. Platz, T. Brosche, I. Kantardjiew, M. Rheinwald and W. Knauf, 1978(a). (Z)-5-Decenyl acetate, a sex attractant for the male turnip moth *Agrotis segetum* (Lepidoptera). *Angew. Chem. Int. Ed. Engl.*, **17**: 768.

Bestmann, H. J., O. Vostrowsky, K. H. Koschatzky, H. Platz, A. Szymanska and W. Knauf, 1978(b). Pheromone XVII. (Z)-11-Hexadecenylacetat, ein Sexuallockstoff des Pheromonsystems der Kohleule *Mamestra brassicae*. *Tetrahedron Lett.*, 605-608.

Bestmann, O. Vostrowsky, H. Platz, T. Brosche, K. H. Koschatzky, 1979. Pheromone XXIII. (Z)-7-Tetradecenylacetat, ein Sexuallockstoff für Männchen von *Amnathes c-nigrum* (Noctuidae, Lepidoptera). *Tetrahedron Lett.*, 497-500.

Bestmann, H. J., T. Brosche, K. H. Koschatzky, K. Michaelis, H. Platz, O. Vostrowsky and W. Knauf, 1980. Pheromone XXX. Identifizierung eines neuartigen Pheromonkomplexes aus der Graseule *Scotia exclamationis* (Lepidoptera). *Tetrahedron Lett.*, **21**: 747-750.

Bestmann, H. J. and O. Vostrowsky, 1982(a). Insect pheromone. *Naturwissenschaften*, **69**: 457-471.

Bestmann, H. J., K. H. Koschatzky, H. Platz, J. Süß O. Vostrowsky, W. Knauf, G. Burghardt and I. Schneider, 1982(b). Pheromones, XL. Synthesis of the pheromone complex of Lasiocampidae species (Lepidoptera); a sex attractant for *Dendrolimus pini*. *Liebigs Ann. Chem.*, 1359-1365.

Bestmann, H. J., T. Brosche, K. H. Koschatzky, K. Michaelis, H. Platz, K. Roth, J. Süß O. Vostrowsky and W. Knauf, 1982(c). Pheromone-XLII. 1,3,6,9-Nonadecatetraen, das Sexualpheromon des Frostspanners *Operophtera brumata* (Geometridae). *Tetrahedron Lett.*, **23**: 4007-4010.

Bestmann, H. J., K. H. Koschatzky and O. Vostrowsky, 1982(d). Stereoselective syntheses of 1,5- and 1,6-alkadienes as sex pheromones and pheromone analogues. *Liebigs Ann. Chem.*, 1478-1494.

Bestmann, H.-J., A. Attygalle, H. Platz and O. Vostrowsky, 1985. (Z)-9-Dodecenyl acetate, a component of the sex pheromone of *Cnephasia longana* Haworth (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **40c**: 667-669.

- Bestmann, H.-J., A. B. Attygalle, T. Brosche, J. Erler, H. Platz, J. Schwarz, O. Vostrowsky, C.-H. Wu, K. E. Kaissling and T.-M. Chen, 1987. Identification of three sex pheromone components of the female Saturniid moth *Antheraea pernyi* (Lepidoptera: Saturniidae). *Z. Naturforsch.*, **42c**, 631-636.
- Bestmann, H. J., A. B. Attygalle, J. Schwarz, O. Vostrowsky and W. Knauf, 1988(a). Identification of sex pheromone components of *Spodoptera sunia* Guenée (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **14**: 683-690.
- Bestmann, H. J., J. Erler and O. Vostrowsky, 1988(b). Determination of diel periodicity of sex pheromone release in three species of Lepidoptera by 'closed-loop-stripping'. *Experientia*, **44**: 797-799.
- Bestmann, H. J., A. B. Attygalle, J. Schwarz, W. Garbe, O. Vostrowsky and I. Tomida, 1989. Pheromones, 71. Identification and synthesis of female sex pheromone of eri-silkworm, *Samia cynthia ricini* (Lepidoptera: Saturniidae). *Tetrahedron Lett.*, **30**: 2911-2914.
- Bestmann, H. J., F. Kern, D. Schäfer, O. Vostrowsky and I. Hasenfuss, 1991. (11Z,13Z)-11,13-Hexadecadienyl acetate, the sex pheromone of females of *Notodonta torva*. *Naturwissenschaften*, **78**: 465-467.
- Bestmann, H. J., F. Kern, A. Mineif, H. Platz and O. Vostrowsky, 1992(a). Pheromone, 84. The sex pheromone complex of the arctiid moth *Arctia caja* (Lepidoptera: Arctiidae). *Z. Naturforsch.*, **47c**: 132-135.
- Bestmann, H. J., J. Erler, W. Garbe, F. Kern, V. Martischonok, D. Schäfer, O. Vostrowsky and L. T. Wasserthal, 1992(b). Pheromone components of the female elephant hawk-moth, *Deilephila elpenor*, and the silver-striped hawk-moth, *Hippotion celerio*. *Experientia*, **48**: 610-613.
- Bestmann, H. J., F. Kern, D. Schäfer and O. Vostrowsky, 1993(a). (11Z,13Z)-11,13-Hexadecadienal, the sex pheromone of females of *Notodonta dromedarius*. *Naturwissenschaften*, **80**: 271-273.
- Bestmann, H. J., F. Kern, G. G. Melikyan, D. Schäfer, O. Vostrowsky, E. V. Babayan and Sh. O. Badanyan, 1993(b). Pheromone, 88. Sex pheromone components of female *Euzophera punicaella* M. (Lepidoptera, Pyralidae). *Z. Naturforsch.*, **48c**: 110-112.
- Bestmann, H. J., F. Kern, E. Janssen and I. Hasenfuss, 1993(c). Pheromone, 93. The sex pheromone of the cosmopterigid moth *Limnaecia phragmitella* (Lepidoptera: Cosmopterigidae). *Z. Naturforsch.*, **48c**: 515-518.

- Bestmann, H. J., A. B. Attygalle, W. Garbe, F. Kern, V. Martichonok, D. Schäfer, O. Vostrowsky and I. Hasenfub 1993(d). Chemical structure and final steps of biosynthesis of the female sex pheromone of *Gastropacha quercifolia* (Lepidoptera: Lasiocampidae). *Insect Biochem. Molec. Biol.*, 23: 791-799.
- Bestmann, H. J., E. Janssen, F. Kern, D. Schäfer and O. Vostrowsky, 1994. Pheromones, 97. The sex pheromone complex of the female arctiid moth *Thyria jacobaeae* (Lepidoptera, Arctiidae). *Z. Naturforsch.*, 49c: 276-279.
- Bhanu, K. R. M., D. R. Hall, R. V. Awalekar, M. Chandrashekharaiiah, T. N. Divya, M. S. Prabhakara and K. P. Jayanth, 2018. Identification and field evaluation of female sex pheromone of leaf-eating caterpillar, *Opisina arenosella* (Lepidoptera: Oecophoridae). *Int. J. Tropic. Insect Sci.*, 38: 274-282.
- Bhardwaj, S. P., 1987. Investigations on the response of lepidopteran sexpheromones of temperate fruitpests in Himachal Pradesh, India. *Agric. Ecosyst. Environ.*, 19: 87-91.
- Biddinger, D. J., W. Roelofs and A. J. Howitt, 1994. The development of a sex pheromone lure for the American plum borer, *Euzophera semifuneralis* (Lepidoptera: Pyralidae), a major pest of cherry in Michigan. *Great Lakes Entomol.*, 26: 311-317.
- Bierl, B. A., M. Beroza and C. W. Collier, 1970. Potent sex attractant of the gypsy moth: its isolation, identification, and synthesis. *Science*, 170: 87-89.
- Bierl, B. A., M. Beroza and C. W. Collier, 1972. Isolation, identification, and synthesis of the gypsy moth sex attractant. *J. Econ. Entomol.*, 65: 659-664.
- Bierl, B. A., M. Beroza, R. T. Staten, P. E. Sonnet and V. E. Adler, 1974. The pink bollworm sex attractant. *J. Econ. Entomol.*, 67: 211-216.
- Bierl, B. A., M. Beroza, V. E. Adler, G. Kasang, H. Schröter and D. Schneider, 1975. The presence of disparlure, the sex pheromone of the gypsy moth, in the female nun moth. *Z. Naturforsch.*, 30 c: 672-675.
- Birch, M. C., 1977. Response of both sexes of *Trichoplusia ni* (Lepidoptera: Noctuidae) to virgin females and to synthetic pheromone. *Ecol. Entomol.*, 2: 99-104.
- Biwer, G., B. Lalanne-Cassou, C. Descoins and D. Samain, 1975. Sex trapping of *Sterrhia biselata* (Lepidoptera, Geometridae,

- Sterrhinae) by (7E,9Z)-dodecadienyl acetate, a sex pheromone for *Lobesia botrana* (Lepidoptera, Tortricidae, Olethreutinae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **280**: 1469-1471.
- Biwer, G., C. Descoins, M. Gallois, E. Priesner, J. P. Chambon, G. Genestier and M. Martinez, 1978(a). Study of the sex pheromone blend produced by the virgin female of the cereal tortrix: *Cnephasia pumicana* Zeller. *Ann. Zool. Ecol. Anim.*, **10**: 129-138.
- Biwer, G. and C. Descoins, 1978(b). Approach to a mechanism of sexual isolation between four species of tortricids of the *Grapholitha* genus. *C. R. Hebd. Seances Acad. Sci., Ser. D*, **286**: 875-877.
- Biwer, G., C. Descoins and M. Gallois, 1979. Volatile components from the sex pheromone-producing gland of *Grapholitha molesta* Busk virgin females (Lepidoptera Tortricidae, Olethreutinae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **288**: 413-416.
- Bjostad, L. B., E. F. Taschenberg and W. L. Roelofs, 1980(a). Sex pheromone of the chokecherry leafroller moth, *Sparganothis directana*. *J. Chem. Ecol.*, **6**: 487-498.
- Bjostad, L. B., L. K. Gaston, L. L. Noble, J. H. Moyer and H. H. Shorey, 1980(b). Dodecyl acetate, a second pheromone component of the cabbage looper moth, *Trichoplusia ni*. *J. Chem. Ecol.*, **6**: 727-734.
- Bjostad, L. B., E. F. Taschenberg and W. L. Roelofs, 1980(c). Sex pheromone of the woodbine leafroller moth, *Sparganothis* sp. *J. Chem. Ecol.*, **6**: 797-804.
- Bjostad, L. B., E. Gurevitz, S. Gothilf and W. L. Roelofs, 1981. Sex attractant for the honeydew moth, *Cryptoblabes gnidiella*. *Phytoparasitica*, **9**: 95-99.
- Bjostad, L. B., C. E. Linn, J.-W. Du and W. L. Roelofs, 1984. Identification of new sex pheromone components in *Trichoplusia ni*, predicted from biosynthetic precursors. *J. Chem. Ecol.*, **10**: 1309-1323.
- Blair, B. W. and J. Tannock, 1978. A further note on the possible pheromone for *Spodoptera trituratora* (Walker) (Lepidoptera, Noctuidae). *Rhod. J. Agric. Res.*, **16**: 221-223.
- Blassioli-Moraes, M. C., M. Borges, A. R. Viana, R. A. Laumann, J. E. Miranda, D. M. Magalhães and M. A. Birkett, 2016. Identification and field evaluation of the sex pheromone of a Brazilian population of *Spodoptera cosmioides*. *Pesq. Agropec. Bras.*, **51**: 545-554.
- Bode, W. M., D. Asquith and J. P. Tette, 1973. Sex attractants and

- traps for tufted apple budmoth and redbanded leafroller males. *J. Econ. Entomol.*, **66**: 1129-1130.
- Bogenschütz, von H., W. Knauf, E. J. Tröger, H. J. Bestmann and O. Vostrowsky, 1985. Pheromones 49: males of Geometridae caught by C₁₉-polyenes in pine stands. *Z. Angew. Entomol.*, **100**: 349-354.
- Boness, M., 1978. Experiences with sex pheromone of Lepidoptera. *Anz. Schädlingsskde.*, **51**: 161-166.
- Boo, K. S. and C. H. Jung 1998(a). Field tests of synthetic sex pheromone of the apple leafminer moth, *Phyllonorycter ringoniella*. *J. Chem. Ecol.*, **24**(12): 1939-1947.
- Boo, K. S. and J. W. Park, 1998(b). Sex pheromone composition of the Asian cornborer moth, *Ostrinia furnacalis* (Guenée) (Lepidoptera: Pyralidae) in South Korea. *J. Asia Pacific Entomol.*, **1**: 77-84.
- Booiij, C. J. H. and S. Voerman, 1984(a). New sex attractants for 35 tortricid and 4 other lepidopterous species, found by systematic field screening in the Netherlands. *J. Chem. Ecol.*, **10**: 135-144.
- Booiij, C. J. H. and S. Voerman, 1984(b). Sex attractants for the fruit-attacking noctuids *Orthosia incerta* and *Orthosia cruda*. *Entomol. Exp. Appl.*, **36**: 15-16.
- Booiij, C. J. H. and S. Voerman, 1984(c). (Z)-11-Hexadecenyl compounds as attractants for male microlepidoptera of the subfamilies Argyresthiinae, Glyphipteryginae, and Crambinae. *Entomol. Exp. Appl.*, **36**: 47-53.
- Booiij, C. J. H., 1984(d). *Cydia indivisa* (Danilevski), een nieuwe bladrollersoort voor de Nederlandse fauna, gevangen met sekslokstoffen (Lepidoptera: Tortricidae). *Entomol. Ber.*, **44**: 133-135.
- Booiij, C. J. H. and S. Voerman, 1984(e). Sex attractant for the poplar shoot-borer, *Gypsonoma aceriana* (Duponchel). *Z. Angew. Entomol.*, **97**: 176-179.
- Booiij, C. J. H. and S. Voerman, 1985. New lepidopteran sex attractants found by systematic field screening of blends containing (Z)-11- and (E)-11-tetradecenal. *J. Chem. Ecol.*, **11**: 1333-1339.
- Booiij, C. J. H., S. Voerman and L. P. M. Willemse, 1986. Conifer-infesting tortricids in Western Europe: a search for more potent and selective sex attractants. *J. Appl. Entomol.*, **102**: 253-259.
- Borek, V., B. Kalinová, I. Valterová, R. Hochmut and J. Vrkoc~, 1991.

- Sex pheromone gland volatiles from *Hypsipyla grandella* females (Lepidoptera, Pyralidae, Phycitinae). *Acta. Entomol. Bochemoslov.*, **88**: 181-186.
- Borg-Karlson, A.-K. and R. Mozūraitis, 1996. Solid phase micro extraction technique used for collecting semiochemicals. Identification of volatiles released by individual signalling *Phyllonorycter sylvella* moths. *Z. Naturforsch.*, **51c**: 599-602. (See Ph-3-2-6)
- Bosa, C. F., A. M. C. Prado, T. Fukumoto, M. Bengtsson and P. Witzgall, 2005. Pheromone-mediated communication disruption in Guatemalan potato moth, *Tecia solanivora*. *Entomol. Exp. Appl.*, **114**: 137-142.
- Bosson, G. A. and M. Gallois, 1982. Analysis of pheromonal secretion in virgin females of *Hypsipyla robusta* (Moore) (Lepidoptera, Pyralidae, Phycitinae) from young mathogany shoots. *C. R. Acad. Sci., Ser. 3*, **294**: 819-822.
- Bourdouxhe, L. 1982. Results of two years of sex trapping of *Cryptophlebia leucotreta* Meyr. in Senegal. *FAO Plant Prot. Bull.*, **30**: 125-129.
- Bouwer, M. C., B. Slippers, D. Degefu, M. J. Wingfield, S. Lawson and E. R. Rohwer, 2015. Identification of the sex pheromone of the tree infesting cossid moth *Coryphodema tristis* (Lepidoptera: Cossidae). *PloS ONE.*, **10**: e0118575.
- Bouwer, M. C., B. Slippers, M. J. Wingfield, J. D. Allison and E. R. Rohwer, 2017. Optimization of pheromone traps for *Coryphodema tristis* (Lepidoptera: Cossidae). *J. Econ. Entomol.*, **110**: 1603-1610.
- Bouwer, M. C., L.-M. Scheepers, B. Slippers, E. R. Rohwer and J. D. Allison, 2021. The sex pheromone of the pine brown-tail moth, *Euproctis terminalis* (Lepidoptera: Erebididae). *J. Chem. Ecol.*, **47**: 732-739.
- Bradshaw, J. W. S., R. Baker, C. Longhurst, J. C. Edwards and J. C. Lisk, 1983. Optimization of a monitoring system for the pine beauty moth, *Panolis flammea* (Denis & Schiffermüller), using sex attractants. *Crop Prot.*, **2**: 63-73.
- Brady, U. E., J. H. Tumlinson III, R. G. Brownlee and R. M. Silverstein, 1971(a). Sex stimulant and attractant in the Indian meal moth and in the almond moth. *Science*, **171**: 802-804.
- Brady, U. E. and D. A. Nordlund, 1971(b). *Cis*-9, *trans*-12 tetradecadien-1-yl acetate in the female tobacco moth *Ephestia elutella* (Hübner) and evidence for an additional component of the sex pheromone. *Life Sci.*, **10**: 797-801.

- Brady, U. E. and M. C. Ganyard, Jr., 1972(a). Identification of a sex pheromone of the female beet armyworm, *Spodoptera exigua*. *Ann. Entomol. Soc. Am.*, **65**: 898-899.
- Brady, U. E. and R. C. Daley, 1972(b). Identification of a sex pheromone from the female raisin moth, *Cadra figulilella*. *Ann. Entomol. Soc. Am.*, **65**: 1356-1358.
- Brady, U. E., 1973. Isolation, identification and stimulatory activity of a second component of the sex pheromone system (complex) of the female almond moth, *Cadra cautella* (Walker). *Life Sci.*, **13**: 227-235.
- Brennan, B. M., 1977. Attraction of *Ereunetis simulans* (Lepidoptera: Tineidae) to the sex attractant of peachtree borer, *Sanninoidae exitiosa* (Lepidoptera: Sesiidae). *Proc. Hawaii. Entomol. Soc.*, **22**: 457-460.
- Breuer, M., H.-G. Kontzog, A. Guerrero, F. Camps and A. de Loof, 2003. Field trials with the synthetic sex pheromone of the oak processionary moth *Thaumetopoea processionea*. *J. Chem. Ecol.*, **29**(11): 2461-2468.
- Brewer, J. W., F. Krampfl and V. Skuhavy', 1985. Forest Lepidoptera attracted to six synthetic pheromones in Czechoslovakia. *Z. Angew. Entomol.*, **100**: 372-381.
- Brown, L. N. and J. W. Snow, 1985(a). The blackberry clearwing borer, *Pennisetia marginata* (Harris): a first report in Florida. *Florida Entomol.*, **68**: 699-700.
- Brown, L. N., 1985(b). New records of the rare clearwing moth, *Alcathoe carolinensis* Engelhardt, (Sesiidae) in Florida. *Florida Entomol.*, **68**: 700-701.
- Brown, L. N., T. D. Eichlin and J. W. Snow, 1985(c). A new species of clearwing moth, *Carmenta laurelae* (Sesiidae), from Florida. *J. Lepido. Soc.*, **39**: 262-265.
- Brown, L. N., T. D. Eichlin and J. W. Snow, 1985(d). Ecological notes on *Synanthedon dominicki* Duckworth and Eichlin (Sesiidae) in Florida and first description of the female. *J. Lepido. Soc.*, **39**: 196-200.
- Brown, L. N. and J. W. Snow, 1986(a). First record of the clearwing moth, *Carmenta odda* (Lepidoptera: Sesiidae) in Florida. *Florida Entomol.*, **69**: 423-424.
- Brown, L. N. 1986(b). First records of three clearwing moths

- (*Synanthedon tabaniformis*, *Synanthedon proxima*, and *Synanthedon castaneae*) in Florida. *Florida Scientist*, **49**: 80-81.
- Brückner, C., E. Beschmann, R. Becker and W. Seufert, 1988. A new highly effective synthetic pheromone mimic for *Lobesia botrana* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **43 c**: 315-318.
- Bûda, V., V. Karalius and P. Ivinskis, 1987. Sex attractants for *Pyrausta nigrilinealis* Hbn. (Lepidoptera, Pyralidae) and leafroller *Apotomis capreana* Hbn. (Lepidoptera, Tortricidae). *Khemoretsepsiya Nasekomykh*, **9**: 84-88.
- Bûda, V., U. Mäeorg, V. Karalius, G. H. L. Rothschild, S. Kolonistova, P. Ivinskis and R. Mozûraitis, 1993. C₁₈ Dienes as attractants for eighteen clearwing (Sesiidae), tineid (Tineidae), and choreutid (Choreutidae) moth species. *J. Chem. Ecol.*, **19**: 799-813.
- Buechi, R., E. Priesner and R. Brunetti, 1982. Sympatric occurrence of two pheromonal strains of the European corn borer, *Ostrinia nubilalis* Hbn., in southern Switzerland. *Mitt. Schweiz. Entomol. Ges.*, **55**: 33-53.
- Buleza, V. V., 1991. Role of sex pheromone components in male reproduction behavior of black cutworm moth *Agrotis ipsilon*. *Dokl. Akad. Nauk SSSR*, **318**: 1265-1268. [*Chem. Abst.*, **116**: 53588t]
- Burghardt, G., W. Knauf, H. J. Bestmann, O. Vostrowsky and K. H. Koschatzky, 1980. Synthetic pheromones and their inhibitors for *Choristoneura murinana* Hbn. (silver-fir shoot tortricid. Lepidoptera: Tortricidae). *Anz. Schädlingskde.*, **53**: 49-51.
- Burns, E. L. and P. E. A. Teal, 1989. Response of male potato stem borer moths, *Hydraecia micacea* (Esper) to conspecific females and synthetic pheromone blends in the laboratory and field. *J. Chem. Ecol.*, **15**: 1365-1378.
- Buser, H.-R., S. Rauscher and H. Arn, 1974. Sex pheromone of *Lobesia botrana*: (E,Z)-7,9-dodecadienyl acetate in the female grape vine moth. *Z. Naturforsch.*, **29 c**: 781-783.
- Buser, H.-R. and H. Arn, 1975. Analysis of insect pheromones by quadrupole mass fragmentography and high-resolution gas chromatography (*Laspeyresia pomonella*, *Lobesia botrana*). *J. Chromatogr.*, **106**: 83-95.
- Buser, H. R., P. M. Guerin, M. Tóth, G. Szöcs, A. Schmid, W. Francke and H. Arn, 1985. (Z,Z)-6,9-Nonadecadien-3-one and (Z,Z,Z)-3,6,9-nonadecatriene: identification and synthesis of sex pheromone components of *Peribatodes rhomboidaria*.

Tetrahedron, Lett., **26**: 403-406.

Butenandt, von A., R. Beckmann, D. Stamm and E. Hecker, 1959. Über den Sexual-Lockstoff des Seidenspinners *Bombyx mori*. Reindarstellung und Konstitution. *Z. Naturforsch.*, **14 b**: 283-284.

Butenandt, A., R. Beckmann and E. Hecker, 1961. Über den Sexuallockstoff des Seidenspinners, I. Der biologische Test und Isolierung des reinen Sexuallockstoffes Bombykol. *Hoppe-Seyler's Z. Physiol. Chem.*, **324**: 71-83.

Butler, L. I., J. E. Halfhill, L. M. McDonough and B. A. Butt, 1977. Sex attractant of the alfalfa looper *Autographa californica* and the celery looper *Anagrapha falcifera* (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **3**: 65-70.

Byers, J. R., E. W. Underhill, W. F. Steck, M. D. Chisholm and P. E. A. Teal, 1981. Biosystematics of the genus *Euxoa* (Lepidoptera: Noctuidae). XV. Sex pheromone cross attractancy among the three closely related species of the *declarata* group. *Can. Entomol.*, **113**: 235-243.

Byers, J. R. and D. L. Struble, 1987. Monitoring population levels of eight species of noctuids with sex-attractant traps in southern Alberta, 1978-1983: specificity of attractants and effect of target species abundance. *Can. Entomol.*, **119**: 541-556.

Byers, J. R. and D. L. Struble, 1990. Identification of sex pheromones of two sibling species in dingy cutworm complex, *Feltia jaculifera* (Gn.) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **16**: 2981-2992.

Byers, J. R. and C. E. Herle, 1997. Identification of sex pheromones of *Leucania anteoclarata* Sm. and *Leucania commoides* Gn. (Lepidoptera: Noctuidae; Hadeninae). *J. Chem. Ecol.*, **23**(2): 473-485.

Cabrera, A., A. Eiras, G. Gries, R. Gries, N. Urdaneta, B. Miras, C. Badji and K. Jaffe, 2001. Sex pheromone of tomato fruit borer, *Neoleucinodes elegantalis*. *J. Chem. Ecol.*, **27**(10): 2097-2107.

Campbell, C. A. M., E. L. Tregidga, D. R. Hall, T. Ando and M. Yamamoto, 2007. Components of the sex pheromone of the currant pug moth *Eupithecia assimidata*, a re-emergent hop pest in UK. *Entomol. Exp. Appl.*, **122**: 265-269.

Campion, D. G., B. W. Bettany, B. F. Nesbitt, P. S. Beevor, R. Lester and R. G. Poppi, 1974. Field studies of the female sex pheromone of the cotton leafworm *Spodoptera littoralis* (Boisd.) in Cyprus. *Bull. Entomol. Res.*, **64**: 89-96.

- Campion, D. G., 1975. Sex pheromones and their uses for control of insects of the genus *Spodoptera*. *Meded. Fac. Landbouww. Rijksuniv. Gent.*, **40**: 283-292.
- Campion, D. G., L. J. McVeigh, J. Polyrakis, S. Michaelakis, G. N. Stravarakis, P. S. Beevor, D. R. Hall and B. F. Nesbitt, 1979. Laboratory and field studies of the female sex pheromone of the olive moth, *Prays oleae*. *Experientia*, **35**: 1146-1147.
- Campion, D. G., P. Hunter-Jones, L. J. McVeigh, D. R. Hall, R. Lester and B. F. Nesbitt, 1980. Modification of the attractiveness of the primary pheromone component of the Ezyptian cotton leafworm, *Spodoptera littoralis* (Boisduval) (Lepidoptera: Noctuidae), by secondary pheromone components and related chemicals. *Bull. Entomol. Res.*, **70**: 417-434.
- Can, F., K. A. Efetov, J. Burman, K. Kaya, E. E. Kucherenko, B. Ulaşlı and G. M. Tarmann, 2019. A study of the Zygaenidae (Lepidoptera) fauna of Central Anatolia, Turkey. *Turk. J. Entomol.*, **43**: 189-199.
- Cao, W.-H. and H.-J. He, 2002. Identification of sex pheromone of the moth, *Tyta luctuosa*. *Acta Entomol. Sinica (Kunchong Xuebao)*, **45**: 30-34.
- Cao, W.-H., R. E. Charlton, J. R. Nechols and M. J. Horak, 2003. Sex pheromone of the noctuid moth, *Tyta luctuosa* (Lepidoptera: Noctuidae), a candidate biological control agent of field bindweed. *Environ. Entomol.*, **32**: 17-22.
- Capek, M., 1979. The influence of orientation of pheromone traps on their attractivity for gypsy moth (*Lymantria dispar* L.). *Lesnictivi*, **25**: 301-306.
- Capizzi, A., C. Tonini, E. Arsura, G. Guglielmetti, P. Massardo and P. Piccardi, 1983. Sex pheromone components of the European goat moth, *Cossus cossus*. *J. Chem. Ecol.*, **9**: 191-200.
- Cardé, R. T., J. Kochansky, J. F. Stimmel, A. G. Wheeler, Jr. and W. L. Roelofs, 1975(a). Sex pheromone of the European corn borer (*Ostrinia nubilalis*): *cis*- and *trans*-responding males in Pennsylvania. *Environ. Entomol.*, **4**: 413-414.
- Cardé, R. T., T. C. Baker and W. L. Roelofs, 1975(b). Behavioural role of individual components of a multichemical attractant system in the Oriental fruit moth. *Nature*, **253**: 348-349.
- Cardé, R. T., T. C. Baker and W. L. Roelofs, 1975(c). Ethological function of components of a sex attractant system for oriental

- fruitmothmales, *Grapholitha molesta* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **1**: 475-491.
- Cardé, R. T., A. M. Cardé, A. S. Hill and W. L. Roelofs, 1977(a). Sex pheromone specificity as a reproductive isolating mechanism among the sibling species *Archips argyrospilus* and *A. mortuanus* and other sympatric tortricine moths (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **3**: 71-84.
- Cardé, R. T., C. C. Doane, T. C. Baker, S. Iwaki and S. Marumo, 1977(b). Attractancy of optically active pheromone for male gypsy moths. *Environ. Entomol.*, **6**: 768-772.
- Cardé, A. M., T. C. Baker and R. T. Cardé, 1979. Identification of a four-component sex pheromone of the female Oriental fruit moth, *Grapholita molesta* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **5**: 423-427.
- Carney, R. L. and A. S. T. Lui, 1982. Insect pheromone. *U. S. Patent* 4,357,474. [*Chem. Abst.*, **98**: 125738b]
- Castellari, P. L., 1985. Oblique-banded leaf roller *Choristoneura lafauryana* Rag. (Lepidoptera, Tortricidae): field tests with synthetic sex pheromone blends. *Bull. Ist. Entomol. Univ. Bologna*, **39**: 243-260.
- Chamberlain, D. J., P. S. Beevor, A. Cork and D. R. Hall, 2003. (E)-8-Dodecenyl acetate: major component of the female sex pheromone of a Macadamia nut borer, *Ecdytolopha torticornis*. *Entomol. Exp. Appl.*, **107**: 91-93.
- Chambon, J. P. and J. d'Aguilar, 1974. Remarks on the specificity of some synthetic sexual pheromones: systematic study of tortricids in orchards. *Ann. Zool. Ecol. Anim.*, **6**: 423-430.
- Chambon, J. P. and G. Biwer, 1978. A propos de la capture d'un Tortricidae nouveau pour la France: *Laspeyresia exquisitana*. *Bull. Soc. Entomol. Fr.*, **83**: 211-213.
- Chang, V. C. S., 1995. Trapping *Cryptophlebia illepidata* and *C. ombrodelta* (Lepidoptera: Tortricidae) in macadamia in Hawaii. *Int. J. Pest Manage.*, **41**(2): 104-108.
- Charlton, R. E., J. A. Wyman, J. R. McLaughlin, J.-W. Du and W. L. Roelofs, 1991. Identification of sex pheromone of tomatopine worm, *Keiferia lycopersicella* (Wals.). *J. Chem. Ecol.*, **17**: 175-183.
- Chen, X., K. Nakamura, T. Nakanishi, T. Nakashima, M. Tokoro, F. Mochizuki and T. Fukumoto, 2006. Female sex pheromone of a carpenter moth, *Cossus insularis* (Lepidoptera: Cossidae). *J.*

- Chem. Ecol.*, **32**(3): 669-679.
- Chen, G.-F., M.-L. Sheng, T. Li, J. G. Millar and Q.-H. Zhang, 2010. Synergistic sex pheromone components of the grey-spotted tussock moth, *Orgyia ericae*. *Entomol. Exp. Appl.*, **136**: 227-234.
- Cheng, Z.-Q., J.-C. Xiao, X.-T. Huang, D.-L. Chen, J.-Q. Li, Y.-S. He, S.-R. Huang, Q.-C. Luo, C.-M. Yang and T.-H. Yang, 1981. Sex pheromone components isolated from Chaina corn borer, *Ostrinia furnacalis* Guenée (Lepidoptera: Pyralidae), (E)- and (Z)-12-tetradecenyl acetates. *J. Chem. Ecol.*, **7**: 841-851.
- Cheng, Z.-Q. 1989. Sex pheromones of two species of corn borers living in China. *Acta Phytopathol. Entomol. Hung.*, **24**: 237-240.
- Cheng, X., J. Wang and J. Zhang, 2007. Synthesis and field tests of sex attractant for *Holcocerus hippophaecolus* (Lepidoptera: Cossidae). *Linze Kexue*, **43**(5): 74-77. [SciFinder, 2007:697769]
- Chi, D. T., H. L. Thi, L. V. Vang, T. T. Thy, M. Yamamoto and T. Ando, 2024. Mass trapping of the diamondback moth (*Plutella xylostella* L.) by a combination of its sex pheromone and allyl isothiocyanate in cabbage fields in southern Vietnam. *J. Pest. Sci.*, **49**: 15-21.
- China, 1978. Isolation, identification, and synthesis of the sex pheromone of *Argyroplote* (*Eucosma*) *schistaceana* Snellen (Lepidoptera: Tortricidae). *K'o Hsueh T'ung Pao*, **23**: 438-441, 435.
- China, 1979. Isolation, identification and synthesis of EAG-active components of the pine-caterpillar moth sex pheromone. *K'o Hsueh T'ung Pao*, **24**: 1004-1008.
- Chisholm, M. D., W. F. Steck, A. P. Arthur and E. W. Underhill, 1975. Evidence for *cis*-11-hexadecen-1-ol acetate as a major component of the sex pheromone of the bertha armyworm, *Mamestra configurata* (Lepidoptera: Noctuidae). *Can. Entomol.*, **107**: 361-366.
- Chisholm, M. D., E. W. Underhill and W. F. Steck, 1979. Field trapping of the diamondback moth *Plutella xylostella* using synthetic sex attractants. *Environ. Entomol.*, **8**: 516-518.
- Chisholm, M. D., E. W. Underhill, W. Steck, K. N. Slessor and G. G. Grant, 1980(a). (Z)-5, (E)-7-Dodecadienal and (Z)-5, (E)-7-dodecadien-1-ol, sex pheromone components of the forest tent caterpillar, *Malacosoma disstria*. *Environ. Entomol.*, **9**: 278-282.
- Chisholm, M. D., W. Steck and E. W. Underhill, 1980(b). Effects of additional double bonds on some olefinic moth sex attractants.

J. Chem. Ecol., **6**: 203-212.

Chisholm, M. D., P. Palaniswamy and E. W. Underhill, 1982. Orientation disruption of male forest tent caterpillar, *Malacosoma disstria* (Hübner) (Lepidoptera: Lasiocampidae), by air permeation with sex pheromone components. *Environ. Entomol.*, **11**: 1248-1250.

Chisholm, M. D., W. F. Steck, E. W. Underhill and P. Palaniswamy, 1983. Field trapping of diamondback moth *Plutella xylostella* using an improved four-component sex attractant blend. *J. Chem. Ecol.*, **9**: 113-118.

Chisholm, M. D., D. W. Reed, E. W. Underhill, P. Palaniswamy and J. W. Wong, 1985(a). Attraction of tortricid moths of subfamily Olethreutinae to field traps baited with dodecadienes. *J. Chem. Ecol.*, **11**: 217-230.

Chisholm, M. D. and D. W. Reed, 1985(b). 9,11-Dodecadienyl alcohols, acetates, or aldehydes as synthetic chemical sex attractants for four Lepidoptera: *Cosmopterix b. gemmiferella* (Clemens), *Dichrorampha simulana* (Clemens), *Tortricidia testacea* (Packard), and an *Ancylis* sp.. *J. Chem. Ecol.*, **11**: 1341-1347.

Cho, J. R., H. H. Park, B. Y. Seo, J. H. Kim, K. H. Yum, J. K. Jung and C. Y. Yang, 2014. Identification and application of the sex pheromones of the rice green caterpillar *Naranga aeneascens* In Korea. *J. Asia-Pac. Entomol.*, **17**: 191-197.

Choi, M.-Y., H. Lim, K. C. Park, R. Adlof, S. Wang, A. Zhang and R. Jurenka 2007. Identification and biosynthetic studies of the hydrocarbon sex pheromone in *Utetheisa ornatrix*. *J. Chem. Ecol.*, **33** (7): 1336-1345.

Choi, K. S., J. M. Lee, J. H. Park, J. R. Cho, J. H. Song, D. S. Kim and K. S. Boo, 2009(a). Sex pheromone composition of the cotton caterpillar, *Palpita indica* (Lepidoptera: Pyralidae), in Korea. *J. Asia-Pac. Entomol.*, **12**: 269-275.

Choi, K. S., J. R. Cho, J. H. Song, D.-S. Kim and K. S. Boo, 2009(b). Sex pheromone composition of the variegated cutworm, *Peridroma saucia* (Lepidoptera: Noctuidae), in Korea. *J. Asia-Pac. Entomol.*, **12**: 71-77.

Choi, K. S., Y. M. Park, K. H. Choi, D. H. Kim and D. S. Kim, 2012. Sex pheromone composition of *Ascotis selenaria* (Lepidoptera: Geometridae) and its regional variation in Korea. *J. Asia-Pac. Entomol.*, **15**: 413-418.

Chow, Y. S., Y. M. Lin and C. L. Hsu, 1977. Sex pheromone of the diamondback moth (Lepidoptera: Plutellidae). *Bull. Inst. Zool.*,

Academia Sinica, **16**: 99-105.

- Chow, Y. S., R. S. Tsai, G. Gries, R. Gries and G. Khaskin, **2001**. Preliminary identification of the sex pheromone of *Orgyia postica* (Walker) (Lepidoptera: Lymantriidae). *Entomol. Sinica*, **8**: 13-20.
- Cichón, L. I., P. Trematerra, M. D. A. Coracini, D. Fernandes, M. Bengtsson and P. Witzgall, **2004**. Sex pheromone of *Argyrotaenia pomililiana* (Lepidoptera: Tortricidae), a leafroller pest of apples in Argentina. *J. Econ. Entomol.*, **97**(3): 946-949.
- Clark, J. D. and K. F. Haynes, **1990**. Sex attractant for the bluegrass webworm (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **83**: 856-859.
- Clarke, C. A., A. Cronin, W. Francke, P. Philipp, J. A. Pickett, L. J. Wadhams and C. M. Woodcock, **1996**. Mating attempts between the scarlet tiger moth, *Callimorpha dominula* L., and the cinnabar moth, *Tyria jacobaeae* L. (Lepidoptera: Arctiidae), involve a common sex pheromone composition. *Experientia*, **52**: 636-638.
- Clearwater, J. R., R. A. Galbreath, M. H. Benn and H. Young, **1986**. Female-produced sexual pheromone of *Sceliodes cordalis* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **12**: 1943-1964.
- Clearwater, J. R., S. P. Foster, S. J. Muggleston, J. S. Dugdale and E. Priesner, **1991**. Intraspecific variation and interspecific differences in sex pheromones of sibling species in *Ctenopseustis obliquana* complex. *J. Chem. Ecol.*, **17**: 413-429.
- Coffelt, J. A., K. W. Vick, P. E. Sonnet and R. E. Doolittle, **1979**. Isolation, identification, and synthesis of a female sex pheromone of the navel orangeworm, *Amyelois transitella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **5**: 955-966.
- Comeau, A. and W. L. Roelofs, **1973**. Sex attraction specificity in the Tortricidae. *Entomol. Exp. Appl.*, **16**: 191-200.
- Conner, W. E., T. Eisner, R. K. V. Meer, A. Guerrero, D. Ghiringelli and J. Meinwald, **1980**. Sex attractant of an arctiid moth (*Utetheisa ornatrix*): a pulsed chemical signal. *Behav. Ecol. Sociobiol.*, **7**: 55-63.
- Coracini, M. D. A., M. Bengtsson, A. Reckziegel, J. Lofqvist, W. Francke, E. F. Vilela, A. E. Eiras, A. Kovaleski and P. Witzgall, **2001**. Identification of a four-component sex pheromone blend in *Bonagota cranaodes* (Lepidoptera: Tortricidae). *J. Econ. Entomol.*, **94**: 911-914.

- Coracini, M. D. A., M. Bengtsson, A. Reckziegel, A. E. Eiras, E. F. Vilela, P. Anderson, W. Francke, J Löfqvist and P. Witzgall, 2003. Behavioural effects of minor sex pheromone components in Brazilian apple leafroller *Bonagota cranaodes* (Lep., Tortricidae). *J. Appl. Entomol.*, **127**: 427-434.
- Cork, A., P. S. Beevor, D. R. Hall, B. F. Nesbitt, G. S. Arida and O. Mochida, 1985(a). Components of the female sex pheromone of the yellow stem borer, *Scirpophaga incertulas*. *Entomol. Exp. Appl.*, **37**: 149-153.
- Cork, A., P. S. Beevor, D. R. Hall, Nesbitt B. F. and D. G. Champion, 1985(b). A sex attractant for the spotted bollworm *Earias vittella*. *Trop. Pest Manag.*, **31**: 156-158.
- Cork, A., D. J. Chamberlain, P. S. Beevor, D. R. Hall, B. F. Nesbitt, D. G. Champion and M. R. Attique, 1988. Components of female sex pheromone of spotted bollworm, *Earias vittella* F. (Lepidoptera: Noctuidae): identification and field evaluation in Pakistan. *J. Chem. Ecol.*, **14**: 929-945.
- Cork, A., J. Murlis and T. Megenasa, 1989. Identification and field testing of additional components of female sex pheromone of African armyworm, *Spodoptera exempta* (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **15**: 1349-1364.
- Cork, A., M. Agyen-Sampong, S. J. Fannah, P. S. Beevor and D. R. Hall, 1991. Sex pheromone of female African white rice stem borer, *Maliarpha separatella* (Lepidoptera: Pyralidae) from Sierra Leone: identification and field testing. *J. Chem. Ecol.*, **17**: 1205-1219.
- Cork, A., K. S. Boo, E. Dunkelblum, D. R. Hall, K. Jee-Rajunga, M. Kehat, E. Kong Jie, K. C. Park, P. Tepgidagarn and Liu Xun, 1992. Female sex pheromone of Oriental tobacco budworm, *Helicoverpa assulta* (Guenée) (Lepidoptera: Noctuidae): identification and field testing. *J. Chem. Ecol.*, **18**: 403-418.
- Cork, A., S. N. Alam, A. Das, C. S. Das, G. C. Ghosh, D. I. Farman, D. R. Hall, N. R. Maslen, K. Vedham, S. J. Phythian, F. M. A. Rouf and K. Srinivasan, 2001. Female sex pheromone of brinjal fruit and shoot borer, *Leucinodes orbonalis* blend optimization. *J. Chem. Ecol.*, **27**(9): 1867-1877.
- Cork, A. and E. A. Lobos, 2003. Female sex pheromone components of *Helicoverpa gelotopoeon*: first heliothine pheromone without (Z)-11-hexadecenal. *Entomol. Exp. Appl.*, **107**: 201-206.
- Coroiu, I., GH. Stan, N. Tomescu, M. C. Roman, A. T. Dragotel, I. Oprean and L. Pop, 1986. Attractant and specificity of some pheromonal synthetic compounds in *Agrotis segetum* (Lepidoptera: Noctuidae).

Rev. Roum. Biol., Ser. Biol. Anim., **31**: 109-118.

- Cortés, A. M. P., P. H. G. Zarbin, D. M. Takiya, J. M. S. Bento, A. S. Guidolin and F. L. Consoli, 2010. Geographic variation of sex pheromone and mitochondrial DNA in *Diatraea saccharalis* (Feb., 1794) (Lepidoptera: Crambidae). *J. Insect Physiol.*, **56**: 1624-1630.
- Cory, H. T., G. E. Daterman, G. D. Daves, Jr., L. L. Sower, R. F. Shepherd and C. J. Sanders, 1982. Chemistry and field evaluation of the sex pheromone of western spruce budworm, *Choristoneura occidentalis*, Freeman. *J. Chem. Ecol.*, **8**: 339-350.
- Cossé, A. A., R. Cyjon, I. Moore, M. Wysoki and D. Becker, 1992. Sex pheromone components of the giant looper, *Boarmia selenaria* Schiff. (Lepidoptera: Geometridae): identification, synthesis, electrophysiological evaluation, and behavioral activity. *J. Chem. Ecol.*, **18**: 165-181.
- Cowles, R. S., J. G. Millar, E. J. Perry, J. S. McElfresh and N. Garrison, 1996. Identification of the sex pheromone of the western poplar clearwing moth (Lepidoptera: Sesiidae). *Environ. Entomol.*, **25**: 109-113.
- Cuevas, P., R. Montoya, X. Belles, F. Camps, J. Coll, A. Guerrero and M. Riba, 1983. Initial field trials with the synthetic sex pheromone of the processionary moth *Thaumetopoea pityocampa* (Denis and schiff.). *J. Chem. Ecol.*, **9**: 85-93.
- Curtis, C. E., P. J. Landolt, R. R. Heath and R. Murphy, 1989. Attraction of western grapeleaf skeletonizer males (Lepidoptera: Zygaenidae) to *S*-(+)-2-butyl-(*Z*)-7-tetradecenoate. *J. Econ. Entomol.*, **82**: 454-457.
- Dahm, K. H., I. Richter, D. Meyer and H. Röller, 1971. The sex attractant of the Indian-meal moth *Plodia interpunctella* (Hübner). *Life Sci.*, **10**: 531-539.
- Dai, X.-J., S.-F. Xu, M.-Z. Wang, Y.-X. Zhu, X.-H. Tang, J.-W. Zhu, J.-W. Du, T.-X. Dong and M.-Z. Du, 1988. E-10,Z-12-Hexadecadienyl acetate - sex pheromone of the mulberry white caterpillar *Rondotia menciaana* Moore (Lepidoptera: Bombycidae). *Kexue Tongbao*, **33**: 1575-1576.
- Daimon, T., T. Fujii, M. Yago, Y.-F. Hsu, Y. Nakajima, T. Fujii, S. Katsuma, Y. Ishikawa and T. Shimada, 2012(a). Female sex pheromone and male behavioral responses of the bombycid moth *Trilocha varians*: comparison with those of the domesticated silkmoth *Bombyx mori*. *Naturwissenschaften*, **99**: 207-215.

- Daimon, T., T. Fujii, T. Fujii, T. Yokoyama, S. Katsuma, T. Shinoda, T. Shimada and Y. Ishikawa, 2012(b). Reinvestigation of the sex pheromone of the wild silkmoth *Bombyx mandarina*: the effects of bombykal and bombykyl acetate. *J. Chem. Ecol.*, **38**:1031-1035.
- Daterman, G. E., L. J. Peterson, R. G. Robbins, L. L. Sower, G. D. Daves, Jr. and R. G. Smith, 1976. Laboratory and field bioassay of the Douglas-fir tussock moth pheromone, (Z)-6-heneicosen-11-one. *Environ. Entomol.*, **5**: 1187-1190.
- Daterman, G. E., R. G. Robbins, T. D. Eichlin and J. Pierce, 1977. Forest Lepidoptera attracted by known sex attractants of western spruce budworms, *Choristoneura* spp. (Lepidoptera: Tortricidae). *Can. Entomol.*, **109**: 875-878.
- Daterman, G. E., H. T. Cory, L. L. Sower and G. D. Daves Jr., 1984. Sex pheromone of a conifer-feeding budworm, *Choristoneura retiniana*, Walsingham. *J. Chem. Ecol.*, **10**: 153-160.
- David, H., B. F. Nesbitt, S. Easwaramoorthy and V. Nandagopal, 1985. Application of sex pheromones in sugarcane pest management. *Proc. Indian Acad. Sci.*, **94**: 333-339.
- Davis, F. M., T. G. Bird, A. E. Knutson and S.-S. Ng, 1986. Evaluation of trapping methods for southwestern corn borer (Lepidoptera: Pyralidae) using synthetic sex pheromone. *J. Econ. Entomol.*, **79**: 1385-1389.
- Davis, H. G. and L. M. McDonough, 1981. Sex attractant for the filbertworm, *Melissopus latiferranus* (Walsingham). *Environ. Entomol.*, **10**: 390-391.
- Davis, H. G., L. M. McDonough, A. K. Burditt, Jr. and B. A. Bierl-Leonhardt, 1984. Filbertworm sex pheromone. Identification and field tests of (E,E)- and (E,Z)-8,10-dodecadien-1-ol acetate. *J. Chem. Ecol.*, **10**: 53-61.
- Davis, H. G., L. M. McDonough, C. L. Smithhisler, D. F. Brown, B. A. Leonhardt, S. Voerman and P. S. Chapman, 1991. Mint root borer, *Fumibotys fumalis* (Guenee): identification and field tests of female sex pheromone gland components. *J. Chem. Ecol.*, **17**: 249-258.
- Davis, H. G., L. M. McDonough, P. S. Chapman, C. L. Smithhisler, D. D. Bowers and S. Voerman, 1993. Sugar beet crown borer, *Hulstia undulatella* (Clemens): identification and field tests of female sex pheromone gland components. *J. Chem. Ecol.*, **19**: 433-440.
- DeBarr, G. L., L. R. Barber, C. W. Berisford and J. C. Weatherby, 1982. Pheromone traps detect webbing coneworms in loblolly pine seed

- orchards. *Southern J. Appl. Forestry.*, **6**: 122-127.
- Decamps, C., P. du Merle, C. Gourio and G. Luquet, 1981. Attraction of species of the genus *Zygaena* by tortricid sex pheromone components (Lepidoptera, Zygaenidae and Tortricidae). *Ann. Soc. Entomol. Fr.*, **17**: 441-448.
- Deland, J.-P., R. Gries, G. Gries, G. J. R. Judd and B. D. Roitberg, 1993. Sex pheromone components of the fruit-tree leaf roller, *Archips argyrospilus* (Walker) (Lepidoptera: Tortricidae), in British Columbia. *J. Chem. Ecol.*, **19**: 2855-2864.
- Delisle, J., 1992. Monitoring the seasonal male flight activity of *Choristoneura rosaceana* (Lepidoptera: Tortricidae) in eastern Canada using virgin females and several different pheromone blends. *Environ. Entomol.*, **21**(5): 1007-1012.
- Deng, J., C. Lan, J. Zhou, Y. Yao, X. Yin, K. Fu, X. Ding, W. Guo, W. Liu, N. Wang and F. Wang, 2023. Analysis of sex pheromone production and field trapping of the Asian corn borer (*Ostrinia furnacalis* Guene) in Xinjiang, China. *J. Integrat. Agric.*, **22**: 1093-1103.
- Descoins, C., E. Priesner, M. Gallois, H. Arn and G. Martin, 1978. Pheromonal secretion by virgin females of *Mamestra brassicae* L. and of *Mamestra oleracea* L. (Lepidopteras Noctuidae, Hadeninae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **286**: 77-80.
- Descoins, C., J. Einhorn, B. Lalanne-Cassou, M. Pralavorio, N. Deschamps and M. Gallois, 1985. Study of the pheromone secretion of the European carnation tortrix: *Cacoecimorpha pronubana* (Hb.), Lep. Tortricidae, Tortricinae. *Agronomie*, **5**: 19-26.
- Descoins, C., B. Lalanne-Cassou, C. Malosse and M.-L. Milat, 1986. Analysis of the sex pheromone produced by the virgin female of *Mocis latipes* (Guenée), Noctuidae, Catocalinae, from Guadeloupe (French Antilla). *C. R. Acad. Sci., Ser. 3*, **302**: 509-512.
- Descoins, C., J. F. Silvain, B. Lalanne-Cassou and H. Chéron, 1988. Monitoring of crop pests by sexual trapping of males in the French West Indies and Guyana. *Agric. Ecosystems Environ.*, **21**: 53-65.
- Descoins, C., B. Lalanne-Cassou, B. Frérot, C. Malosse and M. Renou, 1989. Comparative study of pheromonal secretions produced by arctiid and ctenuchid female moths (insects, Lepidoptera) from the neotropical area. *C. R. Acad. Sci., Ser. 3*, **309**: 577-581.
- Descoins, C., C. Malosse, M. Renou, B. Lalanne-Cassou and J. L. D. d'Aubigny, 1990. Chemical analysis of the pheromone blends produced by males and females of the neotropical moth, *Mocis megas*

- (Guénée) (Lepidoptera, Noctuidae, Catocalinae). *Experientia*, **46**: 536-539.
- Dix, M. E., A. Tagestad, J. D. Stein and M. Jacobson, 1984. Detecting tip mining Olethreutinae (Tortricidae) moths in the northern and central Great Plains with synthetic attractants. *USDA Forest Serv. Res. Note* RM **445**: 5.
- Dix, M. E. and E. W. Underhill, 1988. Sex pheromone of *Retinia metallica* (Busck) (Lepidoptera: Tortricidae): identification and field studies. *Can. Entomol.*, **120**: 721-726.
- Do, N. D., M. Kinjo, T. Taguri, Y. Adachi, R. Yamakawa and T. Ando, 2009. Synthesis and field evaluation of methyl-branched ketones, sex pheromone components produced by Lithosiinae female moths in the family of Arctiidae. *Biosci. Biotechnol. Biochem.*, **73**: 1618-1622.
- Do, N. D., K. Ohbayashi, H. Naka, K. Nakada and T. Ando, 2011. Identification and field evaluation of sex pheromone components of the pear bark miner moth, *Spulerina astauota*. *J. Chem. Ecol.*, **37**: 1222-1230.
- Doolittle, R. E., A. Tagestad and M. E. McKnight, 1976(a). Trapping carpenterworms and aspen carpenterworms with sex attractants in North Dakota. *Environ. Entomol.*, **5**: 267-269.
- Doolittle, R. E., W. L. Roelofs, J. D. Solomon, R. T. Cardé and M. Beroza, 1976(b). (Z,E)-3,5-Tetradecadien-1-ol acetate sex attractant for the carpenterworm moth *Prionoxystus robiniae* (Peck) (Lepidoptera: Cossidae). *J. Chem. Ecol.*, **2**: 399-410.
- Doolittle, R. E. and J. D. Solomon, 1986. Stereoselective synthesis of (Z,E)-3,5-tetradecadienyl acetate: sex attractant for carpenterworm moth, *Prionoxystus robiniae* (Peck) (Lepidoptera: Cossidae) and effect of isomers and monounsaturated acetates on its attractiveness. *J. Chem. Ecol.*, **12**: 619-633.
- Downham, M. C. A., D. R. Hall, D. J. Chamberlain, A. Cork, D. I. Farman, M. Tamo, D. Dahounto, B. Datinon and S. Adetonah, 2003. Minor components in the sex pheromone of legume podborer: *Maruca vitrata* development of an attractive blend. *J. Chem. Ecol.*, **29**(4): 989-1011.
- Du, J.-W., Y.-X. Zhu, T.-P. Zang, S.-F. Xu and X.-J. Dai, 1986. Studies on the precise blending of the pheromone components of the Asian corn borer, *Ostrinia furnacalis* Guenee (Lep. Pyralidae). *Contr. Shanghai Inst. Entomol.*, **6**: 17-22.
- Du, J.-W., C. Löfstedt and J. Löfqvist, 1987(a). Repeatability of

- pheromone emissions from individual female ermine moths *Yponomeuta padellus* and *Yponomeuta rorellus*. *J. Chem. Ecol.*, **13**: 1431-1441.
- Du, J., X. Dai, S. Xu, X. Tang, T. Zhang, Y. Zhu and M. Wang, 1987(b). Studies on the sex pheromone of the yellow rice borer, *Scirpophaga incertulas* (Walker), Pyralidae, Lepidoptera: identification of chemical structure and field trapping tests. *Sci. Sin., Ser. B*, **30**: 967-973.
- Du, T., J. Xiong, Z. Wang and F. Kong, 1989. (7Z,11Z)-7,11-Hexadecadienal: sex attractant of *Phyllocnistis wampella* Liu et Zeng. *Kunchong Zhishi*, **26**: 147-149. [Chem. Abst., **112**: 50570n]
- Du, Y., P. Li, Z. Chen, Y. Lin, Y. Wang and Y. Qin, 2013. Field trapping of male *Phyllonorycter ringoniella* using variable ratios of pheromone components. *Entomol. Exp. Appl.*, **146**: 357-363.
- Duckworth, W. D. and T. D. Eichlin, 1977(a). A new species of clearwing moth from southcentral Texas. *Pan-Pac. Entomol.*, **53**: 175-178.
- Duckworth, W. D. and T. D. Eichlin, 1977(b). Two new species of clearwing moths (Sesiidae) from eastern North America clarified by sex pheromones. *J. Lepi. Soc.*, **31**: 191-196.
- Duff, C. M., G. Gries, K. Mori, Y. Shirai, M. Seki, H. Takikawa, T. Sheng, K. N. Slessor, R. Gries, C. T. Maier and D. C. Ferguson, 2001. Dose pheromone biology of *Lambdina athasaria* and *L. pellucidaria* contribute to their reproductive isolation? *J. Chem. Ecol.*, **27** (3): 431-442.
- Dung, T. T., D. K. Hanh, T. P. Linh, D. T. Chi, T. T. Thy, L. V. Vang and T. V. Phen, 2021. Preliminary identification of the sex pheromones of the citrus fruit borer (*Citripestis sagitiferella* Moore) inhabiting the Mekong Delta of Vietnam. *Int. J. Plant. Anim. Environ. Sci.*, **11**: 194-205.
- Dunkelblum, E., S. Gothilf and M. Kehat, 1980. Identification of the sex pheromone of the cotton bollworm, *Heliothis armigera*, in Israel. *Phytoparasitica*, **8**: 209-211.
- Dunkelblum, E., S. Gothilf and M. Kehat, 1981. Sex pheromone of the tomato looper, *Plusia chalcites* (Esp). *J. Chem. Ecol.*, **7**: 1081-1088.
- Dunkelblum, E., M. Kehat, S. Gothilf, S. Greenberg and B. Sklarsz, 1982. Optimized mixture of sex pheromonal components for trapping of male *Spodoptera littoralis* in Israel. *Phytoparasitica*, **10**: 21-26.

- Dunkelblum, E. and S. Gothilf, 1983. Sex pheromone components of the gamma moth, *Autographa gamma* (L.) (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **38c**: 1011-1014.
- Dunkelblum, E., R. Snir, S. Gothilf and I. Harpaz, 1987(a). Identification of sex pheromone components from pheromone gland volatiles of the tomato looper, *Plusia chalcites* (Esp.). *J. Chem. Ecol.*, **13**: 991-1003.
- Dunkelblum, E., M. Kehat, M. Harel and D. Gordon, 1987(b). Sexual behaviour and pheromone titer of the *Spodoptera littoralis* female moth. *Entomol. Exp. Appl.*, **44**: 241-247.
- Dunkelblum, E. and M. Kehat, 1989. Female sex pheromone components of *Heliothis peltigera* (Lepidoptera: Noctuidae). Chemical identification from gland extracts and male response. *J. Chem. Ecol.*, **15**: 2233-2245.
- Dunkelblum, E. and M. Mazor, 1993. Chemical characterization and species specificity of sex pheromones of *Plusiinae* moths in Israel. *Arch. Insect Biochem. Physiol.*, **22**: 413-424.
- Efetov, K. A., F. Can, T. B. Toshova and M. Subchev, 2010. New sex attractant for *Jordanita anatolica* (Naufock) (Lepidoptera: Zygaenidae: Procridinae). *Acta Zool. Bulg.*, **62**: 315-319.
- Efetov, K. A., E. V. Parshkova, M.Y. Baevsky and A. I. Poddubov, 2014. Sec-butyl ester of dodecenoate: synthesis and attractive properties. *Ukr. Biochem. J.*, **86**: 175-182.
- Efetov, K. A., E. E. Kucherenko, E. V. Parshkova and G. M. Tarmann, 2016. 2-Butyl 2-dodecenoate, a new sex attractant for *Jordanita* (*Tremewania*) *notata* (Zeller, 1847) and some other Procridinae species (Lepidoptera: Zygaenidae). *SHILAP Rev. Lepid.*, **44**: 519-527.
- Efetov, K. A., E. E. Kucherenko and G. M. Tarmann, 2019. New synthetic sex attractants for the males of two endemic Iberian Procridinae species (Lepidoptera: Zygaenidae). *SHILAP Rev. Lepid.*, **47**: 307-315.
- Efetov, K. A. and E. E. Kucherenko, 2020(a). Structural analysis of sex pheromones and attractants in Zygaenidae (Insecta, Lepidoptera): biochemical and evolutionary aspects. *J. Evol. Biochem. Physiol.*, **56**: 381-395.
- Efetov, K. A., E. E. Kucherenko and G. M. Tarmann, 2020(b). An application of the synthetic sex attractants from the series "EFETOV-2" for studying Procridinae in Italy (Lepidoptera:

- Zygaenidae). *SHILAP Rev. Lepid.*, **48**: 733-749.
- Efetov, K. A. and E. E. Kucherenko, **2021**. Enantiomers of 2-butyl dodec-2-enoate as sex attractants. *Rus. J. Bioorg. Chem.*, **47**: 1043-1050.
- Eichlin, T. D. and S. Passoa, **1983**. A new clearwing moth (Sesiidae) from Central America: a stem borer in *Mimosa pigra*. *J. Lepid. Soc.*, **37**: 193-206.
- Einhorn, J., J.-Y. Lallemand, P. Zagatti, M. Gallois, H. Virelizier, J. Riom and P. Menassieu, **1982**. Isolation and identification of the sex pheromone blend of *Hyphantria cunea* (Drury) (Lepidoptera, Arctiidae). *C. R. Acad. Sci., Ser. 2*, **294**: 41-44.
- Einhorn, J., P. Menassier, D. Michelot and J. Riom, **1983**. Sex trapping of the processionary moth *Thaumetopoea pityocampa* Schiff. (Lep., Notodontidae) with synthetic attractants. First experiments in the south-west of France. *Agronomie*, **3**: 499-505.
- Einhorn, J., B. Boniface, M. Renou and M.-L. Milat, **1984(a)**. Study on the sex pheromone of *Arctia villica* L. (Lepidoptera, Arctiidae). *C. R. Acad. Sci., Ser. 3*, **298**: 573-576.
- Einhorn, J., F. Beauvais, M. Gallois, C. Descoins and R. Causse, **1984(b)**. Secondary components of the codling moth (*Cydia pomonella* L.) (Lepidoptera, Tortricidae) sex pheromone. *C. R. Acad. Sci., Ser. 3*, **299**: 773-778.
- Einhorn, J., P. Witzgall, H. Audemard, B. Boniface and R. Causse, **1986**. Secondary components of the codling moth (*Cydia pomonella* L.) (Lepidoptera, Tortricidae) sex pheromone. II. First results on the behavioural effects. *C. R. Acad. Sci., Ser. 3*, **302**: 263-266.
- Eiras, A. E., A. Kovaleski, E. F. Vilela, J. P. Chambon, C. R. Unelius, A.-K. Borg-Karlson, I. Liblikas, R. Mozūraitis, M. Bengtsson and P. Witzgall, **1999**. Sex pheromone of the Brazilian apple leafroller, *Bonagota cranaodes* Meyrick (Lepidoptera, Tortricidae). *Z. Naturforsch.*, **54c**: 595-601.
- El-Sayed, A., J. Gödde, P. Witzgall and H. Arn, **1999**. Characterization of pheromone blend for grapevine moth, *Lobesia botrana* by using flight track recording. *J. Chem. Ecol.*, **25**(2): 389-400.
- El-Sayed, A. M., H. M. Fraser and R. M. Trimble, **2001**. Chemical identification and behavioral activity of (Z)-11-tetradecenal in an eastern North American population of the obliquebanded leafroller. *Can. Entomol.*, **133**: 365-374.
- El-Sayed, A. M., J. Delisle, N. De Lury, L. J. Gut, G. J. R. Judd,

- S. Legrand, W. H. Reissig, W. L. Roelofs, C. R. Unelius and R. M. Trimble, 2003. Geographic variation in pheromone chemistry, antennal electrophysiology, and pheromone-mediated trap catch of North American populations of the obliquebanded leafroller. *Environ. Entomol.*, **32**: 470-476.
- El-Sayed, A. M., A. R. Gibb, D. M. Suckling, B. Bunn, S. Fielder, D. Comeskey, K. A. Moss, L. A. Manning, S. P. Foster, B. D. Morris, T. Ando and K. Mori, 2004(a). Identification and synthesis of the sex pheromone of painted apple moth. *HortResearch Client Report* No: 13463, Contact No: 18879.
- El-Sayed, A. M., L. I. Wainman, E. M. Santangelo, C. R. Unelius and R. M. Trimble, 2004(b). Relative attractiveness of (10E)-dodecen-1-yl acetate and (4E,10E)-dodecadien-1-yl acetate to male spotted tentiform leafminers *Phyllonorycter blancardella* (F.). *J. Chem. Ecol.* **30**: 1827-1838.
- El-Sayed, A. M., A. R. Gibb, D. M. Suckling, B. Bunn, S. Fielder, D. Comeskey, L. A. Manning, S. P. Foster, B. D. Morris, T. Ando and K. Mori, 2005(a). Identification of sex pheromone components of the painted apple moth: a tussock moth with a thermally labile pheromone component. *J. Chem. Ecol.*, **31**(3): 621-646.
- El-Sayed, A. M., L. I. Wainman, E. M. Santangelo, C. R. Unelius and R. M. Trimble, 2005(b). Sex pheromone of a North American population of the spotted tentiform leafminer, *Phyllonorycter blancardella*. *Entomol. Exp. Appl.*, **116**: 143-148.
- El-Sayed, A. M., A. R. Gibb and D. M. Suckling, 2005(c). Chemistry of the sex pheromone gland of the fall webworm, *Hyphantria cunea*, discovered in New Zealand. *New Zealand Plant Protection*, **58**: 31-36.
- El-Sayed, A. M., V. J. Mitchell, L.-A. M. Manning and D. M. Suckling, 2011. New sex pheromone blend for the lightbrown apple moth, *Epiphyas postvittana*. *J. Chem. Ecol.*, **37**: 640-646.
- El-Sayed, A. M., A. R. Gibb, V. J. Mitchell, L. M. Manning, J. Revell, B. Thistleton and D. M. Suckling, 2013. Identification of the sex pheromone of *Conogethes pluto*: a pest of *Alpinia*. *Chemoecology*, **23**: 93-101.
- El-Sayed, A. M. and L.-A. M. Manning, 2022. Identification of the sex pheromone of the pink grass worm, *Tmetolophota atristriga*, reveals possible population differences in male response to sex pheromone. *J. Chem. Ecol.*, **48**: 683-689.
- El-Sayed, A. M. and B. Bunn, 2024. 5-Methylheptadecane: sex pheromone of the broom twig miner, *Leucoptera spartifoliella*, a biological

- control agent for the Scotch broom. *J. Chem. Ecol.*, **50**: 874–883.
- El-Sayed, A. M. and A. R. Gibb, 2025. Identification of components of the sex pheromone of the kōwhai moth, *Uresiphita polygonalis maorialis*, a New Zealand native crambid. *J. Chem. Ecol.*, **51**: 4.
- Evenden, M. L., J. H. Borden, G. A. Van Sickle and G. Gries, 1995. Development of a pheromone-based monitoring system for western hemlock looper (Lepidoptera: Geometridae): effect of pheromone dose, lure age, and trap type. *Environ. Entomol.*, **24**(4): 923–932.
- Evenden, M. L. and R. Gries, 2006. Sex pheromone of the large aspen tortrix, *Choristoneura conflictana* (Lepidoptera: Tortricidae). *Chemoecology* **16**: 115–122.
- Evenden, M.L., B. A. Mori, R. Gries and J. Otani, 2010. Sex pheromone of the red clover casebearer moth, *Coleophora deauratella*, an invasive pest of clover in Canada. *Entomol. Exp. Appl.*, **137**: 255–261.
- Fang, Y.-L., J.-H. Sun, C.-H. Zhao and Z.-N. Zhang, 2005. Sex pheromone components of the sandthorn carpenterworm, *Holococerus hippophaecolus*. *J. Chem. Ecol.* **31**: 39–48.
- Farine, J.-P., B. Frérot and J. Isart, 1981. Chemical isolation factor in pheromonal secretion of two noctuids (Hadeninae): *Mamestra brassicae* (L.) and *Pseudaletia unipuncta* (Haw.). *C. R. Acad. Sci., Ser. 3*, **292**: 101–104.
- Félix, A.-E., G. Genestier, C. Malosse, P.-A. Calatayud, B. L. Rü, J.-F. Silvain and B. Frérot, 2009. Variability in pheromone communication among different haplotype populations of *Busseola fusca*. *J. Chem. Ecol.*, **35**: 618–623.
- Félix, A.-E., P.-A. Calatayud, B. Le Ru, J.-F. Silvain and B. Frérot, 2011. Sex pheromone composition and reproductive isolation in two *Busseola* species (Lepidoptera: Noctuidae) in Kenya. *Chemoecology*, **21**: 107–111.
- Ferrao, P., G. Gries, P. D. C. Wimalaratne, C. T. Maier, R. Gries, K. N. Slessor and J. Li 1998. Sex pheromone of apple blotch leaf miner, *Phyllonorycter crataegella*, and its effect on *P. mespilella* pheromone communication. *J. Chem. Ecol.*, **24**(12): 2059–2078.
- Ferrara, F. A. A., E. F. Vilela, G. N. Jham, A. E. Eiras, M. C. Picanco, A. B. Attygalle, A. Svatoš, R. T. S. Frighetto and J. Meinwald, 2001. Evaluation of the synthetic major component of the sex hormone of *Tuta absoluta* (Meyrick) (Lepidoptera: Gelechiidae).

- J. Chem. Ecol.*, **27**(5): 907-917.
- Filho, M. M., E. F. Vilela, A. B. Attygalle, J. Meinwald, A. Svatoš and G. N. Jham, 2000. Field trapping of tomatomoth, *Tuda absoluta* with pheromone traps. *J. Chem. Ecol.*, **26**(4): 875-881.
- Fitzpatrick, S. M. and J. T. Troubridge, 1992. Relative efficacies of two commercial pheromone blends for monitoring the blackheaded fireworm (Lepidoptera: Tortricidae) of Cranberries. *J. Econ. Entomol.*, **85**(3): 947-949.
- Fleischer, S. J., C. L. Harding, P. E. Blom, J. White and J. Grehan, 2005. *Spodoptera frugiperda* pheromone lures to avoid nontarget captures of *Leucania phragmatidicola*. *J. Econ. Entomol.*, **98**: 66-71.
- Flint, H. M., M. Balasubramanian, J. Campero, G. R. Strickland, Z. Ahmad, J. Barral, S. Barbosa and A. F. Khail, 1979. Pink bollworm: response of native males to ratios of *Z,Z*- and *Z,E*-isomers of gossyplure in several cotton growing areas of the world. *J. Econ. Entomol.*, **72**: 758-762.
- Forno, H., P. Arretz and R. H. Gonzalez, 1975. Sex pheromone in the detection and evaluation of the oriental fruit moth in Chile. *FAO Plant Prot. Bull.*, **23**: 4-11.
- Foster, S. P., J. R. Clearwater, S. J. Muggleston, J. S. Dugdale and W. L. Roelofs, 1986. Probable sibling species complexes within two described New Zealand leafroller moths. *Naturwissenschaften*, **73**: 156-158.
- Foster, S. P. and W. L. Roelofs, 1987(a). Sex pheromone differences in populations of the brownheaded leafroller, *Ctenopseustis obliquana*. *J. Chem. Ecol.*, **13**: 623-629.
- Foster, S. P., J. R. Clearwater and W. L. Roelofs, 1987(b). Sex pheromone of *Planotortrix* species found on mangrove. *J. Chem. Ecol.*, **13**: 631-637.
- Foster, S. P. and J. S. Dugdale, 1988. A comparison of morphological and sex pheromone differences in some New Zealand Tortricinae moths. *Biochem. System. Ecol.*, **16**: 227-232.
- Foster, S. P., J. R. Clearwater and S. J. Muggleston, 1989. Intraspecific variation of two components in sex pheromone gland of *Planotortrix excessana* sibling species. *J. Chem. Ecol.*, **15**: 457-465.
- Foster, S. P., J. R. Clearwater, S. J. Muggleston and P. W. Shaw, 1990. Sex pheromone of a *Planotortrix excessana* sibling species and

- reinvestigation of related species. *J. Chem. Ecol.*, **16**: 2461-2474.
- Foster, S. P., J. S. Dugdale and C. S. White, 1991. Sex pheromones and the status of greenheaded and brownheaded leafroller moths in New Zealand. *New Zealand J. Zool.*, **18**: 63-74.
- Foster, S. P., D. Steven, C. McKenna and S. J. Muggleston, 1993. Sex pheromones of two New Zealand tortricid moths, *Cnephasia jactatana* and *Merophyas leucaniana*. *New. Zeal. J. Zool.*, **20**: 81-88.
- Foster, S. P. and W. P. Thomas, 2000. Identification of a sex pheromone component of the raspberry budmoth, *Heterocrossa rubophaga*. *J. Chem. Ecol.*, **26**(11): 2549-2555.
- Foster, S. P., B. D. Morris, L. D. Charlet, T. Gross and S. Grugel, 2003. Identification of the sex pheromone of *Cochylis arthuri* (Lepidoptera: Cochylidae). *Can. Entomol.*, **135**: 713-719.
- Fouda, H. G., J. N. Seiber and O. G. Bacon, 1975. A potent sex attractant for the potato tuberworm moth. *J. Econ. Entomol.*, **68**: 423-427.
- Francke, W., S. Franke, M. Toth, G. Szöcs, P. Guerin and H. Arn, 1987. Identification of 5,9-dimethylheptadecane as a sex pheromone of the moth *Leucoptera scitella*. *Naturwissenschaften*, **74**: 143-144.
- Francke, W., M. Toth, G. Szöcs, W. Krieg, H. Ernst and E. Buschmann, 1988. Identification and synthesis of dimethylalkanes as sex attractants of female leaf miner moths (Lyonetiidae). *Z. Naturforsch.*, **43c**: 787-789.
- Francke, W., E. Plass, N. Zimmermann, H. Tietgen, T. Tolasch, S. Franke, M. Subchev, T. Toshova, J. A. Pickett, L. J. Wadhams and C. M. Woodcock, 2000. Major sex pheromone component of female herald moth *Scoliopteryx libatrix* is the novel branched alkene (6Z,13)-methylheneicosene. *J. Chem. Ecol.*, **26**(5): 1135-1149.
- Francke, W., S. Franke, J. Bergmann, T. Tolasch, M. Subchev, A. Mircheva, T. Toshova, A. Svatoš, B. Kalinová, Z. Karpati, G. Szöcs and M. Tóth, 2002. Female sex pheromone of *Cameraria ohridella* Desch. and Dim. (Lepidoptera: Gracillariidae): structure confirmation, synthesis and biological activity of (8E,10Z)-8,10-tetradecadienal and some analogues. *Z. Naturforsch.*, **57c**: 739-752.
- Francke, W., V. Karalius, E. Plass, L. Lehmann, A. D. Santos, V. Bûda, A.-K. Borg-Karlson and R. Mozûraitis, 2004. New type of Sesiidae sex pheromone identified from the hornet moth *Sesia apiformis*. *J. Chem. Ecol.*, **30**(4): 805-817.

- Frerot, B., E. Priesner and M. Gallois, 1979(a). A sex attractant for the green budworm moth, *Hedya nubiferana*. *Z. Naturforsch.*, **34c**: 1248-1252.
- Frerot, B., C. Descoins, B. Lalanne-Cassou, P. Saglio and F. Beauvais, 1979(b). Field trapping of tortricids from apple orchards with synthetic sex attractants. *Ann. Zool. Ecol. Anim.*, **11**: 617-636.
- Frerot, B., B. Boniface, J.-P. Chambon and Y. Meritan, 1982(a). Trapping with synthetic sex attractants to determine the distribution of three tortricid moths of apple orchards in the Paris area. *Agronomie*, **2**: 885-893.
- Frerot, B., M. Gallois, M. Lettere, J. Einhorn, D. Michelot and C. Descoins, 1982(b). Sex pheromone of *Pandemis heparana* (Den. and Schiff.) (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **8**: 663-670.
- Frerot, B., M. Renou, M. Gallois and C. Descoins, 1983. A sex attractant for *Archips xylosteana* L. (Lepid., Tortricidae, Tortricinae). *Agronomie*, **3**: 173-178.
- Frerot, B., M. Renou, C. Descoins and M. Gallois, 1984. A sex pheromone for *Argyrotaenia pulchellana* (Haw.) (Lepidoptera, Tortricinae). *Agronomie*, **4**: 565-572.
- Frerot, B., J.-P. Chambon, B. Boniface and C. Descoins, 1985. Sexual trapping of various Lepidoptera males by synthetic attractants. *Ann. Soc. Entomol. Fr.*, **21**: 297-305.
- Frérot, B., C. Malosse, M.-L. Milat, J.-Y. Lallemand, J. Soulié and A. Brunetière, 1986. Isolation and identification of three constituents of the sexual pheromone of *Zeuzera pyrina* L. *C. R. Acad. Sci., Ser. 2*, **302**: 413-415.
- Frérot, B., J.-P. Chambon and C. Cocquempot, 1987(a). Some results on sexual trapping of Lepidoptera Noctuidae and Geometridae. *Ann. Soc. Entomol. Fr.*, **23**: 151-154.
- Frerot, B., M. Gallois, C. Malosse and R. Bues, 1987(b). Identification of the sex pheromone of *Mamestra biren* (Goeze) ex *M. glauca* (Hb.). *C. R. Acad. Sci., Ser. 3*, **304**: 139-142.
- Frérot, B., M. Renou, C. Malosse and C. Descoins, 1988(a). Isolation and identification of pheromone compounds in female moths of *Tyria jacobaeae* (Lepidoptera, Arctiidae). Biological characterization of the absolute configuration of the main component. *Entomol. Exp. Appl.*, **46**: 281-289.
- Frerot, B., J.-R. Pougny, M.-L. Milat, P. Rollin and C. Malosse, 1988(b). Study of the pheromonal secretion of the tiger moth, *Cymbalophora*

- pudica* (Esper) and enantioselectivity of its pheromonal perception. *C. R. Acad. Sci., Ser. 3*, **306**: 157-160.
- Frerot, B., B. Lalanne-Cassou, M. Gallois, C. Malosse, A.-H. Cain and R. Bues, 1988(c). Physico-chemical analysis of the sex pheromone blend produced by the Hadeninae noctuid moth: *Mamestra suasa* (D. et S.). *C. R. Acad. Sci., Ser. 3*, **307**: 785-788.
- Frérot, B., C. Malosse, M. L. Milat, G. Demolin, J. C. Martin, M. Khemici, M. Zamoun and M. Gachi, 1990(a). Chemical analysis of the sex pheromone glands of *Thaumetopoea bonjeani* (Powell) (Lep., Thaumetopoeidae). *J. Appl. Entomol.*, **109**: 210-212.
- Frérot, B., J. P. Chambon, C. Malosse and G. Genestier, 1990(b). A sex attractant for *Lobesia bicinctana* (Dup.) (Lep., Tortricidae). *J. Appl. Entomol.*, **110**: 365-369.
- Frérot, B. and S. P. Foster, 1991. Sex pheromone evidence for two distinct taxa within *Graphania mutans* (Walker). *J. Chem. Ecol.*, **17**: 2077-2093.
- Frérot, B., C. Malosse, R. D. Chenon, P. H. Ducrot and A.-H. Cain, 1993(a). Identification of the sex pheromone components in *Pareuchaetes pseudoinsulata* Rego Barros (Lepidoptera, Arctiidae). *C. R. Acad. Sci., Ser. 2*, **317**: 1045-1050.
- Frérot, B., J. S. Dugdale and S. P. Foster, 1993(b). Chemotaxonomy of some species of moths in the New Zealand genus *Graphania* based on sex pheromones. *New Zealand J. Zool.*, **20**: 71-80.
- Frerot, B. and G. Demolin, 1993(c). Sex pheromone of the processionary moths and biosystematic considerations within the genus *Thaumetopoea* (Thaumetopoeidae Thaumetopoeinae). *Boll. Zool. Agric. Bachic. Ser. II*, **25**(1): 33-40.
- Frérot, B., J.-P. Marro and C. Malosse, 1995. *In vitro* incubation of sex pheromone gland and identification of pheromone components in *Cydia splendana* (Hb). *C. R. Acad. Sci., Life Sciences*, **318**: 447-451.
- Frilli, F., 1974. Observations on the use of *Cydia molesta* pheromones. *Entomologica*, **10**: 31-56.
- Fu, X.-Y. X.-Q. Pan and X.-Z. Meng, 2002. Female sex pheromone of the Chinese tortrix *Cydia trasias*. *Entomol. Exp. Appl.*, **103**: 183-186.
- Fu, X., J. Tabata, Y. Huang, T. Takanashi, S. Ohno, H. Honda, S. Tatsuki and Y. Ishikawa, 2004. Female sex pheromone of *Ostrinia orientalis* - throwing a light on the relationship between *O. orientalis* and

- the European corn borer, *O. nubilalis*. *Chemoecology*, **14**: 175-180.
- Fu, N., F. H. Magsi, Y. Zhao, X. Cai, Z. Li, L. Bian, C. Xiu, Z. Chen and Z. Luo, **2022**. Identification and field evaluation of sex pheromone components and its antagonist produced by a major tea pest, *Archips strobilator* (Lepidoptera: Tortricidae). *Insects*, **13**: 1056.
- Fujii, T., D. Iwai, F. Mochizuki and K. Fukumoto, **2010** (a). Sex pheromones and sex attractants from *Platyptillia ignifera*. *Jpn. Kokai Tokkyo Koho* JP 2010018543. [*SciFinder*. AN 2010:110975]
- Fujii, T., R. Nakano, Y. Takubo, S. Qian, R. Yamakawa, T. Ando and Y. Ishikawa, **2010** (b). Female sex pheromone of a lichen moth *Eilema japonica* (Arctiidae, Lithosiinae). *J. Insect Physiol.*, **56**: 1986-1991.
- Fujii, T., H. Miyake and Y. Nasu, **2013** (a). Sex attractants of *Carposina coreana*. *Jpn. Kokai Tokkyo Koho* JP 2013241377.
- Fujii, T., R. Yamakawa, Y. Terashima, S. Imura, K. Ishigaki, M. Kinjo and T. Ando, **2013** (b). Propionates and acetates of chiral secondary alcohols, novel sex pheromone components produced by a lichen moth *Barsine expressa* (Arctiidae: Lithosiinae). *J. Chem. Ecol.*, **39**: 28-36.
- Fujimura, T., **1976**. Mating behavior and sex attractant of the spotted cutworm moth, *Amathes c-nigrum* L. (Lepidoptera: Noctuidae). *Jap. J. Appl. Entomol. Zool.*, **20**: 133-138. (in Japanese)
- Gago, R., J. D. Allison, J. S. McElfresh, K. F. Haynes, J. McKenney, A. Guerrero and J. G. Millar, **2013**. A tetraene aldehyde as the major sex pheromone component of the promethea moth (*Callosamia promethea* (Drury)). *J. Chem. Ecol.*, **39**: 1263-1272.
- Galbreath, R. A., M. H. Benn, H. Young, V. A. Holt, **1985**. Sex pheromone components in the New Zealand greenheaded leafroller *Planotortrix excessana* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **40 c**: 266-271.
- Gemenio, C. and K. F. Haynes **1998**. Chemical and behavioral evidence for a third pheromone component in a North American population of the black cutworm moth, *Agrotis ipsilon*. *J. Chem. Ecol.*, **24**(6): 999-1011.
- Gemenio, C., A. F. Lutfallah and K. F. Haynes, **2000**. Pheromone blend variation and cross-attraction among populations of the black cutworm moth (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **93**(6): 1322-1328.

- Gentry, C. R., M. Beroza and J. L. Blythe, 1975. Pecanbudmoth: captures in Georgia in traps baited with the pheromone of the oriental fruit moth. *Environ. Entomol.*, **4**: 227-228.
- Germinara, G. S., A. Elgargoti, A. D. Cristofaro and G. Rotundo, 2007. Female sex pheromone of *Sesamia cretica*: chemical and behavioural evidence for a three-component blend. *Entomol. Exp. Appl.*, **124** (2): 213-219.
- Ghizdavu, I., F. Hodosan, N. Popovici, A. Botar and A. Barabas, 1979. A sex attractant for *Autographa (Plusia) gamma* L. moth. *Rev. Roum. Biol., Ser. Biol. Anim.*, **24**: 87-90.
- Ghizdavu, I., F. P. Hodosan, I. Oprean, A. Gocan, H. Ciupe and S. Matic, 1985. Sex attractants for five species of Noctuidae determined using traps. *Rev. Roum. Biol., Ser. Biol. Anim.*, **30**: 25-29.
- Ghizdavu, I., F. P. Hodosan and I. Oprean, 1987. Attractifs sexuels spécifiques pour *Adoxophyes orana* F. v. R. et *Archips crataegana* Hb. *Rev. Roum. Biol., Ser. Biol. Anim.*, **32**: 23-27.
- Gibb, A. R., L. E. Jamieson, D. M. Suckling, P. Ramankutty and P. S. Stevens, 2005. Sex pheromone of the citrus flower moth *Prays nephelomina*: pheromone identification, field trapping trials, and phenology. *J. Chem. Ecol.*, **31**(7): 1633-1644.
- Gibb, A. R., D. M. Suckling, B. D. Morris, T. E. Dawson, B. Bunn, D. Comeskey and J. J. Dymock, 2006(a). (Z)-7-Tricosene and monounsaturated ketones as sex pheromone components of the Australian guava moth, *Coscinoptycha improbana*: identification, field trapping, and phenology. *J. Chem. Ecol.*, **32**(1): 221-237.
- Gibb, A. R., D. Comeskey, L. Berndt, E. G. Brockerhoff, A. M. El-Sayed, H. Jactel and D. M. Suckling, 2006(b). Identification of sex pheromone components of a New Zealand geometrid moth, the common forest looper *Pseudocoremia suavis*, reveals a possible species complex. *J. Chem. Ecol.*, **32**(4): 865-879.
- Gibb, A. R., B. Pinese, D. Tenakanai, A. P. Kawi, B. Bunn, P. Ramankutty and D. M. Suckling, 2007(a). (Z)-11-Hexadecenal and (3Z,6Z,9Z)-tricosatriene: sex pheromone components of the red banded mango caterpillar *Deanolis sublimbalis*. *J. Chem. Ecol.*, **33** (3): 579-589.
- Gibb, A. R., D. M. Suckling, A. M. El-Sayed, B. Bohman, C. R. Unelius, J. J. Dymock, M. L. Larsen and B. E. Willoughby, 2007(b). (11Z,13E)-Hexadecadien-1-yl acetate: sex pheromone of the grass webworm *Herpetogramma licarsisalis* - Identification, synthesis, and field bioassays. *J. Chem. Ecol.*, **33** (4): 839-847.

- Gibb, A. R., D. M. Suckling, S. Fielder, B. Bunn, L. E. Jamieson, M. L. Larsen, G. H. Walter and D. J. Kriticos, 2008. Major sex pheromone components of the Australian gum leaf skeletonizer *Uraba lugens*: (10*E*,12*Z*)-hexadecadien-1-yl acetate and (10*E*,12*Z*)-hexadecadien-1-ol. *J. Chem. Ecol.*, **34** (9): 1125-1133.
- Gigon, F., 1980. Utilisation des pièges à phéromone sexuelle synthétique dans l'étude faunistique des tordeuses (Tortricidae s. str.) de vergers neuchâtelois. *Bulletin de la Société Neuchâteloise des Sciences Naturelles*, **103**: 45-56.
- Gillespie, D. R., G. G. S. King, V. A. Salas-Reyes and K. N. Slessor, 1984. Determination of the sex-pheromone components of *Cheimophila salicella* (Lepidoptera: Oecophoridae), a pest of blueberry in British Columbia. *Can. Entomol.*, **116**: 1397-1402.
- Gothilf, S., M. Kehat, M. Jasobcon and R. Galun, 1978. Sex attractants for male *Heliothis armigera* (Hbn.). *Experientia*, **34**: 853-854.
- Granges, J. and M. Baggiolini, 1971. Une phéromone sexuelle synthétique attractive pour le carpocapse des prunes (*Grapholitha funebrana* Tr. [Lép. Tortricidae]). *Rev. Suisse Vitic. Arboric.*, **3**: 93-94.
- Grant, G. G., 1977. Interspecific pheromone responses of tussock moths and some isolating mechanisms of eastern species. *Environ. Entomol.*, **6**: 739-742.
- Grant, G. G. and K. N. Slessor, 1983. Sex pheromone of the maple leaf roller *Cenopsis acerivorana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **115**: 189-194.
- Grant, G. G., L. MacDonald, D. Frech, K. Hall and K. N. Slessor, 1985. Sex attractants for some eastern species of *Rhyacionia*, including a new species, and *Eucosma gloriola* (Lepidoptera: Tortricidae). *Can. Entomol.*, **117**: 1489-1496.
- Grant, G. G., Y. H. Prévost, K. N. Slessor, G. G. S. King and R. J. West, 1987(a). Identification of the sex pheromone of the spruce coneworm, *Dioryctria reniculelloides* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **16**: 905-909.
- Grant, G. G., D. Frech, L. MacDonald, K. N. Slessor and G. G. S. King, 1987(b). Copulation releaser pheromone in body scales of female whitemarked tussock moth, *Orgyia leucostigma* (Lepidoptera: Lymantriidae): identification and behavioral role. *J. Chem. Ecol.*, **13**: 345-356.
- Grant, G. G., W. H. Fogal, R. J. West, K. N. Slessor and G. E. Miller,

1989. A sex attractant for the spruce seed moth, *Cydia strobilella* (L.), and the effect of lure dosage and trap height on capture of male moths. *Can. Entomol.*, **121**: 691-697.
- Grant G. G., B. Pendrel, K. M. Slessor, X. Z. Meng and W. E. Miller, 1991. Identification of sex pheromone components for two lepidopteran defoliators, the oak olethreutid leafroller, *Pseudexentera spoliata* (Clemens), and the aspen leafroller, *Pseudexentera oregonana* (Walsingham). *Can. Entomol.*, **123**: 1209-1218.
- Grant, G. G., S. A. Katovich, D. J. Hall, D. A. Lombardo and K. N. S. A. Lessor, 1993. Sex pheromone identification and trapping of *Dioryctria resinosella* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **22**(1): 154-161.
- Grant, G. G., D. Langevin, J. Liska, P. Kapitola and J. M. Chong, 1996. Olefin inhibitor of gypsy moth, *Lymantria dispar*, is a synergistic pheromone component of nun moth, *L. monacha*. *Naturwissenschaften*, **83**: 328-330.
- Grant, G. G., P. D. Groot, D. Langevin, S. A. Katovich, K. N. Slessor and W. E. Miller, 2002. Sex attractants and seasonal flight patterns for three *Eucosma* (Lepidoptera: Tortricidae) species sympatric in eastern pine seed orchards and plantations. *Can. Entomol.*, **134**: 391-401.
- Grant, G. G., K. N. Slessor, Weiliu and M. M. Abou-Zaid, 2003. (Z,Z)-6,9-Heneicosadien-11-one, labile sex pheromone of the whitemarked tussock moth, *Orgyia leucostigma*. *J. Chem. Ecol.*, **29**(3): 589-601.
- Grant, G. G., M. D. Coppens, L. K. Hartling, D. O'Shea, D. Winter, J. Gordon, J. Rudderham and W. Liu, 2008. Operational monitoring of the whitemarked tussock moth with Multipher traps generating the unstable pheromone component (Z,Z)-6,9-heneicosadien-11-one from a precursor. *Entomol. Exp. Appl.*, **126**: 174-178.
- Gray, T. G., K. N. Slessor, G. G. Grant, R. F. Shepherd, E. H. Holsten and A. S. Tracey, 1984(a). Identification and field testing of pheromone components of *Choristoneura orae* (Lepidoptera: Tortricidae). *Can. Entomol.*, **116**: 51-56.
- Gray, T. G., K. N. Slessor, R. F. Shepherd, G. G. Grant and J. F. Manville, 1984(b). European pine shoot moth, *Rhyacionia buoliana* (Lepidoptera: Tortricidae): identification of additional pheromone components resulting in an improved lure. *Can. Entomol.*, **116**: 1525-1532.

- Gray, T. G. and G. Gries, 1993. Sex pheromone components of an undescribed *Choristoneura* species (Lepidoptera: Tortricidae) on lodgepole pine in British Columbia. *J. Entomol. Soc. Brit. Columb.*, **90**: 13-18.
- Gray, T. G., R. F. Shepherd, G. Gries and R. Gries, 1996. Sex pheromone component of the western blackheaded budworm, *Acleris gloverana* Walsingham (Lepidoptera: Tortricidae). *Can. Entomol.*, **128**: 1135-1142.
- Green, N., M. Jacobson and J. C. Keller, 1969. Hexalure, an insect sex attractant discovered by empirical screening. *Experientia*, **25**: 682-683.
- Greenfield, M. D. and M. G. Karandinos, 1979. Resource partitioning of the sex communication channel in clearwing moths (Lepidoptera: Sesiidae) of Wisconsin. *Ecol. Monogr.*, **49**: 403-426.
- Greenway, A. R., 1984. Sex pheromone of the pea moth, *Cydia nigricana* (F.) (Lepidoptera: Olethreutidae). *J. Chem. Ecol.*, **10**: 973-982.
- Griepink, F. C., T. A. Beek, J. H. Visser, S. Voerman and A. Groot, 1995. Isolation and identification of sex pheromone of *Symmetrischema tangolias* (Gyen) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **21**: 2003-2013.
- Griepink, F. C., T. A. Beek, M. A. Posthumus, A. Groot, J. H. Visser and S. Voerman, 1996. Identification of the sex pheromone of *Scrobipalpula absoluta*; determination of double bond positions in triple unsaturated straight chain molecules by means of dimethyl disulphide derivatization. *Tetrahedron Lett.*, **37**: 411-414.
- Gries, G., R. Gries, J. H. Borden, J.-X. Li, K. N. Slessor, G. G. S. King, W. W. Bowers, R. J. West and E. W. Underhill, 1991. 5,11-Dimethyl-heptadecane and 2,5-dimethylheptadecane: sex pheromone components of the geometrid moth, *Lambdina fiscellaria fiscellaria*. *Naturwissenschaften*, **78**: 315-317.
- Gries, G., R. Gries, E. W. Underhill and L. Humble, 1993(a). (6Z,9Z-3R,4S)-Epoxy-heptadecadiene: major sex pheromone component of the larch looper, *Semiothisa sexmaculata* (Packard) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **19**: 843-850.
- Gries, G., R. Gries, S. H. Krannitz, J. Li, G. G. S. King, K. N. Slessor, J. H. Borden, W. W. Bowers, R. J. West and E. W. Underhill, 1993(b). Sex pheromone of the western hemlock looper, *Lambdina fiscellaria lugubrosa* (Hulst) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **19**: 1009-1019.
- Gries, G., G. G. S. King, R. Gries, P. D. C. Wimalaratne, T. G. Gray,

- R. F. Shepherd, J.-X. Li, K. N. Slessor and G. Khaskin, 1993(c). 3,13-Dimethylheptadecane: major sex pheromone component of the western false hemlock looper, *Nepytia freemani* Munroe (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **19**: 1501-1510.
- Gries, G., H. L. McBrien, R. Gries, J.-X. Li, G. J. R. Judd, K. N. Slessor, J. H. Borden, R. F. Smith, M. Christie, J. T. Troubridge, P. D. C. Wimalaratne and E. W. Underhill, 1993(d). (E4,E10)-Dodecadienyl acetate: novel sex pheromone component of tentiform leafminer, *Phyllonorycter mespilella* (Hübner) (Lepidoptera: Gracillariidae). *J. Chem. Ecol.*, **19**: 1789-1798.
- Gries, G., J.-X. Li, R. Gries, W. W. Bowers, R. J. West, P. D. C. Wimalaratne, G. Khaskin, G. G. S. King and K. N. Slessor, 1994(a). (E)-11,13-Tetradecadienal: major sex pheromone component of the eastern blackheaded budworm, *Accleris variana* (Fern.) (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **20**: 1-8.
- Gries, R., G. Gries, J.-X. Li, C. T. Maier, C. R. Lemmon and K. N. Slessor, 1994(b). Sex pheromone components of the spring hemlock looper, *Lambdina athasaria* (Walker) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **20**: 2501-2511.
- Gries, G., R. Gries, G. Khaskin, K. N. Slessor, G. G. Grant, J. Liska and P. Kapitola, 1996. Specificity of nun and gypsy moth sexual communication through multiple-component pheromone blends. *Naturwissenschaften*, **83**: 382-385.
- Gries, G., K. N. Slessor, R. Gries, G. Khaskin, P. D. C. Wimalaratne, T. G. Gray, G. G. Grant, A. S. Tracey and M. Hulme, 1997(a). (Z)6, (E)8-Heneicosadien-11-one: synergistic sex pheromone component of Douglas-fir tussock moth, *Orgyia pseudotsugata* (McDunnough) (Lepidoptera: Lymantriidae). *J. Chem. Ecol.*, **23**(1): 19-34.
- Gries, R., G. Gries, G. G. S. King and C. T. Maier, 1997(b). Sex pheromone components of the apple leafminer, *Lyonetia prunifoliella*. *J. Chem. Ecol.*, **23**(4): 1119-1130.
- Gries, R., D. Holden, G. Gries, P. D. C. Wimalaratne, K. N. Slessor and C. Saunders, 1997(c). 3Z-cis-6,7-cis-9,10-Di-epoxy-heneicosene: Novel class of Lepidopteran pheromone. *Naturwissenschaften*, **84**: 219-221.
- Gries, R., E. Dunkelblum, G. Gries, F. Baldilla, C. Hernandez, F. Alvarez, A. Perez, J. Velasco and A. C. Oehlschlager, 1998. Sex pheromone components of *Diatraea considerata* (Heinrich) (Lep., Pyralidae). *J. Appl. Entomol.*, **122**: 265-268.
- Gries, G., J. Clearwater, R. Gries, G. Khaskin, S. King and P. Schaefer

- 1999(a). Synthetic sex pheromone components of white-spotted tussockmoth, *Orgyia thyellina*. *J. Chem. Ecol.*, **25**(5): 1091-1104.
- Gries, G., R. Gries, P. W. Schaefer, T. Gotoh and Y. Higashiura, 1999(b). Sex pheromone components of pink gypsy moth, *Lymantria mathura*. *Naturwissenschaften*, **86**: 235-238.
- Gries, G., P. W. Schaefer, G. Khaskin, R. Hahn, R. Gries and J.-T. Chao, 1999(c). Sex pheromone components of Casuarina moth, *Lymantria xylina*. *J. Chem. Ecol.*, **25**(11): 2535-2545.
- Gries, G., P. W. Schaefer, R. Gries, J. Liska and T. Gotoh, 2001. Reproductive character displacement in *Lymantria monacha* from northern Japan?. *J. Chem. Ecol.*, **27**(6): 1163-1176.
- Gries, G., P. W. Schaefer, R. Gries, Y.-B. Fan, Y. Higashiura and B. Tanaka, 2002. 2-Methyl-(Z)-7-octadecene: sex pheromone of allopatric *Lymantria lucescens* and *L. serva*. *J. Chem. Ecol.*, **28**(3): 469-478.
- Gries, R., G. Khaskin, E. Khaskin, J. L. Foltz, P. W. Schaefer and G. Gries, 2003. Enantiomers of (Z,Z)-6,9-heneicosadien-11-ol: sex pheromone components of *Orgyia detrita*. *J. Chem. Ecol.*, **29**(10): 2201-2212.
- Gries, R., A. Reckziegel, H. Bogenschütz, H.-G. Kontzog, C. Schlegel, W. Francke, J. G. Millar and G. Gries, 2004. (Z,Z)-11,13-Hexadecadienyl acetate and (Z,E)-11,13,15-hexadecatrienyl acetate: synergistic sex pheromone components of oak processionary moth, *Thaumetopoea processionea* (Lepidoptera: Thaumetopoeidae). *Chemoecology*, **14**(2): 95-100.
- Gries, R., G. Khaskin, P. W. Schaefer, R. Hahn, T. Gotoh and G. Gries, 2005(a). (7R,8S)-cis-7,8-Epoxy-2-methyl-octadec-17-ene: a novel trace component from the sex pheromone gland of gypsy moth, *Lymantria dispar*. *J. Chem. Ecol.* **31**: 49-62.
- Gries, R., G. Khaskin, J. Cleawater, D. Hasman, P. W. Schaefer, E. Khaskin, O. Miroshnychenko, G. Hosking and G. Gries, 2005(b). (Z,Z)-6,9-Heneicosadien-11-one: major sex pheromone component of painted apple moth, *Teia anartoides*. *J. Chem. Ecol.*, **31**(3): 603-620.
- Gries, R., G. Khaskin, T. Gotoh, P. W. Schaefer and G. Gries, 2005(c). (7Z,9E)-2-Methyl-7,9-octadecadiene: a sex pheromone component of *Lymantria bantaizana*. *J. Chem. Ecol.*, **31**(4): 879-891.
- Gries, R., G. Gries, P. W. Schäfer, H. J. S. Yoo and M. Greaves, 2005(d). (Z,E)-6,8-Heneicosadien-11-one: major sex-pheromone component

- of *Orgyia vetusta* (Lepidoptera: Lymantriidae). *Can. Entomol.*, **137**: 471-475.
- Gries, R., G. Khaskin, Z.-X. Tan, B.-G. Zhao, G. G. S. King, A. Miroshnychenko, G.-Q. Lin, M. Rhainds and G. Gries, 2006(a). (1*S*)-1-Ethyl-2-methylpropyl 3,13-dimethylpentadecanoate: major sex pheromone component of Paulownia bagworm, *Clania variegata*. *J. Chem. Ecol.*, **32** (8): 1673-1685.
- Gries, R., G. Khaskin, H. Daroogheh, C. Mart, S. Karadag, M. Kubilay Er, R. Britton and G. Gries, 2006(b). (2*S*,12*Z*)-2-Acetoxy-12-heptadecene: Major sex pheromone component of pistachio twig borer, *Kermaniapistaciella*. *J. Chem. Ecol.*, **32** (12): 2667-2677.
- Gries, R., P. W. Schaefer, R. Hahn, G. Khaskin, G. Ramaseshiah, B. Singh, G. K. Hehar and G. Gries, 2007. Sex pheromone components of Indian gypsy moth, *Lymantria obfuscata*. *J. Chem. Ecol.*, **33** (9): 1774-1786.
- Grodner, J., K. Świech, M. Jakubowska and J. Bocianowski, 2018. Attraction of moths of two Noctuidae species to field traps baited with a mixture of two to three homologous acetates in Poland. *J. Econ. Entomol.*, **111**: 1664-1673.
- Groot, A. T., S. Nojima, J. J. Heath, B. Ammagarahalli, M. van Wijk, A. Claßen, R. G. Santangelo, J. Lopez and C. Schal, 2018. Alcohol contributes to attraction of *Heliothis* (= *Chloridea*) *virescens* males to females. *J. Chem. Ecol.*, **44**: 621-630.
- Gu, P., C.-L. Peng, J. Li, C.-H. Feng, P. Li and Y.-J. Du, 2013. Optimization and field evaluation of the sex pheromone of *Trichophysetis cretacea* (Lepidoptera, Crambidae) for pest monitoring. *J. Appl. Entomol.*, **137**: 462-468.
- Guerin, P. M., H. Arn, C. Blaser and M. Lettéré, 1983. *Z,E*-8,10-dodecadien-1-ol, attractant for male *Pammene rhediella*. *Entomol. Exp. Appl.*, **33**: 346-347.
- Guerin, P. M., W. Baltensweiler, H. Arn and H.-R. Buser, 1984. Host race pheromone polymorphism in the larch budmoth. *Can. Entomol.*, **40**: 892-894.
- Guerin, P. M., H. Arn, H. R. Buser and P. J. Charmillot, 1986(a). Sex pheromone of *Adoxophyes orana*: additional components and variability in ratio of (*Z*)-9- and (*Z*)-11-tetradecenyl acetate. *J. Chem. Ecol.*, **12**: 763-772.
- Guerin, P. M., H. Arn, H. R. Buser, P. Charmillot, M. Tóth and G. Szira'ki, 1986(b). Sex pheromone of *Grapholita funebrana*. Occurrence of

- Z-8- and Z-10-tetradecenyl acetate as secondary components. *J. Chem. Ecol.*, **12**: 1361-1368.
- Guerin, P. M., H. R. Buser, M. Tóth, E. Höbaus, A. Schmid and H. Arn, 1986(c). Sex pheromone of *Sparganothis pilleriana*: E- and Z-11-tetradecenyl acetates as essential components. *Entomol. Exp. Appl.*, **40**: 137-140.
- Guerrero, A, F. Camps, J. Coll, M. Riba, J. Einhorn, C. Descoins and J. Y. Lallemand, 1981. Identification of a potential sex pheromone of the processionary moth, *Thaumetopoea pityocampa* (Lepidoptera, Notodontidae). *Tetrahedron Lett.*, **22**: 2013-2016.
- Guerrero, A., R. Canela, M. Riba, M. Eizaguirre, R. Albajes and V. Sarto, 1986. Asex attractant females of the noctuid moth *Athetis hospes*. *Entomol. Exp. Appl.*, **42**: 97-100.
- Guleria, N., S. M. Nebapure, P. D. K. Jayanthi, S. B. Suby and M. G. Deeksha, 2023. Electrophysiological and behavioral responses of spotted stem borer, *Chilopartellus*, to sex pheromone components and their blends. *J. Chem. Ecol.*, **49**: 155-163.
- Guo, G., Z. Li, G. Lin, P. Wu, X. Zhang, H. Liu and X. Yang, 1990(a). Identification and synthesis of insect pheromone. XXVIII. Sex pheromone of poplar pole clearwing moth, *Sphecia siningensis* Hsu. *J. Chem. Ecol.*, **16**: 819-826.
- Guo, G.-Z., H.-Q. Liu, G.-Q. Lin, C.-H. Wu, L.-Z. Meng and D.-M. Chen, 1990(b). Insect sex pheromone of vine tree borer *Paranthrene regalis*. *Youji Huaxue*, **10**: 504-506.
- Guo, Q., S. Yang, F. Zhuo, W. Xiang, C. Feng, S. Zhang and Y. Du, 2022. Effect of age, daily rhythm and mating on pheromone titres and trapping of *Glyphodes pyloalis* (Lepidoptera, Crambidae). *J. Appl. Entomol.*, **146**: 875-884.
- Hai, T. V., L. V. Vang, P. K. Son, S. Inomata and T. Ando, 2002. Sex attractants for moths of Vietnam: field attraction by synthetic lures baited with known lepidopteran pheromones. *J. Chem. Ecol.*, **28**(7): 1473-1481.
- Hall, D. R. and J. S. Read, 1979. Sex attractants for two zygaenid moths. *J. Entomol. Soc. South Afr.*, **42**: 115-119.
- Hall, D. R., P. S. Beever, R. Lester and B. F. Nesbitt, 1980. (E,E)-10,12-Hexadecadienal: a component of the female sex pheromone of the spiny bollworm, *Earias insulana* (Boisd.) (Lepidoptera: Noctuidae). *Experientia*, **36**: 152-154.
- Hall, D. R., P. S. Beever, A. Cork, R. Lester, B. F. Nesbitt, G. K.

- C. Nyirenda, D. D. N. Phiri, B. W. Blair and J. Tannock, 1981. The female sex pheromone of the maize stalk-borer *Busseola fusca* (Fuller) (Lepidoptera: Noctuidae): identification and initial field trials. *Zimbabwe. J. Agric. Res.*, **19**: 111-122.
- Hall, D. R., P. S. Beevor, A. Cork, B. F. Nesbitt and E. A. S. La Croix, 1984. (Z)-8-Dodecenyl acetate: the major component of the female sex pheromone of *Cryptophlebia batrachopa*, a tortricid pest of macadamia in Malawi. *Entomol. Exp. Appl.*, **35**: 33-36.
- Hall, D. R., A. Cork, R. Lester, B. F. Nesbitt and P. Zagatti, 1987. Sex pheromone of rice moth, *Corcyra cephalonica* Stainton. II. Identification and role of female pheromone. *J. Chem. Ecol.*, **13**: 1575-1589.
- Hall, D. R., P. S. Beevor, D. G. Campion, D. J. Chamberlain, A. Cork, R. D. White, A. Almestare and T. J. Henneberry, 1992. Nitrate esters: Novel sex pheromone components of the cotton leafperforator, *Bucculatrix thurberiella* Busck. (Lepidoptera: Lyonetiidae). *Tetrahedron Lett.*, **33**: 4811-4814.
- Hall, D. R., P. S. Beevor, D. G. Campon, D. J. Chamberlain, A. Cork, R. White, A. Almestare, T. J. Henneberry, Nandagopal, V. Wightman, J. A. and G. V. R. Rao, 1993. Identification and synthesis of new pheromones. *Bulletin OILB SROP*, **16**: 1-9.
- Hall, D. R., D. Farman, J. C. Domínguez and J. A. Pajares, 2017. Female sex pheromone of the cone moth, *Dioryctria mendacella*: investigation of synergism between Type I and Type II pheromone components. *J. Chem. Ecol.*, **43**: 433-442.
- Halperin, J. and Y. Golan, 1982. Attraction of males of *Thaumetopoea wilkinsoni* Tams to the sex pheromone of *Th. pityocampa* Schiff. *Phytoparasitica*, **10**: 140-141.
- Hamstead, E. O., R. E. Dolphin and W. Roelofs, 1972. Field test of virgin female and synthetic sex-lure traps for male redbanded leafrollers. *Environ. Entomol.*, **1**: 488-489.
- Han, G.-B., R.-Y. Ma, J.-W. Du and C. Hu, 1999. The sex pheromone communication system of adult jujube leafroller *Ancylis sativa* Liu. *Acta Entomol. Sinica (Kunchong Xuebao)*, **42**: 25-30.
- Han, K.-S., J.-K. Jung,, K.-H. Choi,, S.-W. Lee,, and K.-S. Boo, 2000. Sex pheromone composition and male trapping of the peach fruit moth, *Carposina sasakii* (Matsumura) (Lepidoptera: Carposinidae) in Korea. *J. Asia Pacific Entomol.*, **3**: 83-88.
- Han, K.-S., J.-K. Jung, K.-H. Choi, S.-W. Lee and K.-S. Boo, 2001. Sex pheromone composition and male trapping of the oriental fruit

- moth, *Grapholita molesta* (Lepidoptera: Tortricidae) in Korea. *J. Asia Pacific Entomol.*, **4**: 31-35.
- Han, B., G. Huang, S. Han and M. Wang, 2024. Development of an efficient attractant of *Ectropis grisescens* (Lepidoptera: Geometridae) and evaluation on its control effect in tea plantations. *Acta Ecol. Sinica (Shengtai Xuebao)*, **44**: 2597-2608.
- Hansen, K., 1984. Discrimination and production of disparlure enantiomers by the gypsymoth and the nun moth. *Physiol. Entomol.*, **9**: 9-18.
- Hansson, B. S., M. Tóth, C. Löfstedt, G. Szöcs, M. Subchev and J. Löfqvist, 1990(a). Pheromone variation among eastern European and a western Asian population of the turnip moth *Agrotis segetum*. *J. Chem. Ecol.*, **16**: 1611-1622.
- Hansson, B. S., G. Szöcs, F. Schmidt, W. Francke, C. Löfstedt and M. Tóth, 1990(b). Electrophysiological and chemical analysis of sex pheromone communication system of the mottled umber, *Erannis defoliaria* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **16**: 1887-1897.
- Hansson, B. S., A. Blackwell, E. Hallberg and J. Löfqvist, 1995. Physiological and morphological characteristics of the sex pheromone detecting system in male corn stem borers, *Chilo partellus* (Lepidoptera: Pyralidae). *J. Insect Physiol.*, **41**(2): 171-178.
- Hanula, J. L., C. W. Berisford and G. L. DeBarr, 1984. Pheromone cross-attraction and inhibition among four coneworms, *Dioryctria* spp. (Lepidoptera: Pyralidae) in a loblolly pine seed orchard. *Environ. Entomol.*, **13**: 1298-1301.
- Harris, M. K., A. A. A. Fu, H. Nunez, E. Aranda-Herera, J. A. Moreira, J. S. McElfresh and J. G. Millar, 2008. A new pheromone race of *Acrobasis nuxvorella* (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **101**: 769-776.
- Hathaway, D. O. and G. Tamaki, 1981. Another olethreutine, *Phaneta latens* (Lepidoptera: Tortricidae), attracted to the sex pheromone of the codling moth. *J. Entomol. Soc. Brit. Columbia*, **78**: 24-27.
- Hattori, M., S. Wakamura, K. Igita, K. Yasuda and Tridjaka, 2001. Comparison of the characteristics and sex pheromone of *Etiella behrii* (Zeller), a newly identified pod borer of soybean in Indonesia, with *E. zinckenella* (Treit.) *Jpn. Int. Res. Quart.*, **35**: 19-24.
- Haynes, K. F., 1987. Identification of sex pheromone of calendula

- plumemoth, *Platyptilia williamsii*. *J. Chem. Ecol.*, **13**: 907-916.
- Haynes, K. F., 1990. Identification of sex pheromone of bristly cutworm, *Lacinipolia renigera* (Stephens). *J. Chem. Ecol.*, **16**: 2615-2621.
- Haynes, K. F., K. V. Yeargan, J. G. Millar and B. B. Chastain, 1996. Identification of sex pheromone of *Tetanolita mynesalis* (Lepidoptera: Noctuidae), a prey species of bolas spider, *Mastophora hutchinsoni*. *J. Chem. Ecol.*, **22**: 75-89.
- He, Y., H. Xie and S. Chi, 2016. Identification of female main component sex pheromones of *Orthaga achatina* (Lepidoptera: Pyralidae) and their attractiveness to moths. *Agric. Sci. Technol.*, **17**: 920-922.
- Heath, R. R., J. H. Tumlinson, N. C. Leppla, J. R. McLaughlin, B. Dueben, E. Dundulis and R. H. Guy, 1983. Identification of a sex pheromone produced by female velvetbean caterpillar moth. *J. Chem. Ecol.*, **9**: 645-656.
- Heath, R. R. and P. J. Landolt, 1987. (Z)-5-Decen-1-ol acetate: the major sex pheromone component of *Argyrogramma verruca* (Fabricius) (Lepidoptera: Noctuidae: Plusiinae). *Environ. Entomol.*, **16**: 1030-1033.
- Heath, R. R., P. E. A. Teal, N. D. Epsky, B. D. Dueben, S. D. Hight, S. Bloem, J. E. Carpenter, T. J. Weissling, P. E. Kendra, J. Cibrian-Tovar and K. A. Bloem, 2006. Pheromone-based attractant for males of *Cactoblastis cactorum* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **35**: 1469-1476.
- Hedin, P. A., F. M. Davis, J. C. Dickens, M. L. Burks, T. G. Bird and A. E. Knutson, 1986. Identification of the sex attractant pheromone of the southwestern corn borer *Diatraea grandiosella* Dyar. *J. Chem. Ecol.*, **12**: 2051-2063.
- Hedlin, A. F., J. Weatherston, D. S. Ruth and G. E. Miller, 1983. Chemical lure for male Douglas-fir cone moth, *Barbara colfaxiana* (Lepidoptera: Olethreutidae). *Environ. Entomol.*, **12**: 1751-1753.
- Heikinheimo, O., 1978. Feromonfangst som metod för faunistiska undersökningar. *Norw. J. Entomol.*, **25**: 102.
- Henderson, H. E. and F. L. Warren, 1972. Sex-pheromones, *cis*-dec-5-en-1-yl 3-methylbutanoate as the pheromone from the pine emperor moth (*Nudaurelia cytherea cytherea* Fabr.). *J. Chem. Soc. Chem. Comm.*, 686-687.
- Henderson, H. E., F. L. Warren, O. P. H. Augustyn, B. V. Burger, D. F. Schneider, P. R. Boshoff, H. S. C. Spies and H. Geertsema,

1973. Isolation and structure of the sex-pheromone of the moth, *Nudaurelia cytherea cytherea*. *J. Insect Physiol.*, **19**: 1257-1264.
- Hendricks, D. E., B. A. Leonhardt and T. N. Shaver, 1989. Development of optimized blends of two sex pheromone components impregnated in PVC dispensers for tobacco budworm bait. *Southwestern Entomol.*, **14**: 17-25.
- Hendrikse, A., W. M. Herrebout and C. J. Persoons, 1982. Sex pheromones, attractants and inhibitors in small ermine moths: a comparison of bioassays. *Meded. Fac. Landbouww. Rijksuniv. Gent.*, **47**: 493-502.
- Hendry, L. B., J. Jugovich, L. Roman, M. E. Anderson and R. O. Mumma, 1974. *Cis*-10-tetradecenyl acetate, an attractant component in the sex pheromone of the oak leaf roller moth (*Archips semiferranus* Walker). *Experientia*, **30**: 886-887.
- Hendry, L. B., M. E. Anderson, J. Jugovich, R. O. Mumma, D. Robacker and Z. Kosarych, 1975. Sex pheromone of the oak leaf roller: a complex chemical messenger system identified by mass fragmentography. *Science*, **187**: 355-357.
- Herrebout, W. M., 1978. Sex attractants for *Yponomeuta* species. *Neth. J. Zool.*, **28**: 276.
- Herrera, H., W. Barros-Parada, M. F. Flores, W. Francke, E. Fuentes-Contreras, M. Rodriguez, F. Santis, P. H. G. Zarbin and J. Bergmann, 2016. Identification of a novel moth sex pheromone component from *Chilecomadia valdiviana*. *J. Chem. Ecol.*, **42**: 908-918.
- Hill, A., R. Cardé, A. Comeau, W. Bode and W. Roelofs, 1974. Sex pheromone of the tufted apple bud moth (*Platynota idaeusalis*). *Environ. Entomol.*, **3**: 249-252.
- Hill, A. S. and W. L. Roelofs, 1975(a). Sex pheromone components of the omnivorous leafroller moth, *Platynota stultana*. *J. Chem. Ecol.*, **1**: 91-99.
- Hill, A. S., R. T. Cardé, H. Kido and W. L. Roelofs, 1975(b). Sex pheromone of the orange tortrix moth, *Argyrotaenia citrana* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **1**: 215-224.
- Hill, A. S., C. W. Berisford, U. E. Brady and W. L. Roelofs, 1976. Sex pheromone of the pitch pine tip moth, *Rhyacionia rigidana*. *Environ. Entomol.*, **5**: 959-960.
- Hill, A. S., R. T. Cardé, W. M. Bode and W. L. Roelofs, 1977. Sex pheromone components of the variegated leafroller moth, *Platynota*

- flavedana. *J. Chem. Ecol.*, **3**: 369-376.
- Hill, A. S. and W. L. Roelofs, 1978. Two sex attractants for male speckled green fruitworm moths, *Orthosia hibisci* Guenee (Lepidoptera: Noctuidae). *J. New York Entomol. Soc.*, **86**: 296.
- Hill, A. S. and W. L. Roelofs, 1979(a). Sex pheromone components of the obliquebanded leafroller moth, *Choristoneura rosaceana*. *J. Chem. Ecol.*, **5**: 3-11.
- Hill, A. S., R. W. Rings, S. R. Swier and W. L. Roelofs, 1979(b). Sex pheromone of the black cutworm moth, *Agrotis ipsilon*. *J. Chem. Ecol.*, **5**: 439-457.
- Hill, A. S. and W. L. Roelofs, 1980. A female-produced sex pheromone component and attractant for males in the armyworm moth, *Pseudaletia unipuncta*. *Environ. Entomol.*, **9**: 408-411.
- Hill, A. S., C. W. Berisford, U. E. Brady and W. L. Roelofs, 1981(a). Nantucket pine tip moth, *Rhyacionia frustrana*: identification of two sex pheromone components. *J. Chem. Ecol.*, **7**: 517-528.
- Hill, A. S. and W. L. Roelofs, 1981(b). Sex pheromone of the saltmarsh caterpillar moth, *Estigmene acrea*. *J. Chem. Ecol.*, **7**: 655-668.
- Hill, A. S., B. G. Kovalev, L. N. Nikolaeva and W. L. Roelofs, 1982. Sex pheromone of the fall webworm moth, *Hyphantria cunea*. *J. Chem. Ecol.*, **8**: 383-396.
- Hillier, N. K., P. L. Dixon, W. D. Seabrook and D. J. Larson, 2002. Field testing of synthetic attractants for male *Grapholita libertina* (Lepidoptera: Tortricidae). *Can. Entomol.*, **134**: 657-665.
- Himeno, K. and H. Honda, 1992. (E,Z)- and (E,E)-10,12-Hexadecadienals, major components of female sex pheromone of the cotton leaf-roller, *Notarcha derogata* (Fabricius) (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **27**: 507-515.
- Hirai, Y., H. Kimura, K. Kawasaki and Y. Tamaki, 1978. (Z)-11-Hexadecenyl acetate: a sex-pheromone component of the cabbage armyworm moth, *Mamestra brassicae* Linné. *Appl. Entomol. Zool.*, **13**: 136-137.
- Hirano, C., H. Muramoto and M. Horiike, 1976. Sex pheromone produced by female sweet potato leaf folder moth. *Naturwissenschaften*, **63**: 439.
- Ho, H.-Y., Y.-T. Tao, R.-S. Tsai, Y.-L. Wu, H.-K. Tseng and Y.-S. Chow, 1996. Isolation, identification, and synthesis of sex pheromone

- components of female tea cluster caterpillar, *Andraca bipunctata* Walker (Lepidoptera: Bombycidae) in Taiwan. *J. Chem. Ecol.*, **22**: 271-285.
- Ho, H. Y., R. S. Tsai, E. L. Hsu, Y. S. Chow and R. Kou, 2002. Investigation of possible sex pheromone components of female loreyi leafworm *Acantholeucania loreyi* (Duponchel) (Lepidoptera: Noctuidae) in Taiwan. *Zool. Studies*, **41**(2): 188-193.
- Hobosyan, N. G., K. V. Balyan, L. A. Movsisyan, V. S. Hovsepyan, H. B. Sargsyan, H. R. Pogosyan, H. R. Harutyunyan, A. A. Manvelyan, M. H. Ghazaryan, S. S. Asatryan and H. L. Terlemezyan, 2025. Novel synthetic strategy and field testing tomato pinworm pheromone in Armenia. *Chem. Biodivers.*, **22**: e202403455.
- Hoddle, M. S., J. G. Millar, C. D. Hoddle, Y. Zou and J. S. McElfresh, 2009. Synthesis and field evaluation of the sex pheromone of *Stenoma catenifer* (Lepidoptera: Elachistidae). *J. Econ. Entomol.*, **102**: 1460-1467.
- Hoffmann, M. P., L. M. McDonough and J. B. Bailey, 1983. Field test of the sex pheromone of *Amorbia cuneana* (Walsingham) (Lepidoptera: Tortricidae). *Environ. Entomol.*, **12**(5): 1387-1390.
- Holden, D. G., 2000. Sex pheromone components of satin moth, *Leucoma salicis* (L.) (Lepidoptera: Lymantriidae). Master Thesis, Simon Fraser University, Burnaby, B.C., Canada.
- Honda, H., T. Ando, Y. Ogura, T. Teramine and R. Ueda, 1990. Identification of the sex pheromone of the mulberry pyralid, *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **25**: 265-272.
- Honda, H., K. Himeno and Y. Yoshiyasu, 1994. Chemotaxonomy of the cotton leaf-roller (Lepidoptera: Pyralidae) in Japan with special reference to differences in sex pheromones. *Appl. Entomol. Zool.*, **29**(3): 323-330.
- Honda, H. and T. Kimura, 2004. (E)-10-Hexadecenal as a sex attractant of *Pleuroptia chloropahanta* (Butler) (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **48**: 57-60. (in Japanese)
- Honda, H., R. Yamasaki, Y. Sumiuchi, T. Uehara, S. Matsuyama, T. Ando and H. Naka, 2015. Hybrid sex pheromones of the hibiscus flower-bud borer, *Rehimena surusalis*. *J. Chem. Ecol.*, **41**: 1043-1049.
- Honma, K., K. Kawasaki and Y. Tamaki, 1978. Sex-pheromone activities of synthetic 7-alken-11-ones for male peach fruit moths, *Carposina niponensis* Walsingham (Lepidoptera: Carposinidae). *Jap. J. Appl.*

Entomol. Zool., **22**: 87-91.

- Horák, A. and I. Hrdý, 1988. Attractiveness of (E)-10-dodecen-1-yl acetate for the pea moth, *Cydia nigricana*: modification by codlemone, codlemone acetate and dodecyl acetate. *Acta Entomol. Bohemoslov.*, **85**: 174-181.
- Horak, M., C. P. Whittle, T. E. Bellas and E. R. Rumbo, 1988. Pheromone gland components of some Australian tortricids in relation to their taxonomy. *J. Chem. Ecol.*, **14**: 1163-1175.
- Hrdý, I., J. Marek and F. Krámpal, 1979. Sexual pheromone activity of 8-dodecenyl and 11-tetradecenyl acetates for males of several lepidopteran species in field trials. *Acta Entomol. Bohemoslov.*, **76**: 65-84.
- Hrdý, I., J. Liska, J. Vrkoc and R. Hochmut, 1989. New records on sex attractants for males and faunistic comments on moth (Lepidoptera) from Czechoslovakia. *Acta Entomol. Bohemoslov.*, **86**: 252-268.
- Hrudová, E, 2003. The presence of non-target lepidopteran species in pheromone traps for fruit tortricid moths. *Plant Protect. Sci.*, **39**: 126-131.
- Hsiao, S. N. 1990. Bioassay of sex pheromone of the smaller tea tortrix, *Adoxophyes* sp. (Lepidoptera: Tortricidae). *Chin. J. Entomol.*, **10**: 443-450.
- Hu, Wenli and J. Du, 2005. Sex pheromone secondary components of Indian meal moth *Plodia interpunctella* in China. *Yingyong Shengtai Xuebao*, **16**(9): 1751-1755. [[SciFinder](#), 2005:1285000]
- Hu, D.-H., J. He, Y.-W. Zhou, J.-T. Feng and X. Zhang, 2012. Synthesis and field evaluation of the sex pheromone analogues to soybean pod borer *Leguminivora glycinivorella*. *Molecules*, **17**: 12140-12150.
- Huang, Y.-P., S. Tatsuki, C.-G. Kim, S. Hoshizaki, Y. Yoshiyasu, H. Honda and Y. Ishikawa 1997. Identification of sex pheromone of adzuki bean borer, *Ostrinia sacculalis*. *J. Chem. Ecol.*, **23**(12): 2791-2802.
- Huang, Y., T. Takanashi, S. Hoshizaki, S. Tatsuki, H. Honda, Y. Yoshiyasu and Y. Ishikawa 1998(a). Geographic variation in sex pheromone of Asian corn borer, *Ostrinia furnacalis*, in Japan. *J. Chem. Ecol.*, **24**(12): 2079-2088.
- Huang, Y., S. Tatsuki, C.-G. Kim, S. Hoshizaki and Y. Ishikawa, 1998(b). Identification of the sex pheromone of *Ostrinia*

- palustralis*. *Entomol. Exp. Appl.*, **86**: 313-318.
- Huang, Y., H. Honda, Y. Yoshiyasu, S. Hoshizaki, S. Tatsuki and Y. Ishikawa, 1998(c). Sex pheromone of the butterbur borer, *Ostrinia zaguliaevi*. *Entomol. Exp. Appl.*, **89**: 281-287.
- Huang, Y., T. Takanashi, S. Hoshizaki, S. Tatsuki and Y. Ishikawa, 2002. Female sex pheromone polymorphism in adzuki bean borer, *Ostrinia scapulalis*, is similar to that in European corn borer, *O. nubilalis*. *J. Chem. Ecol.*, **28**(3): 533-539.
- Huang, J., L. J. Gut and M. J. Grieshop, 2022. Attraction of *Choristoneura rosaceana* to pheromone blends in ratios produced in glands or emitted by females. *J. Appl. Entomol.*, **146**: 734-742.
- Hummel, H. E., L. K. Gaston, H. H. Shorey, R. S. Kaae, K. J. Byrne and R. M. Silverstein, 1973. Clarification of the chemical status of the pink bollworm sex pheromone. *Science*, **181**: 873-875.
- Hung, C. C., J. S. Hwang, M. D. Hung, Y. P. Yen and R. F. Hou, 2001. Isolation, identification and field tests of the sex pheromone of the carambola fruit borer, *Eucosma notanthes*. *J. Chem. Ecol.*, **27** (9): 1855-1866.
- Hung, C. C., M. D. Hung and W. L. Wang, 2006. Development of formulations of sex attractant of *Conopomorpha sinensis* on *Litchi chinesis*. *Zhiwu Baohu Xuehui Huikan*, **48**(3): 189-202. [SciFinder, 2007:3272]
- Hussain, A., S. Hladun, M. Vincent, T. J. Wist, N. K. Hillier and B. A. Mori, 2023. Development of a pheromone monitoring system for the goosefoot groundling moth, *Scrobipalpa atriplicella* (von Röslerstamm) in quinoa, *Chenopodium quinoa* (Willdenow). *Crop Prot.*, **165**: 106166.
- Hwang, J. S., C. C. Hung and R. F. Hou, 1996(a). A bioassay method of the sex pheromone of the carambola fruit borer, *Eucosma notanthes* Meyrick. *Zhiwu Baohu Xuehui Huikan*, **38**(2): 119-127. [Chem. Abst., 126: 2969s]
- Hwang, J. S., C. C. Hung, Y. P. Yen and C. N. Chen, 1996(b). Sex attractant and trap designs for trapping litchi fruit borer, *Conopomorpha sinensis* Bradley. *Zhiwu Baohu Xuehui Huikan (Plant Protection Bulletin)*, **38**(2): 129-136.
- Ichikawa, A., T. Yasuda and S. Wakamura, 1995. Absolute configuration of sex pheromone for tea tussock moth, *Euproctis pseudoconspersa* (Strand) via synthesis of (R)- and (S)-10,14-dimethyl-1-pentadecyl isobutyrylates. *J. Chem. Ecol.*, **21**: 627-634.

- Inomata, S., M. Komoda, H. Watanabe, M. Nomura and T. Ando, 2000. Identification of sex pheromones of *Anadevidia peponis* and *Macdunnoughia confusa*, and field tests of their role in reproductive isolation of closely related Plusiinae moths. *J. Chem. Ecol.*, **26**: 443-454.
- Inomata, S., S. Tsuchiya, K. Ikeda, O. Saito and T. Ando, 2002. Identification of sex pheromone components secreted by female moths of *Peridroma saucia* (Noctuidae: Noctuinae). *Biosci. Biotechnol. Biochem.*, **66**(11): 2461-2464.
- Inomata, S., A. Watanabe, M. Nomura and T. Ando, 2005. Mating communication systems of four Plusiinae species distributed in Japan: identification of the sex pheromones and field evaluation. *J. Chem. Ecol.*, **31**(6): 1429-1442.
- Ishikawa, Y., T. Takanashi and Y. Huang, 1999(a). Comparative studies on the sex pheromones of *Ostrinia* spp. In Japan: the burdock bore, *Ostrinia zealis*. *Chemoecology*, **9**: 25-32.
- Ishikawa, Y., T. Takanashi, C.-G. Kim, S. Hoshizaki, S. Tatsuki and Y. Huang, 1999(b). *Ostrinia* spp. in Japan: their host plants and sex pheromones. *Entomol. Exp. Appl.*, **91**: 237-244.
- Islam, MD. A., M. Yamamoto, M. Sugie, H. Naka, J. Tabata, Y. Arita and T. Ando, 2007. Synthesis and characterization of 2,13- and 3,13-octadecadienals for the identification of the sex pheromone secreted by a clearwing moth. *J. Chem. Ecol.*, **33**(9): 1763-1773.
- Islam, MD. A., R. Yamakawa, N. D. Do, N. Numakura, T. Suzuki and T. Ando, 2009. Instrumental analysis of terminal-conjugated dienes for reexamination of the sex pheromone secreted by a nettle moth, *Parasa lepida lepida*. *Biosci. Biotechnol. Biochem.*, **73**: 1156-1162.
- Iwaki, S., S. Marumo, T. Saito, M. Yamada and K. Katagiri, 1974. Synthesis and activity of optically active disparlure. *J. Amer. Chem. Soc.*, **96**: 7842-7844.
- Jaastad, G., M. Bengtsson, P. Anderson, S. Kobro, G. Knudsen and P. Witzgall, 2002. Sex pheromone of apple fruit moth *Argyresthia conjugella* (Lepidoptera: Argyresthiidae). *Agric. Forest Entomol.*, **4**: 233-236.
- Jacobson, M., R. E. Redfern, W. A. Jones, M. H. Aldridge, 1970. Sex pheromones of the southern armyworm moth: isolation, identification, and synthesis. *Science*, **170**: 542-544.

- Jacobson, M. and D. T. Jennings, 1978. Attraction of *Rhyacionia neomexicana* (Dyar) to synthetic pheromones. *J. Environ. Sci. Health*, **A 13**: 429-443.
- Jain, S. C., D. E. Dussourd, W. E. Conner, T. Eisner, A. Guerrero and J. Meinwald, 1983. Polyene pheromone components from an arctiid moth (*Utetheisa ornatrix*): characterization and synthesis. *J. Org. Chem.*, **48**: 2266-2270.
- James, D. G., A. Cosse, L. C. Wright and J. Perez, 2001. Pheromone trapping of *Synanthedon tipuliformis* (Lepidoptera: Sesiidae) in Washington red currants. *Environ. Entomol.*, **30**: 663-666.
- Jennings, D. T. and M.W. Houseweart, 1983. Field attractiveness of (*E*)- and (*Z*)-11-tetradecenal pheromone blends to male spruce budwormmoths, *Choristoneura fumiferana* (Clemens). *J. Chem. Ecol.*, **9**: 1327-1332.
- Jham, G. N., A. A. da Silva, E. R. Lima and P. Viana, 2005. Identification (GC and GC-MS) of unsaturated acetates in *Elasmopalpus lignosellus* and their biological activity (GC-EAD and EAG). *J. Sep. Sci.*, **28**: 281-285.
- Jham, G. N., A. A. da Silva, E. R. Lima and P. A. Viana, 2007. Identification of acetates in *Elasmopalpus lignosellus* pheromone glands using a newly created mass spectral database and Kóvats retention indices. *Quim. Nova*, **30** (4): 916-919.
- Jiang, N.-J., B.-T. Mo, H. Guo, J. Yang, R. Tang and C.-Z. Wang, 2022. Revisiting the sex pheromone of the fall armyworm *Spodoptera frugiperda*, a new invasive pest in South China. *Insect Sci.*, **29**: 865-878.
- Jing, X.-Y, J.-T. Zhang, Y.-Q. Luo, S.-X. Zong, P.-H. Liu and J.-X. Jia, 2010. Identification and field evaluation of the female sex pheromone of the sand *Salix* carpenterworm, *Holcocerus arenicola* Staudinger (Lepidoptera: Cossidae). *Z. Naturforsch.*, **65c**: 403-411.
- Jing, X., G. Li, C. Jia, R. Jin, M. Wang, R. Mu, Y. Bai, Y. Hao and H. Liu, 2025. Identification and field testing of female-produced sex pheromone components of *Dendrolimus suffusus* subspecies *illustratus* De Lajonquière, 1973. *Chem. Biodivers.*, **22**: e00785.
- Jones, R. L. and A. N. Sparks, 1979. (*Z*)-9-Tetradecen-ol acetate. A secondary sex pheromone of the fall armyworm, *Spodoptera frugiperda* (J. E. Smith). *J. Chem. Ecol.*, **5**: 721-725.
- Jordan, T. A., A. Zhang and D. G. Pfeiffer, 2013. Blend chemistry

- and field attraction of commercial sex pheromone lures to grape berry moth (Lepidoptera: Tortricidae), and a nontarget tortricid in vineyards. *Environ. Entomol.*, **42**: 558-563.
- Judd, G. J. R., R. Gries, V. M. Aurelian and G. Gries, 2011. 3Z,13Z-octadecadienyl acetate: sex pheromone of the apple clearwing moth in British Columbia. *Can. Entomol.*, **143**: 236-244.
- Judd, G. J. R., R. Gries, C. Teasdale and G. Gries, 2012. Identification of a sex pheromone component for *Pennisetia marginata* (Lepidoptera: Sesiidae). *Can. Entomol.*, **144**: 798-807.
- Jung, C. R., Y. J. Park and K. S. Boo, 2003. Optimal sex pheromone composition for monitoring *Spodoptera exigua* (Lepidoptera: Noctuidae) in Korea. *J. Asia-Pac. Entomol.* **6**: 175-182.
- Jurb, G. L., Jr. 1973. Catches of *Episimus argutatus* in grape berry moth sex-pheromone traps in Pennsylvania. *J. Econ. Entomol.*, **66**: 1345-1346.
- Kaae, R. S., H. H. Shorey, S. U. McFarland and L. K. Gaston, 1973(a). Sex pheromones of Lepidoptera. XXXVII. Role of sex pheromones and other factors in reproductive isolation among ten species of Noctuidae. *Ann. Entomol. Soc. Am.*, **66**: 444-448.
- Kaae, R. S., H. H. Shorey and L. K. Gaston, 1973(b). Pheromone concentration as a mechanism for reproductive isolation between two lepidopterous species. *Science*, **179**: 487-489.
- Kaissling, K. E., G. Kasang, H. J. Bestmann, W. Stransky and O. Vostrowsky, 1978. A new pheromone of the silkworm moth *Bombyx mori*. Sensory pathway and behavioral effect. *Naturwissenschaften*. **65**: 382-384.
- Kakizaki, M. and H. Sugie, 2003. Sex pheromone of the flax budworm, *Heliothis maritima adaucta* Butler (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **38**(1): 73-78.
- Kalinová, B., A. Minaif and L. Kotera, 1994. Sex pheromone characterization and field trapping of the European corn borer, *Ostrinia nubilalis* (Lepidoptera: Pyralidae), in South Moravia and Slovakia. *Eur. J. Entomol.*, **91**: 197-203.
- Kalinová, B., A. Svatoš, J. Kindl, O. Hovorka, I. Hrdý, J. Kuldova and M. Hoskovec, 2003. Sex pheromone of horse-chestnut leafminer *Cameraria ohridella* and its use in a pheromone-based monitoring system. *J. Chem. Ecol.*, **29**(2): 387-404.
- Kalinová, B., J. Kindl, O. Hovorka, M. Hoskovec and A. Svatoš, 2005. (11Z)-Hexadec-11-enal enhances the attractiveness of

- Diatraea saccharalis* main pheromone component in wind tunnel experiments. *J Appl. Entomol.*, **129** (2): 70-74.
- Kalinová, B., P. Jiros, J. Zd'arek, X. Wen and M. Hoskovec, 2006. GC x GC/TOF MS technique. A new tool in identification of insect pheromones: analysis of the persimmon bark borer sex pheromone gland. *Talanta* **69**: 542-547.
- Kalinová, B., R. R. D. Nascimento, M. Hoskovec, A. L. Mendonça, E. L. Silva, M. R. T. D. Freitas, C. R. Cabral-Jr, C. E. Silva, A. E. G. Sant'Ana and A. Svatoš, 2012. Identification of two components of the female sex pheromone of the sugarcane-borer *Diatraea flavipennella* (Lepidoptera: Crambidae). *J. Appl. Entomol.*, **136**: 203-211.
- Kamm, J. A. and L. M. McDonough, 1979(a). Sex attractant for a plume moth, *Oidaematophorus monodactylus* (L.) (Lepidoptera: Pterophoridae). *Pan-Pacific Entomol.*, **55**: 284-286.
- Kamm, J. A. and L. M. McDonough, 1979(b). Field tests with the sex pheromone of the cranberry girdler. *Environ. Entomol.*, **8**: 773-775.
- Kamm, J. A. and L. M. McDonough, 1980. Synergism of the sex pheromone of the granberry girdler. *Environ. Entomol.*, **9**: 795-797.
- Kamm, J. A., L. M. McDonough and C. L. Smithhisler, 1982. Sex attractant for *Protagrotis obscura*, a pest of grass grown for seed. *Environ. Entomol.*, **11**: 118-120.
- Kamm, J. A. and L. M. McDonough, 1985. Sex attractant for *Diarsia pseudorosaria*, a defoliator of ryegrass (Lepidoptera: Noctuidae). *Pan-Pacific Entomol.*, **61**: 299-302.
- Kamm, J. A., L. M. McDonough and K. E. Rowe, 1989. Effect of dosage and ratio of sex pheromone components on behavior of *Chrysoteuchia topiaria* (Zeller) (Lepidoptera: Pyralidae). *Environ. Entomol.*, **18**: 368-372.
- Kang, S.-K., B.-H. Moon, J.-O. Lee and H.-G. Goh, 1985. A Convergent synthesis of (Z)-13-octadecen-1-yl acetate, the pheromone mimic of the rice leaf folder moth and its biological activity test. *Bull. Korean Chem. Soc.*, **6**: 228-230.
- Kang, S.-K. and K.-Y. Seol, 1995. Sex pheromone of the perilla leaf pyralid moth: isolation and identification. *Bull. Korean Chem. Soc.*, **16**: 688-689.
- Kanno, H., L. P. S. Kuenen, K. A. Klingler, J. G. Millar and R. T. Cardé, 2010. Attractiveness of a four-component pheromone

- blend to male navel orangeworm moths. *J. Chem. Ecol.*, **36**: 584-591.
- Karalius, V., R. Mozūraitis, J. Miatleuski, V. Būda and P. Ivinskis, 2001. Sex attractants for six clearwing and tineid species (Lepidoptera, Sesiidae and Tineidae) from Kazakhstan and Lithuania. *Z. Naturforsch.*, **56c**: 1120-1125.
- Karalius, V., R. Mozūraitis, V. Būda, I. Libikas and A.-K. Borg-Karlson, 2005. Sex attractants for six moth species of the families Brachodidae, Choreutidae and Tineidae from Kazakhstan and Lithuania. *Z. Naturforsch.*, **60c**: 625-631.
- Karalius, V. and V. Būda, 2006. Sex attractants for clearwing moths *Synanthedon mesiaeformis* (Herrich-Schaffer) and *S. flaviventris* (Staudinger) (Lepidoptera, Sesiidae). *Acta Zool. Lit.*, **16**: 198-203.
- Karandinos, M. G., J. H. Tumlinson and T. D. Eichlin, 1977. Field evidence of synergism and inhibition in the Sesiidae sex pheromone system. *J. Chem. Ecol.*, **3**: 57-64.
- Kasang, G., K. E. Kaissling, O. Vostrowsky and H. J. Bestmann, 1978(a). Bombykal, a second pheromone component of the silkworm moth *Bombyx mori* L. *Angew. Chem. Int. Ed. Engl.*, **17**: 60.
- Kasang, G. and D. Schneider, 1978(b). The silkworm moth *Bombyx mori*. Presence of the (*E,E*)-stereoisomer of bombykol in the female pheromone gland. *Naturwissenschaften*, **65**: 337-338.
- Katovich, S. A., P. D. Swedenborg, M. Giblin and E. W. Underhill, 1989. Evidence for (*E,Z*)-8,10-dodecadienyl acetate as the major component of the sex pheromone of the eastern pine seedworm, *Cydia toreuta* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **15**: 581-590.
- Kawahara, A. Y., M. Jones, Q. Jia, S. L. LaPointe and P. A. Stansly, 2013. A synthetic pheromone for *Phyllocnistis citrella* (Lepidoptera: Gracillariidae) attracts multiple leafminer species. *Florida Entomol.*, **96**: 1213-1216.
- Kawazu, K., J. Hasegawa, H. Honda, Y. Ishikawa, S. Wakamura, H. Sugie, H. Kamiwada, T. Kamimuro, Y. Yoshiyasu and S. Tatsuki, 2000. Geographical variation in female sex pheromones of the rice leaf folder moth, *Cnaphalocrocis medinalis*: identification of pheromone components in Japan. *Entomol. Exp. Appl.*, **96**: 103-109.
- Kawazu, K., O. Setokuchi, K. Kohno, K. Takahashi, Y. Yoshiyasu and S. Tatsuki, 2001. Sex pheromone of the rice leaf folder moth, *Cnaphalocrocis medinalis* (Lepidoptera: Crambidae): Synthetic Indian and Philippine blends are not attractive to male *C. medinalis*,

- but are attractive to *C. pilosa* in the South-Western islands n Japan. *Appl. Entomol. Zool.*, **36**: 471-474.
- Kawazu, K., K. Nagata, Z. Zhang, H. Sugie and S. Tatsuki, 2002. Comparison of attractiveness in Japan and China of three synthetic pheromone blends based on geographic variations in the rice leaffolder, *Cnaphalocrocis medinalis* (Lepidoptera: Pyralidae). *Bull. Entomol. Res.*, **92**: 295-299.
- Kawazu, K., H. Honda, S. Nakamura and T. Adati, 2007. Identification of sex pheromone components of the box tree pyralid, *Glyphodes perspectalis*. *J. Chem. Ecol.*, **33**: 1978-1985.
- Kawazu, K., T. Adati, Y. Yosiyasu, K. Sumiarta, W. Susila, P. Sudiarta, H. Purwanto, and S. Tatsuki, 2009. Sex pheromone components of the rice leaffolder, *Cnaphalocrocis medinalis* (Lepidoptera: Crambidae), in Indonesia. *J. Asia Pacific Entomol.*, **12**: 97-99.
- Keaster, A. J., J. A. Grundler, W. S. Craig and M. A. Jackson, 1996. Noctuidmoths and other insects captured in wing-style traps baited with black cutworm (Lepidoptera: Noctuidae) pheromone on offshore oil platforms in the Gulf of Mexico, 1988-1991. *J. Kansas Entomol. Soc.*, **69**: 17-25.
- Keathley, C. P., L. L. Stelinski and S. L. Lapointe, 2013. Attraction of a native Florida leafminer, *Phyllocnistis insignis* (Lepidoptera: Gracillariidae), to pheromone of an invasive citrus leafminer, *P. citrella*: Evidence for mating disruption of a native non-target species. *Florida Entomol.*, **96**: 877-886.
- Kehat, M., S. Greenberg and Y. Tamaki, 1976. Field evaluation of the synthetic sex pheromone, as an attractant for males of the cotton leafworm, *Spodoptera littoralis* (Boisd.), in Israel. *Appl. Entomol. Zool.*, **11**: 45-52.
- Kehat, M., 1977. Field evaluation of purified Z-9,E-11-tetradecadienyl acetate as an attractant for males of *Spodoptera littoralis* (Boisd.) (Lepidoptera, Noctuidae). *Phytoparasitica*, **5**: 35-37.
- Kehat, M., S. Gothilf, E. Dunkelblum and S. Greenberg, 1980. Field evaluation of female sex pheromone components of the cotton bollworm, *Heliothis armigera*. *Entomol. Exp. Appl.*, **27**: 188-193.
- Kehat, M. and E. Dunkelblum, 1990. Behavioral responses of male *Heliothis armigera* (Lepidoptera: Noctuidae) moths in a flight tunnel to combinations of components identified from female sex pheromone glands. *J. Insect Behavior*, **3**: 75-83.
- Kehat, M., A. Eitam, D. Blumberg, E. Dunkelblum, L. Anshelevich and

- D. Gordon, 1992. Sex pheromone traps for detecting and monitoring the raisin moth, *Cadra figulilella*, in date palm plantations. *Phytoparasitica*, **20**: 99-106.
- Khrimian, A., D. R. Lance, M. Schwarz, B. A. Leonhardt and V. Mastro, 2008. Sex pheromone of browntail moth, *Euproctis chrysorrhea* (L.): synthesis and field development. *J. Agric. Food Chem.*, **56**: 2452-2456.
- Kim, J. and I.-K. Park, 2013. Female sex pheromone components of the box tree pyralid, *Glyphodes perspectalis*, in Korea: Field test and development of film-type lure. *J. Asia-Pac. Entomol.*: **16**, 473-477.
- Kim, J., K.-S. Cho, C. Y. Yang, C. G. Park, 2014(a). Identification and field evaluation of the female sex pheromone of *Stathmopoda masinsissa* in Korea. *Chemoecology*, **24**: 253-259.
- Kim, J., H. S. Roh, B. K. Byun and C. G. Park, 2014(b). A co-attractant mixture of (Z)-11-tetradecenyl acetate and (Z)-9-dodecenyl acetate for *Neocalyptis angustilineata* and *Homona magnanima*. *J. Asia-Pac. Entomol.*, **17**: 659-661.
- Kim, J., B.-K. Byun, H.-W. Oh, S. A. Jang and C. G. Park, 2016. *Athetis dissimilis* (Lepidoptera: Noctuidae) is attracted to the sex pheromone of *Euzophera batangensis* (Lepidoptera: Pyralidae). *J. Asia-Pac. Entomol.*, **19**: 841-845.
- Kim, J. and S. Lee, 2023. Identification and field bioassay of *Rusicada privata* (Lepidoptera: Erebidae) sex pheromone. *J. Agric. Food Chem.*, **71**: 8016-8022.
- Kindl, J., P. Jiroš, B. Kalinová, P. Žáček and I. Valterová, 2012. Females of the bumblebee parasite, *Aphomia sociella*, excite males using a courtship pheromone. *J. Chem. Ecol.*, **38**: 400-407.
- King, G. G. S., R. Gries, G. Gries and K. N. Slessor, 1995. Optical isomers of 3,13-dimethylheptadecane: sex pheromone components of the western false hemlock looper, *Nepytia freemani* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **21**: 2027-2045.
- Kinjo, M., M. Nagamine, H. Sugie and Y. Tamaki, 1996. Sex pheromone of the sugarcane shoot borer, *Tetramoera schistaceana* Snellen (Lepidoptera: Tortricidae: Olethreutinae). *Jpn. J. Appl. Entomol. Zool.*, **40**: 191-197.
- Klimetzek, D., G. Loskant, J. P. Vité and K. Mori, 1976. Disparlure: differences in pheromone perception between gypsy moth and nun moth. *Naturwissenschaften*, **63**: 581-582

- Klun, J. A. and T. A. Brindley, 1970. *cis*-11-Tetradecenyl acetate, a sex stimulant of the European corn borer. *J. Econ. Entomol.*, **63**: 779-780.
- Klun, J. A. and J. F. Robinson, 1972. Olfactory discrimination in the European corn borer and several pheromonally analogous moths. *Ann. Entomol. Soc. Am.*, **65**: 1337-1340.
- Klun, J. A., O. L. Chapman, K. C. Mattes, P. W. Wojtkowski, M. Beroza and P. E. Sonnet, 1973. Insect sex pheromones: minor amount of opposite geometrical isomer critical to attraction. *Science*, **181**: 661-663.
- Klun, J. A. and cooperators, 1975. Insect sex pheromones: intraspecific pheromonal variability of *Ostrinia nubilalis* in North America and Europe. *Environ. Entomol.*, **4**: 891-894.
- Klun, J. A. and G. A. Junk, 1977. Iowa European corn borer sex pheromone. Isolation and identification of four C₁₄ esters. *J. Chem. Ecol.*, **3**: 447-459.
- Klun, J. A. and S. Maini, 1979(a). Genetic basis of an insect chemical communication system: the European corn borer. *Environ. Entomol.*, **8**: 423-426.
- Klun, J. A., J. R. Plimmer, B. A. Bierl-Leonhardt, A. N. Sparks and O. L. Chapman, 1979(b). Trace chemicals: the essence of sexual communication systems in *Heliothis* species. *Science*, **204**: 1328-1330.
- Klun, J. A., B. A. Bierl-Leonhardt, M. Schwarz, J. A. Litsinger, A. T. Barrion, H. C. Chiang and Z. Jiang, 1980(a). Sex pheromone of the Asian corn borer moth. *Life Sci.*, **27**: 1603-1606.
- Klun, J. A., J. R. Plimmer, B. A. Bierl-Leonhardt, A. N. Sparks, M. Primiani, O. L. Chapman, G. H. Lee and G. Lepone, 1980(b). Sex pheromone chemistry of female corn earworm moth, *Heliothis zea*. *J. Chem. Ecol.*, **6**: 165-175.
- Klun, J. A., B. A. Bierl-Leonhardt, J. R. Plimmer, A. N. Sparks, M. Primiani, O. L. Chapman, G. Lepone and G. H. Lee, 1980(c). Sex pheromone chemistry of the female tobacco budworm moth, *Heliothis virescens*. *J. Chem. Ecol.*, **6**: 177-183.
- Klun, J. A., K. F. Haynes, B. A. Bierl-Leonhardt, M. C. Birch and J. R. Plimmer, 1981. Sex pheromone of the female artichoke plume moth, *Platyptilia carduidactyla*. *Environ. Entomol.*, **10**: 763-765.
- Klun, J. A., B. A. Leonhardt, J. D. Lopez, Jr. and L. E. Lachance,

1982. Female *Heliothis subflexa* (Lepidoptera: Noctuidae) sex pheromone: chemistry and congeneric comparisons. *Environ. Entomol.*, **11**: 1084-1090.
- Klun, J. A., C. C. Blickenstaff, M. Schwarz, B. A. Leonhardt and J. R. Plimmer, 1983. Western bean cutworm, *Loxagrotis albicosta* (Lepidoptera: Noctuidae): female sex pheromone identification. *Environ. Entomol.*, **12**: 714-717.
- Klun, J. A., B. A. Leonhardt, M. Schwarz, A. Day and A. K. Raina, 1986. Female sex pheromone of the pickleworm, *Diaphania nitidalis* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **12**: 239-249.
- Klun, J. A., J. W. Neal, Jr. and B. A. Leonhardt, 1989. Female sex pheromone of iris borer, *Macronoctua onusta* (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **15**: 1559-1566.
- Klun, J. A., M. Schwarz, B. A. Leonhardt and W. W. Cantelo, 1990. Sex pheromone of the female squashvine borer (Lepidoptera: Sesiidae). *J. Entomol. Sci.*, **25**(1): 64-72.
- Klun, J. A., Y. N. Baranchikov, V. C. Mastro, Y. Hijji, J. Nicholson, I. Ragenovich and T. A. Vshivkova, 2000. A sex attractant for the Siberian moth *Dendrolimus superans sibiricus* (Lepidoptera: Lasiocampidae). *J. Entomol. Sci.*, **35**(2): 158-166.
- Knauf, W., H. J. Bestmann, K. H. Koschatzky, J. Süss and O. Vostrowsky, 1979. Studies on the attractiveness of sex attractants towards *Tortrix viridana* L. and *Panolis flammea* Schiff. *Z. Angew. Entomol.*, **88**: 307-312.
- Knauf, W., H. J. Bestmann and O. Vostrowsky, 1984. Responses of male winter moths (*Operophtera brumata*) to their sex attractant (3Z,6Z,9Z)-1,3,6,9-nonadecatetraene and to some structural analogues. *Entomol. Exp. Appl.*, **35**: 208-210.
- Kochansky, J., R. T. Cardé, J. Liebherr and W. L. Roelofs, 1975(a). Sex pheromone of the European corn borer, *Ostrinia nubilalis* (Lepidoptera: Pyralidae), in New York. *J. Chem. Ecol.*, **1**: 225-231.
- Kochansky, J., J. Tette, E. F. Taschenberg, R. T. Cardé, K.-E. Kaissling and W. L. Roelofs, 1975(b). Sex pheromone of the moth, *Antheraea polyphemus*. *J. Insect Physiol.*, **21**: 1977-1983.
- Kochansky, J. P., W. L. Roelofs and P. Sivapalan, 1978. Sex pheromone of the tea tortrix moth (*Homona coffearia* Neitner). *J. Chem. Ecol.*, **4**: 623-631.
- Kochansky, J., A. Hill, J. W. Neal Jr., J.-A. Bentz and W. L. Roelofs,

1996. The pheromone of the eastern tent caterpillar, *Malacosoma americanum* (F.) (Lepidoptera: Lasiocampidae). *J. Chem. Ecol.*, **22**: 2251-2261.
- Komai, F. 1999. A taxonomic review of the genus *Grapholita* and allied genera (Lepidoptera: Tortricidae) in the Palaearctic region. *Ent. Scand. Suppl.* **55**: 1-226.
- Kohno, K., H. Iida, M. Takeda and K. Honda, 2014. Non-target lepidopteran insects specifically attracted to sex pheromone lures for *Helicoverpa armigera* and *H. assulta* (Lepidoptera: Noctuidae). *Jpn. J. Appl. Entomol. Zool.*, **58**: 343-350. (in Japanese)
- Kong, F.-L., S.-D. Yu, Y.-Q. Yu, X.-H. Hu, J.-W. Xu and P.-C. Zhu, 1990. Sex pheromone of eggplant borer, *Leucinodes orbonalis* Guenée. *Youji Huaxue*, **10**: 258-260.
- Kong, X., Z. Zhang, H. Wang and C. Zhao, 2005. Study on trace component in sex pheromone of *Dendrolimus* spp. *Sepu*, **23**(4): 370-373. [SciFinder, 2005:824663]
- Kong, X. B., Z. Zhang, C. H. Zhao and H. B. Wang, 2007(a). Female sex pheromone of the Yunnan pine caterpillar moth *Dendrolimus houi*: first (*E,Z*)-isomers in pheromone components of *Dendrolimus* spp. *J. Chem. Ecol.*, **33** (7): 1316-1327.
- Kong, X.-B., C.-H. Zhao and R. Wang, 2007(b). Sex pheromone of the larch caterpillar moth, *Dendrolimus superans*, from northeastern China. *Entomol. Exp. Appl.*, **124** (1): 37-44.
- Kong, X.-B., X.-L. Sun, H.-B. Wang, Z. Zhang, C-H. Zhao and K. C. J. H. Booij, 2011. Identification of components of the female sex pheromone of the Simaopine caterpillarmoth, *Dendrolimus kikuchii* Matsumura. *J. Chem. Ecol.*, **37**: 412-419.
- Kong, X.-B., K.-W. Liu, H.-B. Wang, S.-F. Zhang and Z. Zhang. 2012(a). Identification and behavioral evaluation of sex pheromone components of the Chinese pine caterpillar moth, *Dendrolimus tabulaeformis*. *PloS ONE.*, **7**: e33381.
- Kong, X.-B, H.-B. Wang and Z. Zhang, 2012(b). Determination of the absolute configuration of an EAG active component in the sex pheromone gland of *Semiothisa cirrearia* Bremer et Grey (Lepidoptera: Geometridae). *Acta Entomol. Sinica*, **55**: 162-167.
- Konjuchoy, V. P., B. G. Kovalev and N. R. Sattar-Zade, 1983. Isolation and identification of *Heliothis armigera* Hb. sex pheromone components. *Bioorganisk Khimiya*, **9**: 1422-1428.

- Konno, Y., K. Arai, K. Sekiguchi and Y. Matsumoto, 1982. (E)-10-Hexadecenal, a sex pheromone component of the yellow peach moth, *Dichocrocispunctiferalis* Guenée (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **17**: 207-217.
- Konyukhov, V. P. and B. G. Kovalev, 1988. Sex pheromone of lackey moth (*Malacosoma neustria* L.) females. *Bioorg. Khim*, **14**: 268-272. [*Chem. Abst.*, **108**: 164997e]
- Kosaki, A., T. Uehara, H. Naka, S. Matsuyama, T. Ando and H. Honda, 2021. Identification and behavioral assays of sex pheromone components in *Smerinthus tokyonis* (Lepidoptera: Sphingidae). *Appl. Entomol. Zool.*, **56**: 373-378.
- Koshihara, T., H. Yamada, Y. Tamaki and T. Ando, 1978. Field attractiveness of the synthetic sex-pheromone of the diamondback moth, *Plutella xylostella* (L.). *Appl. Entomol. Zool.*, **13**: 138-141.
- Kou, R., Y.-S. Chow, S. Takahashi and R. Yamaoka, 1989. Evidence for the presence and nature of a female sex pheromone of *Brithys crini* Fabricius (Lepidoptera: Noctuidae). *Chung Yang Yen Chiu Yuan Chih Wu Yen Chiu So Chuan K'an* (*Phytochem. Ecol.*), 429-434. [*Chem. Abst.*, **113**: 56115k]
- Kou, R., D. S. Tang, Y. S. Chow and H. K. Tseng, 1990. Sex pheromone components of female smaller tea tortrix moth, *Adoxophyes* sp. (Lepidoptera: Tortricidae) in Taiwan. *J. Chem. Ecol.*, **16**: 1409-1415.
- Kou, R., Y. S. Chow, S. Takahashi and R. Yamaoka, 1991. Evidence for presence and nature of female sex pheromone of *Brithys crini* Fabricius (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **17**: 2035-2042.
- Kou, R., H. Y. Ho, H. T. Yang, Y. S. Chow and H. J. Wu, 1992(a). Investigation of sex pheromone components of female Asian corn borer, *Ostrinia furnacalis* (Hübner) (Lepidoptera: Pyralidae) in Taiwan. *J. Chem. Ecol.*, **18**: 833-840.
- Kou, R., Y.-S. Chow and H.-Y. Ho, 1992(b). Chemical composition of sex pheromone gland extract in female oriental armyworm *Pseudaletia separata* Walker (Lepidoptera: Noctuidae) in Taiwan. *Bull. Inst. Zool., Academia Sinica*, **31**: 246-250.
- Kou, R., 1992c. Calling behavior and pheromone titer in the smaller tea tortrix moth, *Adoxophyes* sp. (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **18**: 855-861.

- Kovaleski, A., M. Botton, O. Nakano, E. F. Vilela and A. E. Eiras, 2003. Concentration and duration of the synthetic sex pheromone of *Bonagota cranaodes* (Meyrick) (Lepidoptera: Tortricidae) in an apple orchard. *Neotropical Entomol.*, **32**(1): 45-48.
- Kovalev, B. G., D. V. Sokolova, T. S. Bolgar and N. P. Sekerskaya, 1985. Identification of the sex pheromone of *Cacocimorpha pronubana*. *Khim. Prir. Soedin.*, 110-112. [*Chem. Abst.*, **102**: 182701j]
- Kovalev, B. G. and L. A. Nikolaeva, 1986. Attractiveness of some epoxy compounds for males of a tiger moth, *Phragmatobia fuliginosa* (Arctiidae), and of a geometrid moth, *Caustoloma (Therapis) flavicaria* (Geometriidae). *Zool Zh.*, **56**: 802-804.
- Kovalev, B. G. and S. Samoilov, 1987. Sex pheromone of female *Eurhodopa advenella* Zck. (Lepidoptera, Pyralidae): identification by GLC and EAG methods. *Khemoretseptsiya Nasekomykh*, **9**: 26-29.
- Kovalev, B. G., J. H. Tamlinson, T. S. Bolgar, T. P. Kolomoets, V. M. Kovalenko and R. R. Heath, 1988. Isolation and identification of a component in the sex pheromone of *Aegeria apiformis*. *Khim. Prir. Soedin.*, 126-128. [*Chem. Abst.*, **108**: 202002m]
- Kovalev, B. G., K. Lofsteot, S. F. Nedopekina, A. M. Sorochinskaya, K. S. Artokhin, E. A. Nesterov and E. Viik, 1991. Isolation and identification of the female sex pheromone of *Heliothis maritima* (Lepidoptera, Noctuidae). *Khim. Prir. Soedin.*, 574-577. [*Chem. Abst.*, **118**: 54297a]
- Kovalev, B. G., T. S. Bolgar, P. A. Zubov, D. G. Zharkov, M. Golosova, E. A. Nesterov and M. S. Tvaradze, 1993. Identification of additional components of the sex pheromone of *Dendrolimus pini*. *Chem. Nat. Compd.*, **29**: 135-136.
- Kovaleva, A. S., L. L. Ivanov, I. B. Piatnova, B. G. Kovalev, V. A. Minyailo and A. K. Minyailo, 1975. Results of testing synthetic compounds as to their attractiveness to insects in nature. *Khemoretseptsiya Nasekomykh*, **2**: 159-164. [*Chem. Abst.*, **85**: 105340]
- Kozaki, J., S. Nakagaki, R. Sato, M. Yukinari, H. Sugie, H. Noguchi and Y. Tamaki, 1984. Sex pheromone of the common apple leafroller, *Hoshinoa longicellana* Walsingham (Lepidoptera: Tortricidae): isolation and identification. *Appl. Entomol. Zool.*, **19**: 382-388.
- Kozlov, M. V., J. Zhu, P. Philipp, W. Francke, E. L. Zvereva, B. S. Hansson and C. Löfstedt, 1996. Pheromone specificity in *Eriocrania semipurpurella* (Stephens) and *E. sangii* (Wood) (Lepidoptera: Eriocraniidae) based on chirality of semiochemicals.

- J. Chem. Ecol.*, **22**: 431-454.
- Krasnoff, S. B., K. W. Vick and J. A. Coffelt, 1984. (Z,E)-9,12-Tetradecadien-1-ol: a component of the sex pheromone of *Ephestia elutella* (Hübner) (Lepidoptera: Pyralidae). *Environ. Entomol.*, **13**: 765-767.
- Krasnoff, S. B. and W. L. Roelofs, 1988. Sex pheromone released as an aerosol by the moth *Pyrrharctia isabella*. *Nature*, **333**: 263-265.
- Krishnakumari, B., A. L. Prasuna, K. N. Jyothi, M. Y. Valli, S. Sighamony, A. R. Prasad and J. S. Yadav 1998. Behavioural and electrophysiological responses of *Achaea janata* Linn. (Lepidoptera: Noctuidae) males to synthetic female-produced sex pheromone components. *J. Entomol. Res.*, **22**(3): 197-202.
- Krivokhizhin, V. I., V. V. Brovko, V. A. Minyailo and G. S. Zolotarenko, 1984. Cis-11-hexadecen-1-yl acetate as a sexual attractant for cutworm moths (Lepidoptera, Noctuidae) in Western Siberia. *Sib. Vestn. S-kh. Nauki*, 33-38. [*Chem. Abst.*, **101**: 124847e]
- Krokos, F. D., A. Ameline, J. Bau, A. Sans, M. Konstantopoulou, B. Frerot, A. Guerrero, M. Eizaguirre, C. Malosse, O. Etchepare, R. Albajes and B. E. Mazomenos, 2002. Comparative studies of female sex pheromone components and male response of the corn stalkborer *Sesamia nonagrioides* in three different populations. *J. Chem. Ecol.*, **28**(7): 1463-1472.
- Kuenen, L. P. S., J. W. Peacock, P. J. Silk and S. S. Wright, 1990. Identification of the primary female sex pheromone component of the mimosa webworm (Lepidoptera: Plutellidae). *Environ. Entomol.*, **19**: 1-7.
- Kuenen, L. P. S., J. S. McElfresh and J. G. Millar, 2010. Identification of critical secondary components of the sex pheromone of the navel orangeworm (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **103**: 314-330.
- Kurihara, M., K. Usui, K. Uchiumi and S. Tatsuki, 1991. Sex pheromone of the lawn grass cutworm moth, *Spodoptera depravata* (Butler) (Lepidoptera: Noctuidae). *Jap. J. Appl. Entomol. Zool.*, **35**: 323-324.
- Kuwahara, Y., C. Kitamura, S. Takahashi, H. Hara, S. Ishii and H. Fukami, 1971(a). Sex pheromone of the almond moth and the Indian meal moth: cis-9,trans-12-tetradecadienyl acetate. *Science*, **171**: 801-802.
- Kuwahara, Y., H. Hara, S. Ishii and H. Fukami, 1971(b). The sex pheromone

- of the Mediterranean flour moth. *Agric. Biol. Chem.*, **35**: 447-448.
- Kuwahara, Y. and J. E. Casida, 1973. Quantitative analysis of the sex pheromone of several phycitid moths by electron-capture gas chromatography. *Agric. Biol. Chem.*, **37**: 681-684.
- Kuwahara, Y. and L. T. M. Yen, 1979. Identification of bombykol in *Bombyx mandarina* females (Lepidoptera: Bombycidae). *Appl. Entomol. Zool.*, **14**: 114-117.
- Kuwahara, Y., N. Mori, S. Yamada and T. Nemoto, 1984. Evaluation of bombykol as the sex pheromone of *Bombyx mandarina* (Lepidoptera: Bombycidae). *Appl. Entomol. Zool.*, **19**: 265-267.
- Lacey, M.J. and C.J. Sanders, 1992. Chemical composition of sex pheromone of oriental fruit moth and rates of release by individual female moths. *J. Chem. Ecol.*, **18**: 1421-1435.
- La Croix, E. A. S., H. Z. Thindwa and D. R. Hall, 1985. Field studies with the synthetic female sex pheromones of *Cryptophlebia batrachopa* and *C. leucotreta*, pests of macadamia in Malawi. *Trop. Pest Manag.*, **31**: 189-191.
- Lalanne-Cassou, B. and B. Frerot, 1980. Analysis of pheromonal secretion of South African carnation tortrix, *Epichoristodes acerbella* (Walk.) (Lepidoptera: Tortricidae). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **291**: 845-848.
- Landolt, P. J. and C. E. Curtis, 1982. Interspecific sexual attraction between *Pyralis farinalis* L. and *Amyelois transitella* (Walker) (Lepidoptera: Pyralidae). *J. Kansas Entomol. Soc.*, **55**: 248-252.
- Landolt, P. J., J. D. Solomon and R. E. Doolittle, 1985. Attraction of male *Prionoxystus piger* (Lepidoptera: Cossidae) to isomers of 3,5-tetradecadien-1-ol acetate. *Florida Entomol.*, **68**: 228-230.
- Landolt, P. J., R. R. Heath, P. E. Sonnet and K. Matsumoto, 1986(a). Attraction of *Harrisina americana* and *Acoloithus falsarius* males (Lepidoptera: Zygaenidae) to (R)-(-)-2-butyl-(Z)-7-tetradecenoate. *Environ. Entomol.*, **15**: 959-962.
- Landolt, P. J., R. R. Heath and N. C. Leppla, 1986(b). (Z,Z,Z)-3,6,9-Eicosatriene and (Z,Z,Z)-3,6,9-heneicosatriene as sex pheromone components of a grass looper, *Mocis disseverans* (Lepidoptera: Noctuidae). *Environ. Entomol.*, **15**: 1272-1274.
- Landolt, P. J. and R. R. Heath, 1986(c). A sex attractant synergist for *Trichoplusia oxygramma* (Lepidoptera: Noctuidae). *Florida*

- Entomol.*, **69**: 425-426.
- Landolt, P. J., J. H. Tumlinson and M. M. Brennan, 1989(a). Attraction of *Amphion floridensis* (Lepidoptera: Sphingidae) to bombykal, (E,Z)-10,12-hexadecadienal. *Florida Entomol.*, **72**: 324-327.
- Landolt, P. J. and R. R. Heath, 1989(b). Lure composition, component ratio, and dose for trapping male *Mocis latipes* (Lepidoptera: Noctuidae) with synthetic sex pheromone. *J. Econ. Entomol.*, **82**: 307-309.
- Landolt, P. J. and R. R. Heath, 1991. Zygaenidae trapped with enantiomers of 2-butyl (Z)-7-tetradecenoate. *J. Lepido. Soc.*, **45**: 63-65.
- Landolt, P. J., M. Tóth, W. Francke and K. Mori, 1996. Attraction of moths (Lepidoptera: Arctiidae, Geometridae, Noctuidae) to enantiomers of several epoxydienes. *Florida Entomol.*, **79**(3): 392-397.
- Landolt, P. J. and C. L. Smithhisler 1998. Isolation and identification of female sex pheromone and development of a sex attractant for *Lacanobia subjuncta*. *J. Chem. Ecol.*, **24**(8): 1369-1379.
- Landolt, P. J., C. Smithhisler, T. Adams and R. S. Zack, 2003. An improved multi-component sex attractant for trapping male western yellowstriped armyworm, *Spodoptera praefica* (Grote) (Lepidoptera: Noctuidae). *Agric. Forest Entomol.*, **5**: 333-339.
- Landolt, P. J., C. Guédot and R. S. Zack, 2011. Spotted cutworm, *Xestia c-nigrum* (L.) (Lepidoptera: Noctuidae) responses to sex pheromone and blacklight. *J. Appl. Entomol.*, **135**: 593-600.
- Landolt, P. J. and D. Roberts, 2012. A sex attractant for trapping *Crambus cypridalis* (Lepidoptera: Crambidae). *Florida Entomol.*, **95**: 194-195.
- Lapointe, S. L., D. G. Hall, Y. Murata, A. L. Parra-pedrazzoli, J. M. S. Bento, E. F. Vilela and W. S. Leal, 2006. Field evaluation of a synthetic female sex pheromone for the leafmining moth *Phyllocnistis citrella* (Lepidoptera: Gracillariidae) in Florida citrus. *Florida Entomol.*, **89** (2), 274-276.
- Lapointe, S. L., W. Barros-Parada, E. Fuentes-Contreras, H. Herrera, T. Kinsho, Y. Miyake, R. P. Niedz and J. Bergmann 2017. Use of mixture designs to investigate contribution of minor sex pheromone components to trap catch of the carpenterworm moth, *Chilecomadia valdiviana*. *J. Chem. Ecol.*, **43**: 1046-1655.

- Lassance, J.-M., G. P. Svensson, M. V. Kozlov, W. Francke and C. Löfstedt, 2019. Pheromones and barcoding delimit boundaries between cryptic species in the primitive moth genus *Eriocrania* (Lepidoptera: Eriocraniidae). *J. Chem. Ecol.*, **45**: 429-439.
- Leal, W. S., J. M. S. Bento, Y. Murata, M. Ono, J. R. P. Parra and E. F. Vilela, 2001. Identification, synthesis, and field evaluation of the sex pheromone of the citrus fruit borer *Ecdytolopha aurantiana*. *J. Chem. Ecol.*, **27**(10): 2041-2051.
- Leal, W. S., A. L. Parra-Pedrazzoli, K.-E. Kaissling, T. I. Morgan, F. G. Zalom, D. J. Pesak, E. A. Dundulis, C. S. Burks and B. S. Higbee, 2005. Unusual pheromone chemistry in the navel orangeworm: novel sex attractants and a behavioral antagonist. *Naturwissenschaften*, **92**: 139-146.
- Leal, W. S., A. L. Parra-Pedrazzoli, A. A. Cossé, Y. Murata, J. M. S. Bento and E. F. Vilela, 2006. Identification, synthesis, and field evaluation of the sex pheromone from the citrus leafminer, *Phyllocnistis citrella*. *J. Chem. Ecol.*, **32**(1): 155-168.
- Lebedeva, K. V., V. V. Voronkova and N. I. Bocharova, 1991. Pheromone of the turnip moth (*Agrotis segetum*). *Agrokhimiya*, 99-114. [*Chem. Abst.*, **116**: 209681f]
- Lebedeva, K. V., N. V. Vendilo, S. A. Kurbatov, V. A. Pletnev, V. L. Ponomarev, Y. B. Pyatnova and N. I. Bocharova, 2000(a). Identification of the pheromone of eastern-meadow cutworm *Mythimna separata* (Lepidoptera: Noctuidae). *Agrokhimiya*, (5): 57-69. [*Chem. Abst.*, **134**: 278087]
- Lebedeva, K. V., N. V. Vendilo, V. A. Pletnev, V. L. Ponomarev, S. A. Kurbatov, V. V. Voronkova and V. A. Shchennikov, 2000(b). Study of the pheromone of the heart-and-dart moth, *Agrotis exclamationis* (Lepidoptera: Noctuidae). *Agrokhimiya*, (8): 71-75. [*Chem. Abst.*, **134**: 262292]
- Lecheva, I., 1999. Possibilities of implementation of synthetic sex pheromones for the geometrid *Erannis bajaria* (Lepidoptera: Geometridae). *Restenievud. Nauki.*, **36**(7-8): 397-400. [*Chem. Abs.*, **132**: 318901]
- Lee, S., D.-W. Lee and K. S. Boo, 2005. Sex pheromone composition of the diamondback moth, *Plutella xylostella* (L.) in Korea. *J. Asia-Pacific Entomol.*, **8**: 243-248.
- Lee, S.-C., E.-C. Koo, D.-H. Lee, S.-M. Seo, M.-J. Huh, I. Nam & J.-H. Park and I.-K. Park, 2020. Identification of the female-produced sex pheromone of the dotted white geometrid *Naxa seriaria* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **46**: 927-934.

- Lee, S.-C., J.-W. Lee, D. H. Lee, M.-J. Huh, I. Nam, J.-H. Park, M. Jung and I.-K. Park, 2022. Identification of sex pheromone components of Korean *Dioryctria abietella* (Lepidoptera: Pyralidae) population and synergism of pheromone and pine cone volatile blends. *J. Econ. Entomol.*, **115**: 178-186.
- Leeuw, I., 1983. *Thyris maculata* (Thyrididae) and three species of clearwing moths (Sesiidae) attracted to an artificial chemical bait. *J. Lepi. Soc.*, **37**: 87-88.
- Legrand, S., M. Botton, M. Coracini, P. Witzgall and C. R. Unelius, 2004. Synthesis and field tests of sex pheromone components of the leafroller *Argyrotaenia sphaleropa*. *Z. Naturforsch.*, **59c**: 708-712.
- Lemmen, J. and M. Evenden, 2009. Peripheral and behavioral plasticity of pheromone response and its hormonal control in a long-lived moth. *J. Exp. Biol.*, **212**: 2000-2006.
- Leonhardt, B. A., J. W. Neal, Jr., J. A. Klun, M. Schwarz and J. R. Plimmer, 1983. An unusual lepidopteran sex pheromone system in the bagworm moth. *Science*, **219**: 314-316.
- Leonhardt, B. A., V. C. Mastro, M. Schwarz, J. D. Tang, R. E. Charlton, A. Pellegrini-Toole, J. D. Warthen, Jr., C. P. Schwalbe and R. T. Cardé, 1991. Identification of sex pheromone of browntail moth, *Euproctis chrysorrhoea* (L.) (Lepidoptera: Lymantriidae). *J. Chem. Ecol.*, **17**: 897-910.
- Leskey, T. C., J. C. Bergh, J. F. Walgenbach and A. J. Zhang, 2006. Attractiveness and specificity of pheromone-baited traps for male dogwood borer, *Synanthedon scitula* Harris (Lepidoptera: Sesiidae). *Environ. Entomol.*, **35**: 268-275.
- Levi-Zada, A., D. Fefer, L. Anshelevitch, A. Litovsky, M. Bengtsson, G. Gindin and V. Soroker, 2011(a). Identification of the sex pheromone of the lesser date moth, *Batrachedra amydraula*, using sequential SPME auto-sampling. *Tetrahedron Lett.*, **52**: 4550-4553.
- Levi-Zada, A., S. Ben-Yehuda, E. Dunkelblum, G. Gindin, D. Fefer, A. Protasov, T. Kuznetsowa, S. Manulis-Sasson and Z. Mendel, 2011(b). Identification and field bioassays of the sex pheromone of the yellow-legged clearwing *Synanthedon vespiformis* (Lepidoptera: Sesiidae). *Chemoecology*, **21**: 227-233.
- Levi-Zada, A., A. Sadowsky, S. Dobrinin, M. David, T. Ticuchinski, D. Fefer, A. Greenberg and D. Blumberg, 2013. Reevaluation of the sex pheromone of the lesser date moth, *Batrachedra amydraula*,

- using autosampling SPME-GC/MS and field bioassays. *Chemoecology*, **23**: 13-20.
- Lewis, T., C. Wall, E. D. M. Macaulay and A. R. Greenway, 1975. The behavioural basis of a pheromone monitoring system for pea moth, *Cydia nigricana*. *Ann. Appl. Biol.*, **80**: 257-274.
- Li, J., R. Gries, G. Gries, K. N. Slessor, G. G. S. King, W. W. Bowers and R. J. West, 1993(a). Chirality of 5,11-dimethylheptadecane, the major sex pheromone component of the hemlock looper, *Lambdina fiscellaria* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **19**: 1057-1062.
- Li, J.-X., G. Gries, R. Gries, J. Bikic and K. N. Slessor, 1993(b). Chirality of synergistic sex pheromone components of the western hemlock looper *Lambdina fiscellaria lugubrosa* (Hulst) (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **19**: 2547-2561.
- Li, Z.-M., E.-Y. Yao, T.-L. Liu, Z.-P. Liu, S.-H. Wang, H.-Q. Zhu, G. Zhao and Z.-L. Ren, 1993(c). Structural elucidation of sex pheromone components of the Geometridae *Semiothisa cinerearia* (Bremer et Grey) in China. *Chinese J. Chem.*, **11**: 251-256.
- Li, W.-W., X.-Z. Meng, Y.-H. Deng and X.-G. Hu, 1993(d). Experimental effects on field trapping of noctuids using several sex attractants. *Entomol. Knowledge*, **30**: 38-40.
- Liebhold, A. M. and W. J. A. Volney, 1985. Effects of attractant composition and release rate on attraction of male *Choristoneura retiniana*, *C. occidentalis*, and *C. carnana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **117**: 447-457.
- Liebhold, A. M. and P. J. Silk, 1991. Capture of *Choristoneura pinus maritima* in traps baited with *C. pinus pinus* pheromone components (Tortricidae). *J. Lepidop. Soc.*, **45**: 172-173.
- Lin, Y. M., Y. S. Chow and H. C. Tzeng, 1982. Field trapping of the diamondback moth *Plutella xylostella* (Linnaeus) and *Pseudaletia separata* Walker using the synthetic sex pheromone of the diamondback moth pest of cruciferous vegetables, Taiwan. *Bull. Inst. Zool. Acad. Sin.*, **21**: 121-127.
- Lin, G.-Q., G.-Z. Guo, Y.-W. Wu, B.-Q. Wu, L.-G. Liu and W.-S. Zhou, 1984. Study on identification and syntheses of insect pheromone XV. Multiple sex pheromone of *Ancylis sativa*. *Kexue Tongbao*, **29**: 1670-1674.
- Lin, G.-Q., 1993. Structural elucidation and chiral syntheses of insect pheromones and extension of the synthetic approaches. *Pure Appl. Chem.*, **65**: 1233-1240.

- Linn, C. E. and W. L. Roelofs, 1983. Effect of varying proportions of the alcohol component on sex pheromone blend discrimination in male Oriental fruit moths. *Physiol. Entomol.*, **8**: 291-306.
- Linn, C. E. Jr., L. B. Bjostad, J. W. Du and W. L. Roelofs, 1984. Redundancy in a chemical signal: Behavioral responses of male *Trichoplusia ni* to a 6-component sex pheromone blend. *J. Chem. Ecol.* **10**: 1635-1658.
- Linn, C. E., Jr., J. Du, A. Hammond and W. L. Roelofs, 1987. Identification of unique pheromone components for soybean looper moth *Pseudoplusia includens*. *J. Chem. Ecol.*, **13**: 1351-1360.
- Linnaaluoto, E. T. and S. Koponen, 1983. Northern moths (Lepidoptera) attracted by synthetic pheromones. *Ann. Entomol. Fenn.*, **49**: 64.
- Liska, J., R. Hochmut and J. Jurik, 1990. Field trials with sexual attractants of *Zeiraphera griseana* (Lepidoptera, Tortricidae) in the High Tatras. *Acta Entomol. Bohemoslov.*, **87**: 81-86.
- Liu, Y.-X. and X.-Z. Meng, 2002(a). Identification of sex pheromone components for *Acleris fimbriana* (Lepidoptera: Tortricidae). *Can. Entomol.* **134**: 511-518.
- Liu, Y.-X. and X.-Z. Meng, 2002(b). Behavioral response of male *Acleris fimbriana* Thunberg et Becklin (Lepidoptera: Tortricidae) to synthetic sex pheromones. *Acta Entomol. Sinica (Kunchong Xuebao)*, **45**: 436-440.
- Liu, Y.-X. and X.-Z. Meng, 2003. Trapping effect of synthetic sex pheromone of *Acleris fimbriana* (Lepidoptera: Tortricidae) in Chinese northern orchard. *Z. Naturforsch.*, **58c**: 421-425.
- Liu, F., L. Guo, Q.-C. Li, S.-F. Zhang, X.-B. Kong, Z. Zhang and F.-Q. Huang, 2019. Identification of the active components of sex pheromone of female moths of *Micromelalopha sieversi* (Lepidoptera: Notodontidae). *Acta Entomol. Sinica*, **62**: 814-822.
- Liu, F., L. Guo, S. Zhang, X. Kong and Z. Zhang, 2021(a). Synthesis and bioactivity of (13Z,15E)-octadecadienal: a sex pheromone component from *Micromelalopha sieversi* Staudinger (Lepidoptera: Notodontidae). *Pest Manag. Sci.*, **77**: 264-272.
- Liu, H., S. Xu, R. Dou, Y. Wang, X. Jing, J. Liu, J. Wang, Z. Wang and J. Zhang, 2021(b). Field trials of synthetic sex attractants of *Illiberis pruni* in China. *Entomol. Exp. Appl.*, **169**: 640-645.
- Liu, H., R. Dou, J. Wang, Y. Zhu, J. Li, J. Liu, Y. Wang, J. Zhang and X. Jing, 2022. Evaluation of sex attractants for monitoring

- of *Illiberis ulmivora* Graeser (Lepidoptera: Zygaenidae: Procridinae): height, distance, placement, climatic conditions. *Entomol. Res.*, **52**: 99-107.
- Löfstedt, C., J. N. C. VanDerPers, J. Löfqvist, B. S. Lanne, M. Appelgren, G. Bergström and B. Thelin, 1982. Sex pheromone components of the turnip moth, *Agrotis segetum*: chemical identification, electrophysiological evaluation and behavioral activity. *J. Chem. Ecol.*, **8**: 1305-1321.
- Löfstedt, C., J. N. C. Van der Pers, J. Löfqvist and B. S. Lanne, 1983. Sex pheromone of the cone pyralid *Dioryctria abietella*. *Entomol. Exp. Appl.*, **34**: 20-26.
- Löfstedt, C. and J. N. C. van der Pers, 1985(a). Sex pheromones and reproductive isolation in four European small ermine moths. *J. Chem. Ecol.*, **11**: 649-666.
- Löfstedt, C., B. S. Lanne, J. Löfqvist, M. Appelgren and G. Bergström, 1985(b). Individual variation in the pheromone of the turnip moth, *Agrotis segetum*. *J. Chem. Ecol.*, **11**: 1181-1196.
- Löfstedt, C., C. E. Linn, Jr. and J. Löfqvist, 1985(c). Behavioral responses of male turnip moths, *Agrotis segetum*, to sex pheromone in a flight tunnel and in the field. *J. Chem. Ecol.*, **11**: 1209-1221.
- Löfstedt, C., W. M. Herrebout and J.-W. Du, 1986(a). Evolution of the ermine moth pheromone tetradecyl acetate. *Nature*, **323**: 621-623.
- Löfstedt, C., J. Löfqvist, B. S. Lanne, J. N. C. Van Der Pers and B. S. Hansson, 1986(b). Pheromone dialects in European turnip moths *Agrotis segetum*. *Oikos*, **46**: 250-257.
- Löfstedt, C. and W. M. Herrebout, 1988. Sex pheromones of three small ermine moths found on the European spindle tree. *Entomol. Exp. Appl.*, **46**: 29-38.
- Löfstedt, C., B. S. Hansson, H. J. Dijkerman and W. M. Herrebout, 1990. Behavioural and electrophysiological activity of unsaturated analogues of the pheromone tetradecyl acetate in the small ermine moth *Yponomeuta rorellus*. *Physiol. Entomol.*, **15**: 47-54.
- Löfstedt, C., B. S. Hansson, M. Tóth, G. Szöcs, V. Buda, M. Bengtsson, N. Ryrholm, M. Svensson and E. Priesner, 1994. Pheromone differences between sibling taxa *Diachrysia chrysitis* (Linnaeus, 1758) and *D. tutti* (Kostrowicki, 1961) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **20**: 91-109.
- Löfstedt, C., J. Zhu, M. V. Kozlov, V. Buda, E. V. Jirle, S. Hellqvist,

- J. Löfqvist, E. Plass, S. Franke and W. Francke, 2004. Identification of the sex pheromone of the currant shoot borer *Lampronia capitella*. *J. Chem. Ecol.*, **30**(3): 643-658.
- Löfstedt, C., G. P. Svensson, E. V. Jirle, O. Rosenberg, A. Roques and J. G. Millar, 2012. (3Z,6Z,9Z,12Z,15Z)-pentacosapentaene and (9Z,11E)-tetradecadienyl acetate: sex pheromone of the spruce coneworm *Dioryctria abietella* (Lepidoptera: Pyralidae). *J. Appl. Entomol.*, **136**: 70-78.
- Loneragan, G. C., B. M. Ponder, W. D. Seabrook and L. R. Kipp, 1989. Sex pheromone components of the blueberry leaf tier moth *Croesia curvalana* Kearfott (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **15**: 2495-2506.
- Lopez, J. D., Jr. and T. N. Shaver, 1990. Plastic laminate dispenser for components of the *Heliothis zea* sex pheromone. *Southwest. Entomol.*, **15**(1): 1-8.
- Lu, P.-F., H.-L. Qiao and Y.-Q. Luo, 2013. Female sex pheromone blends and male response of the legume pod borer, *Maruca vitrata* (Lepidoptera: Crambidae), in two populations of mainland China. *Z. Naturforsch.*, **68 c**: 416-427.
- Lu, J., X. Yao, Y. Shen, C. Du, Q. Guo and Y. Du, 2024. Physiological status of rice leaf-roller *Cnaphalocrocis medinalis* (Lepidoptera: Crambidae) adults trapped by sex pheromone and floral odor. *Insects*, **15**: 637.
- Luo, Z.-X., Z.-Q. Li, X.-M. Cai, L. Bian and Z. M. Chen, 2017. Evidence of premating isolation between two sibling moths: *Ectropis grisescens* and *Ectropis obliqua* (Lepidoptera: Geometridae). *J. Econ. Entomol.*, **110**: 2364-2370.
- Luo, Z., X. Cai, Z. Li, L. Bian, Z. Xin, F. Hussain and Z. Chen, 2019. Identification of sex pheromone of *Miltochrista striata* (Lepidoptera: Arctiidae). *J. Econ. Entomol.*, **112**: 665-670.
- Lynch, R. E., J. A. Klun, B. A. Leonhardt, M. Schwarz and J. W. Garner, 1984. Female sex pheromone of the lesser cornstalk borer, *Elasmopalpus lignosellus* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **13**: 121-126.
- Ma, T., Y. Li, Z. Sun and X. Wen, 2014. (Z,E)-9,12-Tetradecadien-1-ol: A major sex pheromone component of *Euzophera pyriella* (Lepidoptera: Pyralidae) in Xinjiang, China. *Florida Entomol.*, **97**: 496-503.
- Ma, T., Z. Liu, J. Lu, Z. Sun, Y. Li, X. Wen and Y. Cui, 2015. A key compound: (Z)-9-tetradecen-1-ol as sex pheromone active

- component of *Hypsipyla robusta* (Lepidoptera: Pyralidae). *Chemoecology*, **25**: 325-330.
- Ma, T., Q. Xiao, Y.-G. Yu, C. Wang, C.-Q. Zhu, Z.-H. Sun, X.-Y. Chen and X.-J. Wen, **2016**. Analysis of tea geometrid (*Ectropis grisescens*) pheromone gland extracts using GC-EAD and GCxGC/TOFMS. *J. Agric. Food Chem.*, **64**: 3161-3166.
- Ma, T., Z. Liu, C. Wang, S. Zhang, X. Shi, Z. Sun, X. Chen, C. Jia, C. Wang, Y. He and X. Wen, **2017**. Production, identification, and field evaluation of sex pheromone from calling females in *Diaphania angustalis* (Lepidoptera: Crambidae). *Environ. Sci. Pollut. Res. Int.*, **24**: 24485-24493.
- Mafi, S. A., L. V. Vang, Y. Nakata, N. Ohbayashi, M. Yamamoto and T. Ando, **2005**. Identification of the sex pheromone of the citrus leafminer (*Phyllocnistis citrella* Stainton, Lepidoptera: Gracillariidae) with a trial of control by the communication disruption method. *J. Pestic. Sci.*, **30**(4): 361-367.
- Magsi, F. H., Z. Li, X. Cai, M. Yamamoto, L. Bian, Y. Zhao, L. Zhou, C. Xiu, N. Fu, T. Ando, Z. Luo and Z. Chen, **2022**. Identification of a unique three-component sex pheromone produced by the tea black tussock moth, *Dasychira baibarana* (Lepidoptera: Erebiidae: Lymantriinae). *Pest Manag. Sci.*, **78**: 2607-2617.
- Maier, C. T., R. Gries and G. Gries **1998**. Sex pheromone components of pitch pine looper, *Lambdina pellucidaria*. *J. Chem. Ecol.*, **24**(3): 491-500.
- Maini, S., **1973**. Preliminary tests with synthetic sex attractants of the European corn borer. *Inf. Tore Fitopatol.*, **9**: 11-14.
- Malo, E. A., J. C. Rojas, G. Lopez-Guillen and J. F. Barrera, **2009**. Chemical analysis of female volatiles and field response of the coffee leafminer moth (Lepidoptera: Lyonetiidae) to stereoisomers of its major sex pheromone component. *Florida Entomol.*, **92**: 548-553.
- Malosse, C., C. Descoins and B. Frerot, **1993**. Identification of minor component of the sex pheromone of the leopard moth (*Zeuzera pyrina* L.; Lepidoptera, Cossidae) by capillary gas chromatography. *J. High Resol. Chromat.*, **16**: 123-125.
- Marks, R.J., **1976**. Field evaluation of gossyp lure, the synthetic sex pheromone of *Pectinophora gossypiella* (Saund.) (Lepidoptera, Gelechiidae) in Malawi. *Bull. Entomol. Res.*, **66**: 267-278.
- Matsuoka, K., M. Yamamoto, R. Yamakawa, M. Muramatsu, H. Naka, Y. Kondo and T. Ando, **2008**. Identification of novel C₂₀ and C₂₂ trienoic

acids from arctiid and geometrid female moths that produce polyenyl Type II sex pheromone components. *J. Chem. Ecol.*, **34**: 1437-1445.

Masnyk, M., J. Fried and W. L. Roelofs, 1989. Synthesis and pheromone properties of (Z)-7,7-difluoro-8-docecenyl acetate, a difluoro derivative of the sex pheromone of the Oriental fruit moth. *Tetrahedron Lett.*, **30**: 3243-3246.

Mayer, M. S., J. R. McLaughlin and E. R. Mitchell, 1995. Use of paired and single traps to assess perception and discrimination of sex pheromone mixtures in the field by *Trichoplusia ni* (Hubner). *J. Chem. Ecol.*, **21**: 2057-2068.

Mazomenos, B. E., D. Bardas and A. Pantazimazomenou, 1989(a). Sesamianonagrioides sex pheromone: field attractiveness use of pheromone traps for monitoring and control. *Acta Phytopathol. Entomol. Hung.*, **24**: 153-156.

Mazomenos, B. E., 1989(b). Sex pheromone components of corn stalk borer *Sesamia nonagrioides* (Lef.). Isolation, identification, and field tests. *J. Chem. Ecol.*, **15**: 1241-1247.

Mazomenos, B. E., D. Raptopoulos, I. Lefkidou and A. Mazomenos-Pantazi, 1994. Female sex pheromone components of jasmine moth *Palpita unionalis* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **20**: 745-751.

Mazor, M., M. Harel, S. Levsky and E. Dunkelblum, 1991. Sex pheromone components of female *Cornutiplusia circumflexa*. *Entomol. Exp. Appl.*, **60**: 167-172.

Mazor, M. and E. Dunkelblum 1992. Role of sex pheromone components in behavioral reproductive isolation between *Autographa gamma* (L.) and either *Trichoplusia ni* (Hubner) or *Chrysodeixis chalcites* (Esp.) (Lepidoptera: Noctuidae: Plusiinae). *J. Chem. Ecol.*, **18**(12): 2373-2384.

McBrien, H., G. Gries, R. Gries, J. H. Borden, G. J. R. Judd, G. G. S. King and K. N. Slessor, 1992. Sex pheromone components of the eyespotted bud moth, *Spilonota ocellana* (Denis and Schifferrmüller) (Lepidoptera: Olethreutidae). *Can. Entomol.*, **124**: 1391-1394.

McBrien, H. L., G. J. R. Judd, A. K. Minks, J. W. Johnson, R. F. Smith, H. M. A. Thistlewood and J. T. Troubridge, 1994. Response of male eye-spotted bud moth, *Spilonota ocellana* (Lepidoptera: Tortricidae), to different pheromone blends in North America and the Netherlands. *J. Chem. Ecol.*, **20**: 625-630.

McDonough, L. M., D. A. George, B. A. Butt, J. M. Ruth and K. R. Hill, 1972. Sex pheromone of the codling moth: structure and synthesis.

- Science*, **177**:177-178.
- McDonough, L. M. and H. R. Moffitt, 1974. Sex pheromone of the codling moth. *Science*, **183**: 978.
- McDonough, L. M. and J. A. Kamm, 1979. Sex pheromone of the cranberry girdler, *Chrysoteuchia topiaria* (Zeller) (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **5**: 211-219.
- McDonough, L. M., J. A. Kamm and B. A. Bierl-Leonhardt, 1980. Sex pheromone of the armyworm, *Pseudaletia unipuncta* (Haworth) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **6**: 565-572.
- McDonough, L. M., M. P. Hoffmann, B. A. Bierl-Leonhardt, C. L. Smithhisler, J. B. Bailey and H. G. Davis, 1982(a). Sex pheromone of the avocado pest, *Amorbia cuneana* (Walsingham) (Lepidoptera: Tortricidae): structure and synthesis. *J. Chem. Ecol.*, **8**: 255-265.
- McDonough, L. M., D. S. Moreno, B. A. Bierl-Leonhardt and B. A. Butt, 1982(b). Isolation and identification of a component of the female sex pheromone gland attractive to male *Xylomyges curialis*. *Environ. Entomol.*, **11**: 660-662.
- McDonough, L. M., J. A. Kamm, D. A. George, C. L. Smithhisler and S. Voerman, 1982(c). Sex attractant for the western lawn moth, *Tehama bonifatella* Hulst. *Environ. Entomol.* **11**: 711-714.
- McDonough, L. M., J. B. Bailey, M. P. Hoffmann, B. A. Leonhardt, D. F. Brown, C. L. Smithhisler and K. Olsen, 1986. *Sabulodes caberata* Guenée (Lepidoptera: Geometridae). Components of its sex pheromone gland. *J. Chem. Ecol.*, **12**: 2107-2116.
- McDonough, L. M., H. G. Davis and S. Voerman, 1987. Blackheaded fireworm: laboratory and field studies of its sex pheromone. *J. Chem. Ecol.*, **13**: 1235-1242.
- McDonough, L. M., R. D. Eikenbary, M. T. Smith, H. G. Davis, C. L. Smithhisler, G. A. Hedger, R. K. Campbell, J. A. Payne, W. Reid and J. R. McVay, 1990(a). Sex pheromone of hickory shuckworm *Cydia caryana*. Development of an effective field lure. *J. Chem. Ecol.*, **16**: 317-324.
- McDonough, L. M., H. G. Davis, C. L. Smithhisler, S. Voerman and P. S. Chapman, 1990(b). Apple ermine moth, *Yponomeuta malinellus* Zeller. Two components of female sex pheromone gland highly effective in field trapping tests. *J. Chem. Ecol.*, **16**: 477-486.
- McDonough, L. M., H. G. Davis, P. S. Chapman and C. L. Smithhisler, 1993. Response of male codling moths (*Cydia pomonella*) to

- components of conspecific female sex pheromone glands in flight tunnel tests. *J. Chem. Ecol.*, **19**: 1737-1748.
- McDonough, L. M., A. L. Averill, H. G. Davis, C. L. Smithhisler, D. A. Murray, P. S. Chapman, S. Voerman, L. J. Dapsis and M. M. Averill, **1994**. Sex pheromone of cranberry fruitworm, *Acrobasis vaccinii* Riley (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **20**: 3269-3279.
- McDonough, L. M., H. G. Davis, P. S. Chapman and C. L. Smithhisler, **1995**. Codling moth, *Cydia pomonella*, (Lepidoptera: Tortricidae): is its sex pheromone multicomponent? *J. Chem. Ecol.*, **21**: 1065-1071.
- McElfresh, J. S. and J. G. Millar **1999(a)**. Sex pheromone of the common sheep moth, *Hemileuca eglanterina*, from the San Gabriel Mountains of California. *J. Chem. Ecol.*, **25**(4): 687-709.
- McElfresh, J. S. and J. G. Millar **1999(b)**. Sex pheromone of Nuttall's sheep moth, *Hemileuca nuttalli*, from the Eastern Sierra Nevada Mountains of California. *J. Chem. Ecol.*, **25**(4): 711-726.
- McElfresh, J. S. and J. G. Millar **1999(c)**. Sex attractant pheromone of saturniid moth, *Coloradia velda*. *J. Chem. Ecol.*, **25**(5): 1067-1078.
- McElfresh, J. S. and J. G. Millar, **1999(d)**. Geographic variation in sex pheromone blend of *Hemileuca electra* from southern California. *J. Chem. Ecol.*, **25**(11): 2505-2525.
- McElfresh, J. S., X. Chen, D. W. Ross and J. G. Millar, **2000**. Sex pheromone blend of the pandora moth (Lepidoptera: Saturniidae), an outbreak pest in pine forests (Pinaceae). *Can. Entomol.*, **132**: 775-787.
- McElfresh, J. S., J. G. Millar and D. Rubinoff, **2001(a)**. (E4,Z9)-Tetradecadienal, a sex pheromone for three north American moth species in the genus *Saturnia*. *J. Chem. Ecol.*, **27**(4): 791-806.
- McElfresh, J. S., A. M. Hammond and J. G. Millar, **2001(b)**. Sex pheromone components of the buck moth *Hemileuca maia*. *J. Chem. Ecol.*, **27**(7): 1409-1422.
- McElfresh, J. S. and J. G. Millar, **2008**. Sex pheromone of the saturniid moth, *Hemileuca burnsi*, from the western Mojave Desert of California. *J. Chem. Ecol.*, **34** (9): 1115-1124.
- McElfresh, J. S., J. A. Moreira, E. E. Grafton-Cardwell, D. H. Headrick, J. M. Heraty, M. Guillen and J. G. Millar, **2009**. Novel lepidopteran sex pheromone components from *Marmara gulosa* (Lepidoptera:

- Gracillariidae). *J. Ecocn. Entomol.*, **102**: 574-584.
- McElfresh, J. S. and J. G. Millar, 2024. Sex pheromone of the saturniid moth *Hemileuca nevadensis* from southern California. *J. Chem. Ecol.*, **50**: 330-337.
- McLaren, G. F. and D. M. Suckling, 1993. Pheromone trapping of orchard lepidopterous pests in Central Otago, New Zealand. *New Zealand J. Crop Horticul. Sci.*, **21**: 25-31.
- McLaughlin, J. R., E. R. Mitchell, M. Beroza and B. A. Bierl, 1975. Effect of *E-Z* concentration of 7-dodecenyl acetate on captures of four Noctuid species in pheromone traps. *J. Ga. Entomol. Soc.*, **10**: 338-341.
- McLaughlin, J. R. and R. R. Heath, 1989. Field trapping and observations of male velvetbean caterpillar moths and trapping of *Mocis* spp. (Lepidoptera: Noctuidae: Catacolinae) with calibrated formulations of sex pheromone. *Environ. Entomol.*, **18**: 933-938.
- McLeod, D. G. R., T. Nagai, A. N. Starratt, C. Bonenfant, E. W. Rud and S. Belloncik, 1982. Field and laboratory evaluation of a sex attractant for the white cutworm, *Euxoa scandens* (Lepidoptera: Noctuidae). *Can. Entomol.*, **114**: 1101-1103.
- McNair, C., G. Gries, and R. Gries 1999. Sex pheromone components of *Enarmonia formosana* (Lepidoptera: Tortricidae). *Can. Entomol.* **131**: 85-92.
- McQuate, G. T., A. Cossé, C. D. Sylva and J. A. MacKay, 2019. Field evaluation of a binary sex pheromone for sweetpotato vine borer (Lepidoptera: Crambidae) in Hawaii. *J. Insect Sci.*, **19**: iez008.
- Meijer, G. M., F. J. Ritter, C. J. Persoons, A. K. Minks and S. Voerman, 1972. Sex pheromone of summer fruit tortrix moth *Adoxophyes orana*: two synergistic isomers. *Science*, **175**: 1469-1470.
- Meng, X.-Z., 1983(a). Synthesis of the three active components of the sex pheromone of the masson pine caterpillar moth - (*Z,E*)-5,7-dodeca-dien-1-ol, acetate and propionate. *Kexue Tongbao*, **28**: 141.
- Meng, X.-Z. and H.-M. Wang, 1983(b). Synthesis and field experiment of sex attractant for pine caterpillar, *Dendrolimus tabulaeformis*. *Scientia Silvae Sinicae*, **19**: 137-140.
- Meng, W., W. Zhang, D. Zhang, J. Zheng, D. Huang, Q. Guo, and Y. Du, 2025. Geographic variation of the sex pheromone of *Mythimna separata* (Lepidoptera: Noctuidae) associated with overwintering sites. *Environ. Entomol.*, **54**: 1016-1026.

- Meyer, W. L., G. L. Debarr, C. W. Berisford, L. R. Barber and W. L. Roelofs, 1982. Identification of the sex pheromone of the webbing coneworm moth, *Dioryctria disclusa* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **11**: 986-988.
- Meyer, W. L., R. S. Cameron, A. Tamhankar, G. DeBarr, C. W. Berisford and W. L. Roelofs, 1984. Sex pheromone of the blister coneworm moth, *Dioryctria clarioralis* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **13**: 854-858.
- Meyer, W. L., G. L. Debarr, J. L. Hanula, B. Kovalev, R. S. Cameron, C. W. Berisford and W. L. Roelofs, 1986. Z-11-Hexadecenyl acetate, a sex pheromone component for the southern pine coneworm, *Dioryctria amatella* (Lepidoptera: Pyralidae). *Environ. Entomol.*, **15**: 316-320.
- Meyer, W. L., W. S. Cranshaw and T. D. Eichlin, 1988. Flight patterns of clearwing borers in Colorado based on pheromone trap captures. *Southwest. Entomol.*, **13**: 39-46.
- Meyer, W. L., and Cranshaw, W. S. 1994. Capture of clearwing borers (Lepidoptera: Sesiidae) with three synthetic attractants in Colorado. *Southwest Entomol.*, **19**: 71-76.
- Millar, J. G., E. W. Underhill, M. Giblin and D. Barton, 1987. Sex pheromone components of three species of *Semiothisa* (Geometridae), (Z,Z,Z)-3,6,9-heptadecatriene and two monoepoxydiene analogs. *J. Chem. Ecol.*, **13**: 1371-1383.
- Millar, J. G., M. Giblin, D. Barton and E. W. Underhill, 1990(a). (3Z,6Z,9Z)-Nonadecatriene and enantiomers of (3Z,9Z)-cis-6,7-epoxy-nonadecadiene as sex attractants for two geometrid and one noctuid moth species. *J. Chem. Ecol.*, **16**: 2153-2166.
- Millar, J. G., M. Giblin, D. Barton and E. W. Underhill, 1990(b). 3Z,6Z,9Z-Trienes and unsaturated epoxides as sex attractants for geometrid moths. *J. Chem. Ecol.*, **16**: 2307-2316.
- Millar, J. G., M. Giblin, D. Barton, A. Morrison and E. W. Underhill, 1990(c). Synthesis and field testing of enantiomers of 6Z,9Z-cis-3,4-epoxydienes as sex attractants for geometrid moths. *J. Chem. Ecol.*, **16**: 2317-2339.
- Millar, J. G., M. Giblin, D. Barton, D. A. Reynard, G. B. Neill and E. W. Underhill, 1990(d). Identification and field testing of female-produced sex pheromone components of the spring cankerworm, *Paleacrita vernata* Peck (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **16**: 3393-3409.

- Millar, J. G., M. GIBLIN, D. Barton and E. W. Underhill, 1991(a). Synthesis and field screening of chiral monounsaturated epoxides as lepidopteran sex attractants and sex pheromone components. *J. Chem. Ecol.*, **17**: 911-929.
- Millar, J. G., M. GIBLIN, D. Barton, J. W. Wong and E. W. Underhill, 1991(b). Sex attractants and sex pheromone components of noctuid moths *Euclidea cuspidea*, *Caenurgina distincta*, and geometrid moth *Eupithecia annulata*. *J. Chem. Ecol.*, **17**: 2095-2111.
- Millar, J. G., M. GIBLIN, D. Barton and E. W. Underhill, 1991(c). Chiral lepidopteran sex attractants: blends of optically active C₂₀ and C₂₁ diene epoxides as sex attractants for geometrid and noctuid moths (Lepidoptera). *Environ. Entomol.*, **20**(2): 450-457.
- Millar, J. G., M. GIBLIN, D. Barton and E. W. Underhill, 1992(a). Sex pheromone components of the geometrid moths *Lobophora nivigerata* and *Epirrhoe sperryi*. *J. Chem. Ecol.*, **18**: 1057-1068.
- Millar, J. and R. E. Rice, 1992(b). Reexamination of the female sex pheromone of the peach twig borer: field screening of minor constituents of pheromone gland extracts and of pheromone analogs. *J. Econ. Entomol.*, **85**: 1709-1716.
- Millar, J. G., A. E. Knudson, J. S. McElfresh, R. Gries, G. Gries and J. H. Davis, 1996. Sex attractant pheromone of the pecan nut casebearer (Lepidoptera: Pyralidae). *Bioorg. Medic. Chem.*, **4**: 331-339.
- Millar, J., J. S. McElfresh and F. de A. Marques, 2002. Unusual acetylenic sex pheromone of grape leaf folder (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **95**(4): 692-698.
- Millar, J. G., G. G. Grant, J. S. McElfresh, W. Strong, C. Rudolph, J. D. Stein and J. A. Moreira, 2005. (3Z,6Z,9Z,12Z,15Z)-Pentacosapentaene, a key pheromone component of the fir coneworm moth, *Dioryctria abietivorella*. *J. Chem. Ecol.*, **31**(5): 1229-1234.
- Millar, J. G., M. Hoddle, J. S. McElfresh, Y. Zou and C. Hoddle, 2008. (9Z)-9,13-Tetradecadien-11-ynal, the sex pheromone of the avocado seed moth, *Stenoma catenifer*. *Tetrahedron Lett.*, **49**: 4820-4823.
- Millar, J. G., J. S. McElfresh, C. Romero, M. Vila, N. Mari-Mena and C. Lopez-Vaamonde, 2010. Identification of the sex pheromone of a protected species, the Spanish moon moth *Graellsia isabellae*. *J. Chem. Ecol.*, **36**: 923-932.

- Millar, J. G., K. F. Haynes, A. T. Dossey, J. S. McElfresh and J. D. Allison, 2016. Sex attractant pheromone of the luna moth, *Actias luna* (Linnaeus). *J. Chem. Ecol.*, **42**: 869-876.
- Miller, J. R., T. C. Baker, R. T. Cardé and W. L. Roelofs, 1976. Reinvestigation of oak leaf roller sex pheromone components and the hypothesis that they vary with diet. *Science*, **192**: 140-143.
- Miller, J. R. and W. L. Roelofs, 1978. Gypsymoth responses to pheromone enantiomers as evaluated in a sustained-flight tunnel. *Environ. Entomol.*, **7**: 42-44.
- Miller, D. R., J. G. Millar, A. Mangini, C. M. Crowe and G. G. Grant, 2010. (3Z,6Z,9Z,12Z,15Z)-Pentacosapentaene and (Z)-11-hexadecenyl acetate: sex attractant blend for *Dioryctria amatella* (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **103**: 1216-1221.
- Minks, A. K. and S. Voerman, 1973(a). Sex pheromone of the summerfruit tortrix moth, *Adoxophyes orana*: trapping performance in the field. *Entomol. Exp. Appl.*, **16**: 541-549.
- Minks, A. K., W. L. Roelofs, F. J. Ritter and C. J. Persoons, 1973(b). Reproductive isolation of two tortricid moth species by different ratios of a two-component sex attractant. *Science*, **180**: 1073-1074.
- Minks, A. K., S. Voerman and M. Van de Vrie, 1976. A sex attractant for the cherry-bark tortrix moth, *Enarmonia formosana*. *Entomol. Exp. Appl.*, **19**: 301-302.
- Minks, A. K., S. Voerman and J. Theunissen, 1994. Improved sex attractant for the leek moth, *Acrolepiopsis assectella* Zell. (Lep., Acrolepiidae). *J. Appl. Entomol.*, **117**: 243-247.
- Minyailo, V. A., B. G. Kovalev, E. I. Kirov and S. A. Bakhvalov, 1976. Comparison of the attractiveness of disparlure, sex pheromone of females of *Porthetria dispar*, its *trans*-isomer and females of *Porthetria monacha* for males of the latter species. *Zool. Zhur.*, **55**: 1410-1411.
- Minyailo, V. A., B. G. Kovalev, E. I. Kirov and A. K. Minyailo, 1977. On the attractiveness of disparlure, sex pheromone of the *Porthetria dispar* females, for the *Zanclognatha lunaris* males. *Zool. Zhur.*, **56**: 309-310.
- Mitchell, E. R., J. C. Webb, A. H. Baumhover, R. W. Hines, J. W. Stanley, R. G. Endris, D. A. Lindquist and S. Masuda, 1972. Evaluation of cylindrical electric grids as pheromone traps for loopers and tobacco hornworms. *Environ. Entomol.*, **1**: 365-368.

- Mitchell, E. R. and J. H. Tumlinson, 1973(a). An attractant for males of *Spodoptera dolichos* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **66**: 917-918.
- Mitchell, 1973(b). Nocturnal activity of adults of three species of loopers, based on collections in pheromone traps. *Environ. Entomol.*, **2**: 1078-1080.
- Mitchell, E. R. and R. E. Doolittle, 1976. Sex pheromones of *Spodoptera exigua*, *S. eridania*, and *S. frugiperda*: bioassay for field activity. *J. Econ. Entomol.*, **69**: 324-326.
- Mitchell, E. R., J. H. Tumlinson and A. H. Baumhover, 1978. *Heliothis virescens*: attraction of males to blends of (Z)-9-tetradecen-1-ol formate and (Z)-9-tetradecenal. *J. Chem. Ecol.*, **4**: 709-716.
- Mitchell, E. R., H. Sugie and J. H. Tumlinson, 1983. *Spodoptera exigua*: capture of feral males in traps baited with blends of pheromone components. *J. Chem. Ecol.*, **9**: 95-104.
- Mitchell, E. R., J. H. Tumlinson and J. N. McNeil, 1985. Field evaluation of commercial pheromone formulations and traps using a more effective sex pheromone blend for the fall armyworm (Lepidoptera: Noctuidae). *J. Econ. Entomol.*, **78**: 1364-1369.
- Mitchell, E. R. and J. H. Tumlinson, 1994. Response of *Spodoptera exigua* and *S. eridania* (Lepidoptera: Noctuidae) males to synthetic pheromone and *S. exigua* females. *Florida Entomol.*, **77**(2): 237-247.
- Mizukoshi, T., 2006. Non-target species caught by synthetic sex pheromone trap of lepidopterous pests on apple trees in Hokkaido. *Jpn. J. Appl. Entomol. Zool.* **50** (3): 231-239.
- Mochida, O., G. S. Arida, S. Tatsuki and J. Fukami, 1984. A field test on a third component of the female sex pheromone of the rice striped stem borer, *Chilo suppressalis*, in the Philippines. *Entomol. Exp. Appl.*, **36**: 295-296.
- Mochizuki, F., T. Shibuya, T. Ihara and S. Wakamura, 1993. Electrophysiological responses of the male antenna to compounds found in the female sex pheromone gland of *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **28**: 489-496.
- Mochizuki, F., M. Takai, T. Shibuya, and S. Wakamura, 1994. Field trap responses of male *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae) to (Z,E)-9,12-tetradecadien-1-ol and (Z)-9-tetradecenyl acetate. *Appl. Entomol. Zool.*, **29**: 436-438.

- Mochizuki, F. and K. Fukumoto, 2003. Insect attractants for *Grapholita dimorpha*. *Jpn. Kokai Tokkyo Koho*, JP 2003-238311 [2002-42519].
- Mochizuki, F. and Y. Yumoto, 2009(a). Hexadecenal mixture-containing sex pheromones of *Pleuroptya ruralis*, and their use as attractants. *Jpn. Kokai Tokkyo Koho* JP 2009018994. [*SciFinder*. AN 2009:111797]
- Mochizuki, F. Y. Aoki, H. Sugie, J. Tabata, A. Takahashi and H. Higuchi, 2009(b). Sex attractant of the bean webworm, *Pleuroptya ruralis* (Cmbidae: Pyraustinae). *Jpn. J. Appl. Entomol. Zool.*, **53**: 53-56.
- Molnár, B. P., A. Tröger, T. B. Toshova, M. Subchev, E. J. van Nieukerken, J. C. Koster, G. Szöcs, M. Tóth and W. Francke, 2012. Identification of the female-produced sex pheromone of *Tischeria ekebladella*, an oak leafmining moth. *J. Chem. Ecol.*, **38**: 1298-1305.
- Molnár, P. B., C. Bognár, A. L. Erdei, T. Fujii, P. Vági, J. K. Jósvai and Z. Kárpáti 2018. Identification of the female-produced sex pheromone of an invasive greenhouse pest, the European pepper moth (*Duponchelia fovealis*) *J. Chem. Ecol.*, **44**: 257-267.
- Monti, L., B. Lalanne-Cassou, P. Lucas, C. Malosse and J.-F. Silvain, 1995. Differences in sex pheromone communication systems of closely related species: *Spodoptera latifascia* (Walker) and *S. descoinsi* Lalanne-Cassou & Silvain (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **21**: 641-660.
- Moreira, J. A., J. S. McElfresh and J. G. Millar, 2006. Identification, synthesis, and field testing of the sex pheromone of the citrus leafminer, *Phyllocnistis citrella*. *J. Chem. Ecol.*, **32**(1): 169-194.
- Moreira, J. A., T. Neppe, M. M. D. Paiva, A. M. Deobald, L. G. Batista-Pereira, M. W. Paixão and A. G. Corrêa, 2013. Studies towards the identification of the sex pheromone of *Thyrinteina arnobia*. *J. Braz. Chem. Soc.*, **24**: 1933-1941.
- Mori, K., H. Harada, P. Zagatti, A. Cork and D. R. Hall, 1991. Synthesis and biological activity of four stereoisomers of 6,10,14-trimethyl-2-pentadecanol, the female-produced sex pheromone of rice moth (*Corcyra cephalonica*). *Liebigs Ann. Chem.*, 259-267.
- Mori, K., T. Tashiro, B. Zhao, D. M. Suckling and A. M. El-Sayed, 2010. Pheromone synthesis. Part 243: Synthesis and biological evaluation of (3R,13R,1'S)-1'-ethyl-2'-methylpropyl 3,13-dimethylpentadecanoate, the major component of the sex

- pheromone of Paulownia bagworm, *Clania variegata*, and its stereoisomers. *Tetrahedron*, **66**: 2642-2653.
- Möttus, E., Z. Klukowski, I. Liblikas, M. Stachowiak and J. Viidalepp, 1998. Sex attractant dispensers for monitoring antler moth *Cerapteryx graminis* (L.) (Lep., Noctuidae): Effect of component ratio and dispenser type. *J. Appl. Entomol.*, **122**: 193-200.
- Mozūraitis, R., V. Buda, A.-K. Borg-Karlson, P. Ivinskis, V. Karalius, M. Laanmaa and D. Pleys 1998. New sex attractants and inhibitors for 17 moth species from the families Gracillariidae, Tortricidae, Yponomeutidae, Oecophoridae, Pyralidae and Gelechiidae. *J. Appl. Entomol.*, **122**: 441-452.
- Mozūraitis, R., V. Buda and J. Metleuski 1999(a). New sex attractants for five *Chamaesphecia* species (Lepidoptera, Sesiidae) from the Ukraine and Turkmenistan. *Z. Naturforsch.*, **54c**: 253-258.
- Mozūraitis, R., A.-K. Borg-Karlson, V. Buda and P. Ivinskis, 1999(b). Sex pheromone of the spotted tentiform leafminer moth *Phyllonorycter blancardella* (Fabr.) (Lep., Gracillariidae). *J. Appl. Entomol.*, **123**: 603-606.
- Mozūraitis, R., V. Buda, V. Jonusaite, A.-K. Borg-Karlson and R. Noreika, 2000. Sex pheromones of *Phyllonorycter acerifoliella* and *Ph. heegerella* and communication peculiarities in three species of leafmining moths. *Entomol. Exp. Appl.*, **94**: 15-23.
- Mozūraitis, R., V. Buda, I. Liblikas, C. R. Unelius and A.-K. Borg-Karlson, 2002. Parthenogenesis, calling behavior, and insect-released volatiles of leafminer moth *Phyllonorycter emberizaepenella*. *J. Chem. Ecol.*, **28**(6): 1191-1208.
- Mozūraitis, R., V. Karalius, V. Buda and A.-K. Borg-Karlson, 2006. Inter- and intraspecific activities of compounds driven from sex pheromone glands of currant borer, *Synanthedon tipuliformis* (Clerck) (Lepidoptera: Sesiidae). *Z. Naturforsch.*, **61 c**, 278-284.
- Mozūraitis, R. and V. Karalius, 2007. Identification of minor sex pheromone components of the poplar clearwing moth *Paranthrene tabaniformis* (Lepidoptera, Sesiidae). *Z. Naturforsch.*, **62c**: 138-142.
- Mozūraitis, R., I. Liblikas and R. Noreika, 2008. Sex pheromone communication of tentiform leaf-miners *Phyllonorycter insignitella* and *Ph. nigrescentella* from two related species groups. *Chemoecol.*, **18**: 171-176.
- Mozūraitis, R. and V. Buda, 2013. Intra- and interspecific activities

- of semiochemicals from the sex pheromone gland of the Welsh clearwing, *Synanthedon scoliaeformis*. *J. Chem. Ecol.*, **39**: 1066-1069.
- Muggleston, S. J. and S. P. Foster, 1989. Sustained-flight tunnel responses of male light brown apple moth to synthetic sex pheromone. *Physiol. Entomol.*, **14**: 443-449.
- Mukunthan, N. and B. Singaravelu, 2005. Evaluation of synthetic sex pheromone of sugarcane top borer (*Scirpophaga excerptalis* Walk). *Sugar Tech.*, **7**: 168-170.
- Murakami, Y., H. Sugie, T. Fukumoto and F. Mochizuki, 2005. Sex pheromone of *Grapholita dimorpha* Komai (Lepidoptera: Tortricidae), and its utilization for monitoring. *Appl. Entomol. Zool.*, **40**(3): 521-527.
- Muraki, Y. T. Taguri, R. Yamakawa and T. Ando, 2014. Synthesis and field evaluation of stereoisomers and analogues of 5-methylheptadecan-7-ol, an unusual sex pheromone component of the lichen moth, *Miltochrista calamine*. *J. Chem. Ecol.*, **40**: 250-258.
- Muraki, Y., R. Yamakawa, M. Yamamoto, H. Naka, A. Honma, J. Mappes, K. Suisto and T. Ando, 2017. GC/FT-IR analysis of novel 4,6,9-triene and 2,4,6,9-tetraene occurring in a female pheromone gland of *Arctia plantaginis* (Erebidae: Arctiinae). *Am. J. Chem. Anal.*, **8**: 645-656.
- Myerson, J., W. F. Haddon and E. L. Soderstrom, 1982. sec-Butyl (Z)-7-tetradecenoate. A novel sex pheromone component from the western grapeleaf skeletonizer, *Harrisina brillians*. *Tetrahedron Lett.*, **23**: 2757-2760.
- Nagaki, M., K. Miyata, T. Musashi, J. Kawakami, N. Ohya and H. Sagami, 2009. Insect pheromone-like activity of several isoprenoids against *Phyllonorycter ringoniella* (Matsumura). *Trans. Mater. Res. Soc. Jpn.*, **34**: 575-578.
- Nagano, M., Y. Sato, M. Sakai and H. Fujiwara, 1979. Cross-attractiveness of synthetic sex pheromones of lepidopterous species. *J. Takeda Res. Lab.*, **38**: 209-215.
- Nagayama, A., S. Wakamura, N. Taniai and N. Arakaki, 2006. Reinvestigation of sex pheromone components and attractiveness of synthetic sex pheromone of the pink borer, *Sesamia inferens* Walker (Lepidoptera: Noctuidae) in Okinawa. *Appl. Entomol. Zool.* **41** (3): 399-404.
- Nagy, A., I. Szarukán, B. Bohman, S. Szanyi, L. Kozák, A. Szilágyi,

- Z. Imrei, J. Vuts, E. Matula, Z. Varga and M. Tóth, 2023. Catches of *Euxoa tritici* in pheromone traps for *Anarsia lineatella* are due to the presence of (Z)-5-decenyl acetate as an impurity. *Entomol. Exp. Appl.*, **171**: 821-827.
- Naka, H., L. V. Vang, S. Inomata, T. Ando, T. Kimura, H. Honda, K. Tsuchida and H. Sakurai, 2003. Sex pheromone of the persimmon fruit moth, *Stathmopoda masinissa*: identification and laboratory bioassay of (4E,6Z)-4,6-hexadecadien-1-ol derivatives. *J. Chem. Ecol.*, **29**(11): 2447-2459.
- Naka, H., T. Nakazawa, M. Sugie, M. Yamamoto, Y. Horie, R. Wakasugi, Y. Arita, H. Sugie, K. Tsuchida and T. Ando, 2006. Synthesis and characterization of 3,13- and 2,13-octadecadienyl compounds for identification of the sex pheromone secreted by a clearwing moth, *Nokona pernix*. *Biosci. Biotechnol. Biochem.*, **69**(2): 508-516.
- Naka, H., S. Inomata, K. Matsuoka, M. Yamamoto, H. Sugie, K. Tsuchida, Y. Arita and T. Ando, 2007. Sex pheromone of two Melittini species, *Macroscelia japona* and *M. longipes*: identification and field attraction. *J. Chem. Ecol.*, **33** (3): 591-601.
- Naka, H., Y. Horie, F. Mochizuki, L. V. Vang, M. Yamamoto, T. Saito, T. Watarai, K. Tsuchida, Y. Arita and T. Ando, 2008. Identification of the sex pheromone secreted by *Synanthedon Hector* (Lepidoptera: Sesiidae). *Appl. Entomol. Zool.*, **43**: 467-474.
- Naka, H., M. Mochizuki, K. Nakada, N. D. Do, T. Yamauchi, Y. Arita and T. Ando, 2010. Female sex pheromone of *Glossosphecia romanovi* (Lepidoptera: Sesiidae): Identification and field attraction. *Biosci. Biotechnol. Biochem.*, **74**: 1943-1946.
- Naka, H., T. Suzuki, T. Watarai, Y. Horie, F. Mochizuki, A. Mochizuki, K. Tsuchida, Y. Arita and T. Ando, 2013. Identification of the sex pheromone secreted by *Synanthedon tenuis* (Lepidoptera: Sesiidae). *Appl. Entomol. Zool.*, **48**: 27-33.
- Naka, H., S. Kubota, F. Kuchiki, H. Shibasaki, M. Yamamoto, Y. Soejima, Y. Arita and T. Ando, 2018. Identification and field attraction of the female sex pheromone of a kiwifruit pest, *Nokona feralis* (Lepidoptera: Sesiidae). *Biosci. Biotechnol. Biochem.*, **82**: 1468-1472.
- Nam, I., J.-H. Park, J.-W. Lee, D. H. Lee, W. Jung, U. Lee and I.-K. Park, 2022. Field confirmation of (Z)-9-heptacosene and (3Z,6Z,9Z)-tricosatriene as key sex pheromone components of Korean *Conogethes punctiferalis* Guenée (Lepidoptera: Crambidae). *J. Econ. Entomol.*, **115**: 1976-1986.
- Nässig, W. A. and W. Thomas, 1991. *Cnephasia eculllyana* Réal, 1951,

- a species native to Central Europe (Lepidoptera, Tortricidae). *Nota lepid.*, **14**: 41-51.
- Nasu, Y., K. Tamashima, M. Shibao, S. Yoshimatsu and T. Naito, 2010. Discovery of *Carposina niponensis* Walsingham and Carposinids caught by synthetic sex pheromone trap for *C. sasakii* Matsumura in Japan (Lepidoptera: Carposinidae). *Jpn. J. Appl. Entomol. Zool.*, **54**: 115-126.
- Navarro, I., E. Mas, G. Fabriàs and F. Camps, 1997. Identification and biosynthesis of (*E,E*)-10,12-tetradecadienyl acetate in *Spodoptera littoralis* female sex pheromone gland. *Bioorg. Med. Chem.*, **5**: 1267-1274. (See Ph-4-1-102: pheromone biosynthesis)
- Neal, J. W., Jr, 1981. Timing insecticide control of rhododendron borer with pheromone trap catches of males. *Environ, Entomol.*, **10**: 264-266.
- Neal, J. W., Jr., J. A. Klun, B. A. Bierl-Leonhardt and M. Schwarz, 1982. Female sex pheromone of *Choristoneura parallela* (Lepidoptera: Tortricidae). *Environ, Entomol.*, **11**: 893-896.
- Neal, J. W., Jr. and T. D. Eichlin, 1983. Seasonal response of six male Sesiidae of woody ornamentals to clearwing borer (Lepidoptera: Sesiidae) lure. *Environ, Entomol.*, **12**: 206-209.
- Negishi, T., T. Ishiwatari, M. Uchida and S. Asano, 1979. A new synergist, 11-methyl-*cis*-9,12-tridecadienyl acetate, for the smaller tea tortrix sex attractant. *Appl. Entomol. Zool.*, **14**: 478-483.
- Nesbitt, B. F., P. S. Beevor, R. A. Cole, R. Lester and R. G. Poppi, 1973. Sex pheromones of two noctuid moths. *Nature*, **244**: 208-209.
- Nesbitt, B. F., P. S. Beevor, R. A. Cole, R. Lester and R. G. Poppi, 1975a. The isolation and identification of the female sex pheromones of the red bollworm moth, *Diparopsis castanea*. *J. Insect Physiol.*, **21**: 1091-1096.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester and V. A. Dyck, 1975b. Identification of the female sex pheromones of the moth, *Chilo suppressalis*. *J. Insect Physiol.*, **21**: 1883-1886.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester and V. A. Dyck, 1976. Identification of the female sex pheromone of the purple stem borer moth, *Sesamia inferens*. *Insect Biochem.*, **6**: 105-107.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester, M. Sternlicht and S. Goldenberg, 1977. Identification and synthesis of the female sex pheromone of the citrus flower moth, *Prays citri*. *Insect Biochem.*, **7**: 355-359.

- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester, J. C. Davies and K. V. Seshu Reddy, 1979(a). Components of the sex pheromone of the female spotted stalk borer, *Chilo partellus* (Swinhoe) (Lepidoptera:Pyralidae): identification and preliminary field trials. *J. Chem. Ecol.*, **5**: 153-163.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall and R. Lester, 1979(b). Female sex pheromone components of the cotton bollworm, *Heliothis armigera*. *J. Insect Physiol.*, **25**: 535-541.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall, R. Lester and J. R. Williams, 1980(a). Components of the sex pheromone of the female sugar cane borer, *Chilosacchariphagus* (Bojer) (Lepidoptera:Pyralidae). Identification and field trials. *J. Chem. Ecol.*, **6**: 385-394.
- Nesbitt, B. F., P. S. Beevor, D. R. Hall and R. Lester, 1980(b). (Z)-9-Hexadecenal: a minor component of the female sex pheromone of *Heliothis armigera* (Hübner) (Lepidoptera, Noctuidae). *Entomol. Exp. Appl.*, **27**: 306-308.
- Nesbitt, B. F., P. S. Beevor, A. Cork, D. R. Hall, R. Lester, B. W. Blair and J. Tannock, 1980(c). Identification of the female sex pheromone of the maize stalk borer, *Busseola fusca*: a preliminary note. *Trop. Pest Manag.*, **26**: 327.
- Nesbitt, B. F., P. S. Beevor, A. Cork, D. R. Hall, R. M. Murillo and H. R. Leal, 1985. Identification of components of the female sex pheromone of the potato tuber moth, *Scrobipalopsis solanivora*. *Entomol. Exp. Appl.*, **38**: 81-85.
- Nesbitt, B. F., P. S. Beevor, A. Cork, D. R. Hall, H. David and V. Vandagopal, 1986. The female sex pheromone of sugarcane stalk borer, *Chilo auricilius*. Identification of four components and field tests. *J. Chem. Ecol.*, **12**: 1377-1388.
- Newton, P. J., C. D. Thomas, V. C. Mastro and C. P. Schwalbe, 1993. Improved two-component blend of the synthetic female sex pheromone of *Cryptophlebia leucotreta*, and identification of an attractant for *C. peltastica*. *Entomol. Exp. Appl.*, **66**: 75-82.
- Nielsen, D. G., F. F. Purrington, J. H. Tumlinson, R. E. Doolittle and C. E. Yonce, 1975. Response of male clearwing moths to caged virgin females, female extracts, and synthetic sex attractants. *Environ. Entomol.*, **4**: 451-454.
- Nielsen, D. G., F. F. Purrington, R. L. Campbell, T. R. Wilmot, J. Capizzi and J. H. Tumlinson, 1978(a). Sex attractants for sequoia pitch moth and strawberry crown moth. *Environ. Entomol.*, **7**: 544-546.

- Nielsen, D. G. and F. F. Purrington, 1978(b). Field attraction of male *Podosesia syringae* and *P. aureocincta* to blends of synthetic compounds. *Environ, Entomol.*, **7**: 708-710.
- Nielsen, D. G., F. F. Purrington and G. F. Shambaugh, 1979. EAG and field responses of sesiid males to sex pheromones and related compounds. "Pheromone of the Sesiidae," ed. by J. W. Neal, Jr., ARR-NE-6, U. S. Department of Agriculture, Beltsville, Maryland, pp. 11-26. (see USDA)
- Niogret, J., A. Ekayanti and A. Zhang, 2022. Sex Pheromone of cocoa pod borer, *Conopomorpha cramerella*: field activity evaluation of pheromone formulations in an Indonesia plantation. *Insects*, **13**: 663.
- Niwa, C. G., L. L. Sower and G. E. Daterman, 1987. Chemistry and field evaluation of the sex pheromone of ponderosa pine tip moth, *Rhyacionia zozana* (Lepidoptera: Tortricidae). *Environ, Entomol.*, **16**: 1287-1290.
- Niwa, C. G. and L. L. Sower, 1992. Sex pheromone chemistry of the southwestern pine tip moth (Lepidoptera: Tortricidae). *Environ, Entomol.*, **21**: 1410-1413.
- Noguchi, H., Y. Tamaki and T. Yushima, 1979. Sex pheromone of the tea tortrix moth: isolation and identification. *Appl. Entomol. Zool.*, **14**: 225-228.
- Noguchi, H., Y. Tamaki, S. Arai, M. Shimoda and I. Ishikawa, 1981. Field evaluation of synthetic sex pheromone of the oriental tea tortrix moth, *Homona magnanima* Diakonoff (Lepidoptera: Tortricidae). *Jap. J. Appl. Entomol. Zool.*, **25**: 170-175.
- Noguchi, H., H. Sugie, Y. Tamaki and Y. Oomasa, 1985. Sex-pheromone components and related compounds released by virgin females of *Adoxophyes* sp. and *Adoxophyes orana fasciata* (Lepidoptera: Tortricidae). *Jap. J. Appl. Entomol. Zool.*, **29**: 278-283.
- Nojima, S., A. Classen, A. T. Groot and C. Scha, 2018. Qualitative and quantitative analysis of chemicals emitted from the pheromone gland of individual *Heliothis subflexa* females. *PLoS ONE*, **13**: e0202035.
- Novák, L., M. Tóth, J. Balla and C. Szántay, 1979. Sex pheromone of the cabbage armyworm, *Mamestra brassicae*: isolation, identification and stereocontrolled synthesis. *Acta Chim. Acad. Sci. Hung.*, **102**: 135-140.
- Novotný, Von J., J. Patočka, I. Hrdý and J. Vrkoč, 1990. Monitoring of *Tortrix viridana* L. by means of the synthetic sexual pheromone.

- Anz. Schädlin. Pflanz. Umwelt.*, **63**: 125-129.
- Nunez, S., J. J. De Vlieger, J. J. Rodriguez, C. J. Persoons and I. Scatoni, 2002. Sex pheromone of South American tortricid moth *Argyrotaenia sphaleropa*. *J. Chem. Ecol.*, **28**(2): 425-432.
- Ode11, T. M., C.-H. Xu, P. W. Schaefer, B. A. Leonhardt, D.-F. Yao and X.-D. Wu, 1992. Capture of gypsy moth, *Lymantria dispar* (L.), and *Lymantria mathura* (L.) males in traps baited with disparlure enantiomers and olefin precursor in the People's Republic of China. *J. Chem. Ecol.*, **18**: 2153-2159.
- Oetting, R. D., 1977. Immature stages and biology of *Chionodes psiloptera* a pest in bluegrass seed fields. *Pan-Pac. Entomol.*, **53**: 258-268.
- Ohmasa, Y., S. Wakamura, S. Kozaki, H. Sugie, M. Horiike, C. Hirano and S. Mori, 1991. Sex pheromone of the fruit-piercing moth, *Oraesia excavata* (Butler) (Lepidoptera: Noctuidae): isolation and identification. *Appl. Entomol. Zool.*, **26**: 55-62.
- Ohshima, M., Y. Kuwahara, S. Matsuyama and T. Suzuki, 1993. Identification of the sex pheromone of the European grain moth *Nemapogon granellus*: (3Z,13Z)-3,13-octadecadienyl acetate. *J. Pesticide Sci.*, **18**: 163-167.
- Ohta, K., S. Tatsuki, K. Uchiumi, M. Kurihara and J. Fukami, 1976. Structures of sex pheromones of rice stem borer. *Agric. Biol. Chem.*, **40**: 1897-1899.
- Oleander, A., D. R. Hall, D. P. Bray and J. P. J. Burman1, 2019. Identification of female sex pheromone for monitoring the barred tooth striped moth, *Trichopteryx polycommata*, a priority conservation species. *J. Chem. Ecol.*, **45**: 649-656.
- Olenici, N., I. Capuse, I. Olenici, I. Oprean and E. Mihalciuc, 2007. Non-target Lepidoptera species in pheromone traps baited with attractants for several tortricid moths. *Analele ICAS.*, **50**: 185-201.
- Oliver, J. E., J. C. Dickens, M. Zlotina, V. C. Mastro and G. I. Yurchenko 1999. Sex attractant of the rosy Russian gypsy moth (*Lymantria mathura* Moore). *Z. Naturforsch.*, **54c**: 387-394.
- Ono, T., R. E. Charlton and R. T. Cardé, 1990. Variability in pheromone composition and periodicity of pheromone titer in potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **16**: 531-542.
- Ono, T., P. Chouvalitwongporn and T. Saito, 1997. Comparison of the

- sex pheromone system between Japanese and Thai populations of the potato tuberworm moth, *Phthorimaea operculella* (Lepidoptera: Gelechiidae). *Appl. Entomol. Zool.*, **32**: 514-517.
- Ono, M., Y. Murata, J. M. S. Vento, W. S. Leal, E. F. Vilela, J. R. P. Para and A. J. Aires, **2001**. Dodecenyl acetate and dodecenol as sex attractants for citrus borer moth. *Jpn. Kokai Tokkyo Koho*, JP 2001-81002A.
- Ostrauskas, H., **2003**. Moths caught in pheromone traps for southern armyworm (*Spodoptera eridania* Cr.), fall armyworm (*S. frugiperda* Sm.), and Egyptian cotton leafworm (*S. littoralis* Bsd.) (Noctuidae, Lepidoptera) during 1999-2001 in Lithuania. *Acta Zool. Lit.*, **13**: 411-424.
- Ostrauskas, H., **2004**. Moths caught in pheromone traps for American white moth (*Hyphantria cunea* Dr.) (Archtiidae, Lepidoptera) in Lithuania during 2001. *Acta Zool. Litua.*, **14**: 66-74.
- Ostrauskas, H., P. Ivinskis and V. Bûda, **2010(a)**. Moths caught in traps during a survey of *Choristoneura fumiferana* (Clemens, 1865) - Lepidoptera, Tortricidae - in Lithuania. *Acta Zool. Lit.*, **20**: 156-161.
- Ostrauskas, H., P. Ivinskis and V. Bûda, **2010(b)**. Moths caught in traps during the survey of *Choristoneura Rosaceana* (Lepidoptera, Tortricidae) in Lithuania. *Acta Zool. Lit.*, **20**: 242-249.
- Palacio-Cortes, A. M., M. G. Fonseca and P. H. G. Zarbin, **2014**. Toward the identification of the sex pheromone of *Diatraea indigenella* Dyar & Heinrich (Lepidoptera: Crambidae): calling behavior and chemical structure of a major component. *Neotrop. Entomol.*, **43**: 526-531.
- Park, K. C., A. Cork, K. S. Boo and D. R. Hall, **1994**. Biological activity of female sex pheromone of the oriental tobacco budworm, *Helicoverpa assulta* (Guenée) (Lepidoptera: Noctuidae): electroantennography, wind tunnel observation and field trapping. *Korean J. Appl. Entomol.*, **33**(1): 26-32.
- Park, K. C., A. Cork and K. S. Boo, **1996**. Intrapopulational changes in sex pheromone composition during scotophase in oriental tobacco budworm, *Helicoverpa assulta* (Guenée) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **22**(6): 1201-1210.
- Park, S.-C. and S. P. Foster, **1998**. Comparative sex pheromone-associated behavior of *Planotortrix notophaea* and *P. octo* (Lepidoptera: Tortricidae). *J. Asia Pacific Entomol.*, **1**: 91-98.

- Park, J. H., K. S. Han, K. Mori and K. S. Boo, 2002. Right stereoisomers for sex pheromone components of the apple leafminer, *Lyonetia prunifoliella*, in Korea. *J. Chem. Ecol.*, **28**(12): 2515-2525.
- Park, H.-K., H.-R. Lee, C.-S. Jung and S.-H. Koh, 2019. Synergic effect of sex pheromone (7R,8S)-*cis*-7,8-epoxy-2-methyloctadec-17-ene on (+)-disparlure of the Asian gypsy moth, *Lymantria dispar*, in Korea. *J. Forest. Res.*, **30**: 1119-1122.
- Pasek, J. E. and M. E. Dix, 1989. Life history of a ponderosa pine coneworm, *Dioryctria auranticella* (Lepidoptera: Pyralidae). *J. Econ. Entomol.*, **82**: 879-885.
- Peltotalo, P. and T. Tuovinen, 1986. Specificity of pheromone preparates for lepidopterous pests. *Ann. Agric. Fenniae*, **25**: 139-146.
- Pelozuelo, L., C. Malosse, G. Genestier, H. Guenego and B. Frerot, 2004. Host-plant specialization in pheromone strains of the European cornborer *Ostrinia nubilalis* in France. *J. Chem. Ecol.*, **30**(2): 335-352.
- Peña, A., H. Arn, H.-R. Buser, S. Rauscher, F. Bigler, R. Brunetti, S. Maini and M. Tóth, 1988. Sex pheromone of European corn borer, *Ostrinia nubilalis*: polymorphism in various laboratory and field strains. *J. Chem. Ecol.*, **14**: 1359-1366.
- Peng, C.-L., P. Gu, J. Li, Q.-Y. Chen, C.-H. Feng, H.-H. Luo and Y.-J. Du, 2012. Identification and field bioassay of the sex pheromone of *Trichophysetis cretacea* (Lepidoptera: Crambidae). *J. Econ. Entomol.*, **105**: 1566-1572.
- Persoons, C. J., A. K. Minks, S. Voerman, W. L. Roelofs and F. J. Ritter, 1974. Sex pheromones of the moth, *Archips podana*: isolation, identification and field evaluation of two synergistic geometrical isomers. *J. Insect. Physiol.*, **20**: 1181-1188.
- Persoons, C. J., S. Voerman, P. E. J. Verwiel, P. J. F. Nooijen, W. J. Nooijen, F. J. Ritter and A. K. Minks, 1976(a). Sex pheromone of the potato tuberworm moth, *Phthorimaea operculella*: isolation and identification. *Med. Fac. Landbouww. Rijksuniv. Gent.*, **41**: 945-948.
- Persoons, C. J., S. Voerman, P. E. J. Verwiel, F. J. Ritter, W. J. Nooyen and A. K. Minks, 1976(b). Sex pheromone of the potato tuberworm moth, *Phthorimaea operculella*: isolation, identification and field evaluation. *Entomol. Exp. Appl.*, **20**: 289-300.
- Persoons, C. J., F. J. Ritter, D. Hainaut and J. P. Demoute, 1976(c).

- Sex pheromone of the false codling moth *Cryptophlebia* (= *Argyroplote*) *leucotreta* (Lepidoptera: Tortricidae) *trans*-8-dodecenyl acetate, a corrected structure. *Med. Fac. Landbouww. Rijksuniv. Gent.*, **41**: 937-943.
- Persoons, C. J., F. J. Ritter and W. J. Nooyen, **1977**. Sex pheromone of the false codling moth *Cryptophlebia leucotreta* (Lepidoptera: Tortricidae). Evidence for a two-component system. *J. Chem. Ecol.*, **3**: 717-722.
- Persoons, C. J., C. van der Kraan, W. J. Nooijen, F. J. Ritter, S. Voerman and T. C. Baker, **1981**. Sex pheromone of the beet armyworm, *Spodoptera exigua*: isolation, identification and preliminary field evaluation. *Entomol. Exp. Appl.*, **30**: 98-99.
- Persoons, C. J., S. Voerman, C. van der Kraan and C. J. H. Booij, **1984**. Sex pheromone of the leafroller moth *Syndemis musculana* (Tortricidae): preliminary identification and field evaluation. *Entomol. Exp. Appl.*, **35**: 101-103.
- Persoons, C. J., J. D. Vos, J. S. Yadav, A. R. Prasad, S. Sighomony, K. N. Jyothi and A. L. Prasuna, **1993**. Indo-Dutch cooperation on pheromones of Indian agricultural pest insects: sex pheromone components of *Diacrisia obliqua* (Arctiidae), *Achaea janata* (Noctuidae) and *Amsacta albistriga* (Arctiidae). *Bulletin OILB SROP*, **16**: 136-140.
- Piccardi, P., A. Capizzi, G. Cassani, P. Spinelli, E. Arsurra and P. Massardo, **1977**. A sex pheromone component of the old world bollworm *Heliothis armigera*. *J. Insect Physiol.*, **23**: 1443-1445.
- Picimbon, J. F., C. Gadenne, J. M. Bécard, J. L. Clément and L. Sreng, **1997**. Sex pheromone of the French black cutworm moth, *Agrotis ipsilon* (Lepidoptera: Noctuidae): identification and regulation of a multicomponent blend. *J. Chem. Ecol.*, **23**(1): 221-230.
- Pineda-Ríos, J. M., J. C. Tovar, J. M. Sámano, L. F. S. Abarca, R. M. L. Romero and E. de J. A. Suárez, **2016**. La composición y proporción de los componentes de la feromona sexual de *Hypsipyla grandella* Zeller (Lepidoptera: Pyralidae) varía dependiendo de la técnica utilizada para su aislamiento. *Entomotropica*, **31**: 172-185.
- Pires, E. V., A. L. Mendonça, L. Vanícková, N. S. J. Serra, R. C. C. da Silva, D. C. dos Santos, R. S. Campos, A. E. G. Sant'Ana and R. R. do Nascimento, **2016**. Identification and field and laboratory tests of the sex pheromone of *Cerconota anonella* Sepp. (Lepidoptera: Oecophoridae). *J. Appl. Entomol.*, **140**: 72-80.

- Pistillo, O. M., I. D'Isita, A. D. Palma and G. S. Germinara, 2024. Identification of the sex pheromone of the asparagus moth, *Parahypopta caestrum* (Lepidoptera, Cossidae). *J. Chem. Ecol.*, **50**: 321-329.
- Pletnev, V. A., V. L. Ponomarev, N. V. Vendilo, S. A. Kurbatov and K. V. Lebedeva, 2000. Search of the pheromone of Siberian silkworm *Dendrolimus superans sibiricus* (Lepidoptera: Lasiocampidae). *Agrokhimiya*, (6): 67-72. [*Chem. Abst.*, **134**: 263662]
- Polavarapu, S. and G. Lonergan, 1998. Sex pheromone of *Choristoneura parallela* (Lepidoptera: Tortricidae): components and development of a pheromone lure for population monitoring. *Environ. Entomol.*, **27**(5): 1242-1249.
- Ponder, B. M. and W. D. Seabrook, 1991. Sensitivity of blueberry leaf tier moths (Lepidoptera: Tortricidae) (Kearfott) to their own sex pheromone: mating bioassay, electroantennogram, and trap attractancy studies. *Can. Entomol.*, **123**: 231-238.
- Pope, M. M., L. K. Gaston and T. C. Baker, 1982. Composition, quantification, and periodicity of sex pheromone gland volatiles from individual *Heliothis virescens* females. *J. Chem. Ecol.*, **8**: 1043-1055.
- Pope, M. M., L. K. Gaston and T. C. Baker, 1984. Composition, quantification, and periodicity of sex pheromone volatiles from individual *Heliothis zea* females. *J. Insect Physiol.*, **30**: 943-945.
- Priesner, E., M. Jacobson, H. J. Bestmann, 1975. Structure-response relationships in noctuid sex pheromone reception. *Z. Naturforsch.*, **30c**: 283-293.
- Priesner, E., H.-J. Bestmann, O. Vostrowsky and P. Rösel, 1977. Sensory efficacy of alkyl-branched pheromone analogues in noctuid and tortricid Lepidoptera. *Z. Naturforsch.*, **32c**: 979-991.
- Priesner, E., H. Bogenschütz, W. Altenkirch and H. Arn, 1978. A sex attractant for the pine beauty moth, *Panolis flammea*. *Z. Naturforsch.*, **33c**: 1000-1002.
- Priesner, E., H. Bogenschütz and H. Arn, 1980(a). A sex attractant for the European fir budworm moth, *Choristoneura murinana*. *Z. Naturforsch.*, **35c**: 390-398.
- Priesner, E., 1980(b). Sex attractant system in *Polia pisi* L. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **35c**: 990-994.
- Priesner, E. and H. Bogenschütz, 1982(a). Further field experiments

- on the specificity of synthetic sex-attractants in *Choristoneura murinana*. *Z. Angew. Entomol.*, **94**: 331-351.
- Priesner, E., W. Altenkirch, W. Baltensweiler and H. Bogenschütz, 1982(b). Evaluation of (Z)-5-decen-1-ol as an attractant for male larch casebearer moths, *Coleophora laricella*. *Z. Naturforsch.*, **37c**: 953-966.
- Priesner, E. and M. Schroth, 1983. Supplementary data on the sex attractant system *Panolis flammea*. *Z. Naturforsch.*, **38c**: 870-873.
- Priesner, E., C.M. Naumann and J. Stertenbrink, 1984(a). Specificity of synthetic sex-attractants in *Zygaena* moths. *Z. Naturforsch.*, **39c**: 841-844.
- Priesner, E., 1984(b). Analysis of a sex-attractant system in the noctuid moth *Rhyacia baja* Schiff. *Z. Naturforsch.*, **39c**: 845-848.
- Priesner, E., 1984(c). The pheromone receptor system of male *Eulia ministrana* L., with notes on other Cnephasiini moths. *Z. Naturforsch.*, **39c**: 849-852.
- Priesner, E., H. Bogenschütz, R. Albert, D. W. Reed and M. D. Chisholm, 1984(d). Identification and field evaluation of a sex pheromone of the European pine moth. *Z. Naturforsch.*, **39c**: 1192-1195.
- Priesner, E., 1985(a). Specificity of sexual attractants in *Xestia triangulum* Hufn. and *X. ditrapezium* Schiff. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **40c**: 939-942.
- Priesner, E., 1985(b). Inhibitors of sexual attraction in the moth *Agrotis exclamationis*. *Z. Naturforsch.*, **40c**: 943-945.
- Priesner, E., 1985(c). Species-specific sex-attractants for male *Diachrysia chrysitis* (L.) and *D. tutti* (Koster.) (Lepidoptera, Noctuidae: Plusiinae). *Mitt. Schweiz. Entomol. Ges.*, **58**: 373-391.
- Priesner, E., G. Dobler and S. Voerman, 1986(a). Synergism of positional isomers in sex-attractant systems of clearwing moths (Sesiidae). *Entomol. Exp. Appl.*, **41**: 311-313.
- Priesner, E., P. Witzgall and S. Voerman, 1986(b). Field attraction response of raspberry clearwing moths, *Pennisetia hylaeiformis* Lasp. (Lepidoptera: Sesiidae), to candidate pheromone chemicals. *J. Appl. Entomol.*, **102**: 195-210.
- Priesner, E., 1987(a). (Z)-5-Dodecen-1-ol, another inhibitor of pheromonal attraction in *Coleophora laricella*. *Z. Naturforsch.*,

42c: 1349-1351.

- Priesner, E., 1987(b). (Z)-3-Tetradecenyl acetate as a sex-attractant component in Gelechiinae and Anomologinae (Lepidoptera: Gelechiidae). *Z. Naturforsch.*, **42c**: 1352-1354.
- Priesner, E., H. Bogenschütz and C. Löfstedt, 1988(a). Secondary sex pheromone components of *Choristoneura murinana* Hb. (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **43c**: 918-928.
- Priesner, E., 1988(b). (Z)-3- and (Z)-5-Decenyl acetates, sex-attractant components for male *Eustrotia uncula* Cl. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **43c**: 961-962.
- Priesner, E., 1989(a). Two-component sexual attractant for male *Batrachedra pinicolella* (Zell.) (Lepidoptera: Batrachedridae). *Z. Naturforsch.*, **44c**: 1061-1062.
- Priesner, E., D. W. Reed, E. W. Underhill and H. Bogenschütz, 1989(b). (Z,Z)-7,9-Dodecadienyl acetate, sex pheromone of *Epinotia tedella* Clerck (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **15**: 2457-2464.
- Priesner, E. and K. Spatenka, 1990. The distribution of *Pennisetia bohémica* Kralicek & Povolny, 1974 (Lepidoptera: Sesiidae) in central Europe as determined by pheromone trapping. *Mitt. Schweiz. Entomol. Ges.*, **63**: 87-98.
- Priesner, E. and Q.-H. Zhang, 1991. Field response of male eastern larch casebearer *Coleophora dahurica* Flkv. (Lep., Coleophoridae) to synthetic sex-attractant and attraction-inhibitors. *J. Appl. Entomol.*, **112**: 321-326.
- Priesner, E., 1993. Pheromone test an einer südbayerischen Population von *Synanthedon soffneri* Spatenka, 1983 (Lepidoptera: Sesiidae). *NachrBl. Bayer. Entomol.*, **42**(4): 97-107.
- Qian, C., Z. Su, W. Xie, X. Dong, Y. Li, P. Yuan, S. Wang, X. Wen, C. Wang and T. Ma, 2024. Structure, synthesis, and bioassays of sex pheromone for *Pyrausta machaeralis* (Lepidoptera: Crambidae). *J. Agric. Food Chem.*, **72**: 3904-3912.
- Pu, G.-Q., M. Yamamoto, Y. Takeuchi, H. Yamazawa and T. Ando 1999. Resolution of epoxydienes by reversed-phase chiral HPLC and its application to stereochemistry assignment of mulberry looper sex pheromone. *J. Chem. Ecol.*, **25**(5): 1151-1162.
- Qin, X.-R., T. Ando, M. Yamamoto, M. Yamashita, K. Kusano and H. Abe, 1997. Resolution of pheromonal epoxydienes by chiral HPLC,

- stereochemistry of separated enantiomers, and their field evaluation. *J. Chem. Ecol.*, **23**(5): 1403-1417.
- Qu, H., Y. Long, L. Chen, Z. Luo, H. Chen, X. Wang, L. Long, J. Tian, T. Jing and L. Chen, 2024. Metabolomic profiling identifies metabolites in the pheromone glands of *Agriophara rhombata* associated with the synthesis and release of female pheromone signals. *Heliyon*, **10**: e40768.
- Quartey, G. K. and T. H. Coaker, 1993. Role of sex pheromone components in the orientation behaviour of *Ephestia cautella*. *Entomol. Exp. Appl.*, **66**: 237-245.
- Quero, C., F. Camps and A. Guerrero, 1995. Behavior of processionary males (*Thaumetopoea pityocampa*) induced by sex pheromone and analogs in a wind tunnel. *J. Chem. Ecol.*, **21**: 1957-1969.
- Quero, C., D. A. Malo, G. Fabriàs, F. Camps, P. Lucas, M. Renou and A. Guerrero, 1997. Reinvestigation of female sex pheromone of processionary moth (*Thaumetopoea pityocampa*): No evidence for minor components. *J. Chem. Ecol.*, **23**(3): 713-726.
- Quero, C., J. Bau, A. Guerrero, M. Breuer, A. D. Loof, H.-G. Kontzog and F. Camps, 2003. Sex pheromone of the oak processionary moth *Thaumetopoea processionea*. Identification and biological activity. *J. Agric. Food Chem.*, **51**: 2987-2991.
- Radžiutė, S., S. Orlovskytė, J. Būdienė, L. Blažytė-Čereškienė, G. Bumbulytė and V. Būda, 2023. Stored product pest *Nemapogon variatella* (Clemens): identification of sex pheromone, putative antagonist and mitotype characterization. *J. Appl. Entomol.*, **147**: 857-867.
- Rahmani, R., D. Carrasco, G. P. Svensson, H. Roweck, N. Ryrholm, M. C. Larsson and E. Hedenstroem, 2020. Identification and Synthesis of Putative Pheromone Components of the Threatened Salt Marsh Bagworm Moth, *Whittleia retiella* (Lepidoptera: Psychidae) *J. Chem. Ecol.*, **46**: 115-127.
- Rahn, R. 1979. The leek moth, *Acrolepiopsis* (*Acrolepia*) *assectella* Zell.: possibility of generalization of agricultural advertisements with sex traps. *Ann. Zool. Ecol. Anim.*, **11**: 611-616.
- Raina, A. K., J. A. Klun, M. Schwarz, A. Day, B. A. Leonhardt and L. W. Douglass, 1986(a). Female sex pheromone of the melonworm, *Diaphania hyalinata* (Lepidoptera: Pyralidae), and analysis of male responses to pheromone in a flight tunnel. *J. Chem. Ecol.*, **12**: 229-237.

- Raina, A. K., J. A. Klun, J. D. Lopez and B. A. Leonhardt, 1986(b). Female sex pheromone of *Heliothis phloxiphaga* (Lepidoptera: Noctuidae): chemical identification, male behavioral response in the flight tunnel, and field tests. *Environ. Entomol.*, **15**: 931-935.
- Ramachandran, R., P. Caballero and Z. R. Khan, 1990. Pheromone components of rice leaffolders (LF) *Cnaphalocrocis medinalis* and *Marasmia patnalis*. *Int. Rice Res. New.*, **15**: 25-26.
- Raman, K. V., 1982. Field trials with the sex pheromone of the potato tuberworm. *Environ. Entomol.*, **11**: 367-370.
- Ramaswamy, S. B., S. A. Randle and W. K. Ma, 1985. Field evaluation of the sex pheromone components of *Heliothis virescens* (Lepidoptera: Noctuidae) in cone traps. *Environ. Entomol.*, **14**: 293-296.
- Rao, A. G., K. Krishnaiah, D. R. Hall, A. Cork and D. D. R. Reddy, 1993. Evaluation of sex pheromone components of rice leaf folder, *Cnaphalocrocis medinalis* Guenee. *Current Sci.*, **65**: 355-356.
- Rao, A. G., D. D. R. Reddy, K. Krishnaiah, P. S. Beevor, A. Cork and D. R. Hall, 1995. Identification and field optimisation of the female sex pheromone of the rice leaffolder, *Cnaphalocrocis medinalis* in India. *Entomol. Exp. Appl.*, **74**: 195-200.
- Rauscher, S., H. Arn and P. Guerin, 1984. Effects of dodecyl acetate and Z-10-tridecenyl acetate on attraction of *Eupoecilia ambiguella* males to the main sex pheromone component, Z-9-dodecenyl acetate. *J. Chem. Ecol.*, **10**: 253-264.
- Razov, J., K. A. Efetov, K. Franin, T. B. Toshova and M. A. Subchev, 2017. The application of sex pheromone traps for recording the Procridinae fauna (Lepidoptera: Zygaenidae) in Croatia. *Entomol. Gaz.*, **68**: 49-53.
- Read, J. S., P. H. Hewitt, F. L. Warren and A. C. Myberg, 1974. Isolation of the sex pheromone of the moth *Argyroplote leucotreta*. *J. Insect Physiol.*, **20**: 441-450.
- Read, J. S. and P. S. Beevor, 1976. Analytical studies on the sex pheromone complex of *Ephestia cautella* (Walker) (Lepidoptera: Phycitidae). *J. Stored Prod. Res.*, **12**: 55-57.
- Read, J. S. and C. P. Haines, 1979. Secondary pheromone components and synergism in stored-products Phycitinae. *J. Chem. Ecol.*, **5**: 251-257.
- Reed, D. W., M. D. Chisholm and E. W. Underhill, 1984. Sex attractant

- for three species of the genus *Oncocnemis*: *O. chandleri* (Grt.), *O. cibalis* (Grt.), and *O. mackiei* (B. & Benj.) (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **10**: 1701-1706.
- Reed, D. W. and M. D. Chisholm, 1985(a). Field trapping of three *Epinotia* species with (Z,Z)-7,9-dodecadienyl acetate (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **11**: 1389-1398.
- Reed, D. W. and M. D. Chisholm, 1985(b). Attraction of moth species of Tortricidae, Gelechiidae, Geometridae, Drepanidae, Pyralidae, and Gracillariidae families to field traps baited with conjugated dienes. *J. Chem. Ecol.*, **11**: 1645-1657.
- Reed, D. W., E. W. Underhill and E. M. Giblin, 1987. Attraction of sphingid moths (Lepidoptera: Sphingidae) to 10,12-hexadecadienyl aldehydes and acetates: evidence of pheromone components. *J. Chem. Ecol.*, **13**: 931-942.
- Redfern, R. E., E. Cantu, W. A. Jones and M. Jacobson, 1971. Response of the male southern armyworm in a field cage to prodenialure A and prodenialure B. *J. Econ. Entomol.*, **64**: 1570-1571.
- Renou, M., C. Descoins, E. Priesner, M. Gallois and M. Lettère, 1979. Z-7-Tetradecenal, the main component of the sex pheromone of the olive moth *Prays oleae*. *C. R. Hebd. Seances Acad. Sci., Ser. D*, **288**: 1559-1562.
- Renou, P. M., Ch. Descoins, J. Y. Lallemand, E. Priesner, M. Lettère and M. Gallois, 1980. (E)-3-dodecenyl acetate, main component of the sex pheromone of the sugar beet moth, *Scrobipalpa ocellatella* Boyd. (Lepidoptera, Gelechiidae). *Z. Angew. Entomol.*, **90**: 275-289.
- Renou, M., C. Descoins, E. Priesner, M. Gallois and M. Lettère, 1981(a). A study of the sex pheromone of the leek moth, *Acrolepiopsis assectella* (Lepidoptera: Acrolepiidae). *Entomol. Exp. Appl.*, **29**: 198-208.
- Renou, M., B. Lalanne-Cassou, B. Frérot, M. Gallois and C. Descoins, 1981(b). Composition of the pheromonal secretion produced by virgin females of *Mamestra (Polia) pisi* L. (Lepidoptera, Noctuidae, Hadeninae). *C. R. Acad. Sci., Ser. 3*, **292**: 1117-1119.
- Renou, M., B. Lalanne-Cassou, J.-C. Doré and M.-L. Milat, 1988. Electroantennographic analysis of sex pheromone specificity in neotropical Catocalinae (Lepidoptera: Noctuidae): a multivariate approach. *J. Insect Physiol.*, **34**: 481-488.
- Renou, M., 1991. Sex pheromone reception in the moth, *Mamestra thalassina*. Characterization and distribution of two types of

- olfactory hairs. *J. Insect Physiol.*, **37**: 617-626.
- Rhainds, M., G. Gries, J.-X. Li, R. Gries, K. N. Slessor, C. M. Chinchilla and A. C. Oehlschlager, 1994. Chiral esters: sex pheromone of the bagworm, *Oiketicus kirbyi* (Lepidoptera: Psychidae). *J. Chem. Ecol.*, **20**: 3083-3096.
- Riba, M., J. A. Rosell, M. Eizaguirre, R. Canela and A. Guerrero, 1990. Identification of a minor component of the sex pheromone of *Leucoptera malifoliella* (Lepidoptera, Lyonetiidae). *J. Chem. Ecol.*, **16**: 1471-1483.
- Rice, R. E., and R. A. Jones, 1975. Peach twig borer: field use of a synthetic sex pheromone. *J. Econ. Entomol.*, **68**: 358-360.
- Ritter, F. J. and C. J. Persoons, 1975. Recent development in insect pheromone research, in particular in the Netherlands. *Neth. J. Zool.*, **25**: 261-275.
- Robacker, D. C. and K. J. R. Johnson, 1989. Pheromone emission and blend percentages in *Eoreuma loftini* determined by two methods. *J. Chem. Ecol.*, **15**: 105-120.
- Rocchini, L. A., B. S. Lindgren and R. G. Bennett, 2003. Douglas-fir pitch moth, *Synanthedon novaroensis* (Lepidoptera: Sesiidae) in North-Central British Columbia: flight period and the effect of trap type and pheromone dosage on trap catches. *Environ. Entomol.*, **32**: 208-213.
- Rodríguez, S. A. and M. A. Nazareno 2018. Identification of the sex pheromone of female *Eurata patagiata*. *Southwestern Entomol.*, **43**: 81-90.
- Roelofs, W. L. and H. Arn. 1968(a). Sex attractant of the red-banded leafroller moth. *Nature*, **219**: 513.
- Roelofs, W. L. and A. Comeau, 1968(b). Sex pheromone perception. *Nature*, **220**: 600-601.
- Roelofs, W. L., A. Comeau and R. Selle, 1969(a). Sex pheromone of the oriental fruit moth. *Nature*, **224**: 723.
- Roelofs, W. L. and A. Comeau, 1969(b). Sex pheromone specificity, taxonomic and evolutionary aspects in Lepidoptera. *Science*, **165**: 398-400.
- Roelofs, W. L. and A. Comeau, 1970(a). Lepidopterous sex attractants discovered by field screening tests. *J. Econ. Entomol.*, **63**: 969-974.

- Roelofs, W. L. and J. P. Tette, 1970(b). Sex pheromone of the oblique-banded leaf roller moth. *Nature*, **226**: 1172.
- Roelofs, W. L. and A. Comeau, 1971(a). Sex attractants in Lepidoptera. "Pesticide Chemistry, vol. III, Chemical Releasers in Insects," ed. by A. S. Tobot, Cordon of Beach, New York, N. Y., pp. 91-114.
- Roelofs, W. L. and R. T. Cardé, 1971(b). Hydrocarbon sex pheromone in tiger moths (Arctiidae). *Science*, **171**: 684-686.
- Roelofs, W. L., J. P. Tette, E. F. Taschenberg and A. Comeau, 1971(c). Sex pheromone of the grape berry moth: identification by classical and electroantennogram methods, and field tests. *J. Insect Physiol.*, **17**: 2235-2243.
- Roelofs, W. L., R. Cardé, G. Benz and G. von Salis, 1971(d). Sex attractant of the larch bud moth found by electroantennogram method. *Experientia*, **27**: 1438-1439.
- Roelofs, W., A. Comeau and G. M. Hill, 1971(e). Sex attractant of the codling moth: characterization with electroantennogram technique. *Science*, **174**: 297-299.
- Roelofs, W. L., R. T. Cardé, R. J. Bartell and P. G. Tierney, 1972. Sex attractant trapping of the European corn borer in New York. *Environ. Entomol.*, **1**: 606-608.
- Roelofs, W., J. Kochansky, R. Cardé, H. Arn and S. Rauscher, 1973. Sex attractant of the grape vine moth, *Lobesia botrana*. *Mitt. Schweiz. Entomol. Ges.*, **46**: 71-73.
- Roelofs, W. L. and R. T. Cardé, 1974(a). Oriental fruit moth and lesser appleworm attractant mixtures refined. *Environ. Entomol.*, **3**: 586-588.
- Roelofs, W. L. and R. T. Cardé, 1974(b). Sex pheromones in the reproductive isolation of lepidopterous species. "North-Holland Research Monograph Frontiers of Biology, vol. 32, Pheromones," ed. by M. C. Birch, North-Holland Publishing Co., Amsterdam, pp. 96-114.
- Roelofs, W. L., A. S. Hill, R. T. Cardé and T. C. Baker, 1974(c). Two sex pheromone components of tobacco budworm moth, *Heliothis virescens*. *Life Sci.*, **14**: 1555-1562.
- Roelofs, W. L., A. Hill, R. Cardé, J. Tette, H. Madsen and J. Vakenti, 1974(d). Sex pheromones of the fruit tree leafroller moth, *Archips argyrospilus*. *Environ. Entomol.*, **3**: 747-751.
- Roelofs, W., A. Hill and R. Cardé, 1975(a). Sex pheromone components

- of the redbanded leafroller, *Argyrotaenia velutinana* (Lepidoptera: Tortricidae). *J. Chem. Ecol.*, **1**: 83-89.
- Roelofs, W., J. Kochansky, E. Anthon, R. Rice and R. Cardé, 1975(b). Sex pheromone of the peach twing borer moth (*Anarsia lineatella*). *Environ, Entomol.*, **4**: 580-582.
- Roelofs, W. L., J. P. Kochansky, R. T. Carde, G. G. Kennedy, C. A. Henrick, J. N. Labovitz and V. L. Corbin, 1975(c). Sex pheromone of the potato tuberworm moth, *Phthorimaea operculella*. *Life Sci.*, **17**: 699-706.
- Roelofs, W., A. Hill, A. Cardé, R. Cardé, H. Madsen and J. Vakenti, 1976(a). Sex pheromone of the European leafroller, *Archips rosanus*. *Environ, Entomol.*, **5**: 362-364.
- Roelofs, W., A. Carde, A. Hill and R. Carde, 1976(b). Sex pheromone of the threelined leafroller, *Pandemis limitata*. *Environ, Entomol.*, **5**: 649-652.
- Roelofs, W. L., R. F. Lagier and S. C. Hoyt, 1977(a). Sex pheromone of the moth, *Pandemis pyrusana*. *Environ, Entomol.*, **6**: 353-354.
- Roelofs, W. L., W. H. Reissig and R. W. Weires, 1977(b). Sex attractant for the spotted tentiform leaf miner moth, *Lithocolletis blancardella*. *Environ, Entomol.*, **6**: 373-374.
- Roelofs, W. L., A. S. Hill, C. W. Berisford and J. F. Godbee, 1979. Sex pheromone of the subtropical pine tip moth, *Rhyacionia subtropica*. *Environ, Entomol.*, **8**: 894-895.
- Roelofs, W. L., A. J. Tamhankar, A. Comeau, A. S. Hill and E. F. Taschenberg, 1980. Moth activity periods and identification of the sex pheromone of the uglynest caterpillar, *Archips cerasivoranus*. *Ann. Entomol. Soc. Am.*, **73**: 631-634.
- Roelofs, W. L. and R. L. Brown, 1982(a). Pheromones and evolutionary relationships of Tortricidae. *Ann. Rev. Ecol. Syst.*, **13**: 395-422.
- Roelofs, W. L., A. S. Hill, C. E. Linn, J. Meinwald, S. C. Jain, H. J. Herbert and R. F. Smith, 1982(b). Sex pheromone of the winter moth, a geometrid with unusually low temperature precopulatory responses. *Science*, **217**: 657-659.
- Roelofs, W. L. and L. Bjostad, 1984. Biosynthesis of lepidopteran pheromones. *Bioorg. Chem.*, **12**: 279-298.
- Rojas, J. C., L. Cruz-López, E. A. Malo, O. Diaz-Gomez, G. Calyecac and J. C. Tovar, 2006. Identification of the sex pheromone of *Copitarsia decolora* (Lepidoptera: Noctuidae). *J. Econ. Entomol.*,

99: 797-802.

- Rossi, R., A. Carpita, P. Piccardi, C. Tonini, G. Guglielmetti, A. Niccoli and G. A. Fenili, 1982. Identification of (Z)-11-tetradecen-1-yl acetate as a component of *Epichoristodes acerbellasex* pheromone. *Frust. Entomol. Nuova serie*, **2**: 113-120.
- Rothschild, G. H. L., 1975. Attractants for monitoring *Pectinophora scutigera* and related species in Australia. *Environ. Entomol.*, **4**: 983-985.
- Rothschild, G. H. L., 1978. Attractants for *Heliothis armigera* and *H. punctigera*. *J. Aust. Entomol. Soc.*, **17**: 389-390.
- Rothschild, G. H. L., B. F. Nesbitt, P. S. Beevor, A. Cork, D. R. Hall and R. A. Vickers, 1982. Studies of the female sex pheromone of the native budworm, *Heliothis punctiger*. *Entomol. Exp. Appl.*, **31**: 395-401.
- Rotundo, G. and E. Tremblay, 1982. Preliminary observations on the sexual pheromone of *Opogona sacchari* (Bojer) (Lepidoptera, Hieroxestidae). *Boll. Lab. Entomol. Agrar. F.S.*, **39**: 123-132.
- Rotundo, G., 1983. Identification of the primary component of the sex pheromone of *Blepharita solieri* (Bdv.) (Lepidoptera, Noctuidae). *Boll. Lab. Entomol. Agrar. F.S.*, **40**: 179-186.
- Rotundo, G., R. Giacometti and A. Rotundo, 1984. Identification of the sex attractants of chestnut tortricids by means of electroantennographic and field studies. *Boll. Lab. Entomol. Agrar. F.S.*, **41**: 211-223.
- Rotundo, G. and E. Tremblay, 1985(a). Field evaluation of female sex pheromone of the artichoke moth, *Gortyna xanthenes*. *Entomol. Exp. Appl.*, **38**: 201-204.
- Rotundo, G. and R. Giacometti, 1985(b). Tortricidae Olethreutini caught with attractant traps in a chestnut-growing area. *Boll. Lab. Entomol. Agrar. F.S.*, **42**: 167-189.
- Rotundo, G. and R. Giacometti, 1988. Individualization of an attractive mixture for males of *Cydia fagiglandana* Z. (Lep. Tortricidae) by means of a field study. *Boll. Lab. Entomol. Agrar. F.S.*, **45**: 81-97.
- Rotundo, G., R. Giacometti and A. de Cristofaro, 1991. Individual mixtures attractive to males of *Pammene fasciana* L. (Lep., Tortricidae) using electroantennograms and field studies. *Boll. Lab. Entomol. Agrar. F.S.*, **48**: 89-104.

- Rotundo, G., G. S. Germinara and A. de Cristofaro, 2004. Chemical electrophysiological and behavioral investigations on the sex pheromone of lackey moth, *Malacosoma neustria*. *J. Chem. Ecol.* **30**: 2057-2069.
- Rudziński, K. J., D. Staszek, M. Asztemborska, L. Sukovata, J. Raczko, M. Cieślak, A. Kolk and R. Szmigielski, 2022. Newly discovered components of *Dendrolimus pini* sex pheromone. *Insects*, **13**: 1063.
- Ruiz, K. P., A. Bruce, N. E. Chérémond, C. A. Stratton, E. G. Murrell, S. Gillette and W. R. Morrison III, 2022. Field trapping and flight capacity of *Eucosma giganteana* (Riley) (Lepidoptera: Tortricidae) in response to behaviorally active congeneric semiochemicals in novel silflower agroecosystems. *Insects*, **13**: 350.
- Rule, G. S. and W. L. Roelofs, 1989. Biosynthesis of sex pheromone components from linolenic acid in arctiid moths. *Arch. Insect Biochem. Physiol.*, **12**: 89-97.
- Ryall, K., P. J. Silk, J. Wu, P. Mayo, M. A. Lemay and D. MaGee, 2010. Sex pheromone chemistry and field trapping studies of the elm spanworm *Ennomos subsignaria* (Hübner) (Lepidoptera: Geometridae). *Naturwissenschaften*, **97**: 717-724.
- Safonkin, A. F. and V. V. Buleza, 1988. Role of sexual feromones in interspecific isolation of leafrollers (Lepidoptera: Tortricidae). *Zhural Obschiy Biologiya (J. General Biology)*, **49**: 396-400.
- Safonkin, A. F. and T. A. Triseleva, 2003. Identification of the sex pheromone components of the apple surface eating tortrix *Archips podana* Scop. (Lepidoptera: Tortricidae). *Agrokhimiya*, (3): 62-64.
- Saglio, P., E. Priesner, C. Descoins and M. Gallois, 1977(a). A synthetic sex attractant for *Sparganothis pilleriana* (Schiffermüller). *C. R. Hebd. Seances Acad. Sci., Ser. D*, **284**: 2007-2010.
- Saglio, P., C. Descoins, M. Gallois, M. Lettere, D. Jaouen and J. Mercier, 1977(b). Study of the sex pheromone of *Eupoecilia (Clysia) ambiguella* Hb. Lepidoptera Tortricidae, Cochylidae. *Ann. Zool. Entomol. Anim.*, **9**: 553-562.
- Sakin, V., 2023. Probability analysis of pheromone trapping for olive moths, *Prays oleae* Bern. (Lepidoptera: Yponomeutidae) and a regional decision-making algorithm for spraying for fruit generation. *Phytoparasitica*, **51**: 993-1006.

- Sanders, C. J., G. E. Daterman, R. F. Shepherd and H. Cerezke, 1974. Sex attractants for two species of western spruce budworm, *Choristoneura biennis* and *C. viridis* (Lepidoptera: Tortricidae). *Can. Entomol.*, **106**: 157-159.
- Sanders, C. J. and J. Weatherston, 1976. Sex pheromone of the eastern spruce budworm (Lepidoptera: Tortricidae): optimum blend of *trans*- and *cis*-tetradecenal. *Can. Entomol.*, **108**: 1285-1290.
- Sanders, C.J. 1984. Sex pheromone of the spruce budworm (Lepidoptera: Tortricidae): evidence for a missing component. *Can. Entomol.*, **116**: 93-100.
- Sanders, C. J., 1993. Non-target Lepidoptera captured in traps baited with spruce budworm pheromone in Ontario. *Can. Entomol.*, **125**: 1067-1076.
- Sans, A., M. Riba, M. Eizaguirre and C. Lopez, 1997. Electroantennogram, wind tunnel and field responses of male Mediterranean corn borer, *Sesamia nonagrioides*, to several blends of its sex pheromone components. *Entomol. Exp. Appl.*, **82**(2): 121-127.
- Santangelo, E. M., M. Coracini, P. Witzgall, A. G. Correa and C. R. Unelius, 2002. Identification, syntheses, and characterization of the geometric isomers of 9,11-hexadecadienal from female pheromone glands of the sugar cane borer *Diatraea saccharalis*. *J. Nat. Prod.*, **65**: 909-915.
- Sartwell, C., G. E. Daterman and L. L. Sower, 1985. A synthetic attractant for female moths of a biotype in the *Cydia pipperana* complex (Lepidoptera: Tortricidae). *Can. Entomol.*, **117**: 1151-1152.
- Sasaerila, Y., G. Gries, G. Khaskin, R. Gries and Hardi, 1997. Identification of sex pheromone components of nettle caterpillar, *Setothosea asigna*. *J. Chem. Ecol.*, **23**(9): 2187-2196.
- Sasaerila, Y., R. Gries, G. Gries, G. Khaskin, S. King and T. C. Boo, 2000(a). Decadienoates: sex pheromone components of nettle caterpillars *Darna trima* and *D. bradleyi*. *J. Chem. Ecol.*, **26**(8): 1969-1981.
- Sasaerila, Y., R. Gries, G. Gries, G. Khaskin and Hardi, 2000(b). Sex pheromone components of nettle caterpillar, *Setora nitens*. *J. Chem. Ecol.*, **26**(8): 1983-1990.
- Sato, R., K. Yaginuma and M. Kumakura, 1979. Sex attractant of the stone leek miner, *Acrolepia alliella* Semenov et Kuznesov: (Z)-11-hexadecenyl acetate, (Z)-11-hexadecenal and (Z)-11-hexadecenol. *Jap. J. Appl. Entomol. Zool.*, **23**: 115-117.

- Sato, R., N. Abe, P. Sonnet, H. Sugie and Y. Tamaki, 1985. Biological activity of (R)- and (S)-14-methyl-1-octadecene, as the chiral component of the sex pheromone of the peach leafminer moth, *Lyonetia clerkella* Linné (Lepidoptera: Lyonetiidae). *Appl. Entomol. Zool.*, **20**: 411-415.
- Sato, R., N. Abe, H. Sugie, M. Kato, K. Mori and Y. Tamaki, 1986. Biological activity of the chiral sex pheromone of the peach leafminer moth, *Lyonetia clerkella* Linné (Lepidoptera: Lyonetiidae). *Appl. Entomol. Zool.*, **21**: 478-480.
- Sato, Y., S. Takahashi, M. Sakai and T. Kodama, 1980. Attractiveness of the synthetic sex pheromone to the males of the armyworm, *Leucania separata* Walker and the lorey leafworm, *Leucania loreyi* Duponchel (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **15**: 334-340.
- Saveer, A. M., E. Hatano, A. Wada-Katsumata, R. L. Meagher and C. Schal, 2023. Nonanal, a new fall armyworm sex pheromone component, significantly increases the efficacy of pheromone lures. *Pest Manag. Sci.*, **79**: 2831-2839.
- Schaefer, P. W., G. Gries, R. Gries and D. Holden 1999. Pheromone components and diel periodicity of pheromonal communication in *Lymantria fumida*. *J. Chem. Ecol.*, **25**(10): 2305-2312.
- Schal, C., R. E. Charlton and R. T. Cardé, 1987. Temporal patterns of sex pheromone titers and release rates in *Holomelina lamae* (Lepidoptera: Arctiidae). *J. Chem. Ecol.*, **13**: 1115-1129.
- Schläger, S., F. Beran, A. T. Groot, C. Ulrichs, D. Veit, C. Paetz, B. R. M. Karumuru, R. Srinivasan, M. Schreiner and I. Mewi, 2015. Pheromone blend analysis and cross-attraction among populations of *Maruca vitrata* from Asia and West Africa. *J. Chem. Ecol.*, **41**: 1155-1162.
- Schlamp, K. K., R. Gries, G. Khaskin, K. Brown, E. Khaskin, G. J. R. Judd and G. Gries, 2005. Pheromone components from body scales of female *Anarsia lineatella* induce contacts by conspecific males. *J. Chem. Ecol.* **31**(12): 2897-2911.
- Schmidt, B. C., J. Roland and D. Wakarchuk, 2003. Evaluation of synthetic pheromones for monitoring forest tent caterpillar (Lepidoptera: Lasiocampidae) populations. *Environ. Entomol.*, **32**(1): 214-219.
- Schneider, D., S. Schulz, R. Kittmann and P. Kanagaratman, 1992. Pheromones and glandular structures of both sexes of the weed-defoliator moth *Pareuchaetes pseudoinsulata* Rego Barros (Lep., Arctiidae). *J. Appl. Entomol.*, **113**: 280-294.

- Schneider, D., S. Schulz, E. Priesner, J. Ziesmann and W. Francke 1998. Autodetection and chemistry of female and male pheromone in both sexes of the tiger moth *Panaxia quadripunctaria*. *J. Comp. Physiol. A*, **182**: 153-161.
- Schönherr, von J., 1972. Attraction of disparlure to the nun moth, *Lymantria monacha* L. *Z. Angew. Entomol.*, **71**: 260-263.
- Schwarz, M., J. A. Klun, E. R. Hart, B. A. Leonhardt and J. C. Weatherby, 1983(a). Female sex pheromone of the yellowheaded fireworm, *Acleris minuta* (Lepidoptera: Tortricidae). *Environ. Entomol.*, **12**: 1253-1256.
- Schwarz, M., J. A. Klun, B. A. Leonhardt and D. T. Johnson, 1983(b). (*E,Z*)-2,13-Octadecadien-1-ol acetate. A new pheromone structure for sesioid moths. *Tetrahedron Lett.*, **24**: 1007-1010.
- Scott, C. D., M. L. Winston, K. N. Slessor, G. G. S. King and G. G. Grant, 1984. The biology and pheromone-based monitoring of the driedfruit moth, *Vitula edmandsae serratilineella* (Lepidoptera: Pyralidae). *Can. Entomol.*, **116**: 1007-1013.
- Sekul, A. A. and A. N. Sparks, 1967. Sex pheromone of the fall armyworm moth: isolation, identification, and synthesis. *J. Econ. Entomol.*, **60**: 1270-1272.
- Seol, K. Y., H. Honda, K. Usui, T. Ando and Y. Matsumoto, 1987. 10,12,14-Hexadecatrienyl acetate: sex pheromone of the mulberry pyralid, *Glyphodes pyloalis* Walker (Lepidoptera: Pyralidae). *Agric. Biol. Chem.*, **51**: 2285-2287.
- Senda, S., T. Omata, Y. Ninomiya, M. Aburatani and T. Takeuchi, 1991. Attractant compositions for fall webworm. *European Patent Application*, EP 474,486.
- Sharp, J. L., J. R. McLaughlin, J. James, T. D. Eichlin and J. H. Tumlinson, 1978. Seasonal occurrence of male Sesiidae in north central Florida determined with pheromone trapping methods. *Florida Entomol.*, **61**: 245-250.
- Shaver, T. N., H. E. Brown, H. J. Williams, L. T. Woods and J. Worley, 1988. Components of female sex pheromone of *Eoreuma loftini* Dyar. *J. Chem. Ecol.*, **14**: 391-399.
- Shaver, T. N., D. E. Hendricks and J. D. Lopez, Jr., 1989. Influence of (*Z*)-11-hexadecen-1-ol on field performance of *Heliothis virescens* pheromone in a PVC dispenser as evidenced by trap capture. *J. Chem. Ecol.*, **15**: 1637-1644.

- Shaver, T. N., H. E. Brown and D. E. Hendricks, 1990. Development of pheromone lure for monitoring field populations of *Eoreuma loftini* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **16**: 2393-2399.
- Shearer, P. and H. Riedl, 1994. Comparison of pheromone trap bioassays for monitoring insecticide resistance of *Phyllonorycter elmaella* (Lepidoptera: Gracillariidae). *J. Econ. Entomol.*, **87**(6): 1450-1454.
- Shibasaki, H., M. Yamamoto, Q. Yan, H. Naka, T. Suzuki and T. Ando, 2013. Identification of the sex pheromone secreted by a nettle moth, *Monema flavescens*, using gas chromatography/Fourier transform infrared spectroscopy. *J. Chem. Ecol.*, **39**: 350-357.
- Shimuzu, N. and Y. Kuwahara, 2009. Female sex pheromone of a Japanese population of allium leafminer, *Acrolepiopsis sapporensis* (Lepidoptera: Acrolepiidae). *J. Pestic. Sci.*, **34**: 181-183.
- Shoji, A., Y. Adachi, H. Naka, T. Ando and Y. Arita, 2009. *Nokona regalis* (Lepidoptera, Sesiidae): a late collection record in autumn and a sex attractant. *Trans. Lipid. Soc. Jpn*, **60**: 193-195.
- Shorey, H. H., L. K. Gaston and J. S. Roberts, 1965. Sex pheromones of noctuid moths. VI. Absence of behavioral specificity for the female sex pheromones of *Trichoplusia ni* versus *Autographa californica*, and *Heliothis zea* versus *H. virescens* (Lepidoptera: Noctuidae). *Ann. Entomol. Soc. Am.*, **58**: 600-603.
- Shry, Y. L. and Y. S. Chow, 1985. Field attraction experiment of the synthetic sex pheromone of the smaller tea tortrix and oriental tea tortrix in Taiwan. *J. Agric. Assoc. China*, **132**: 108.
- Siderhurst, M. S., E. B. Jang, A. H. Hara and P. Conant, 2007. *n*-Butyl (*E*)-7,9-decadienoate: sex pheromone component of the nettle caterpillar, *Darna pallivitta*. *Entomol. Exp. Appl.*, **125** (1): 63-69.
- Silk, P. J., S. H. Tan, C. J. Wiesner, R. J. Ross and G. C. Lonergan, 1980. Sex pheromone chemistry of the eastern spruce budworm, *Choristoneura fumiferana*. *Environ. Entomol.*, **9**: 640-644.
- Silk, P. J., C. J. Wiesner, S. H. Tan, R. J. Ross and G. G. Grant, 1982. Sex pheromone chemistry of the western spruce budworm *Choristoneura occidentalis* Free. *J. Chem. Ecol.*, **8**: 351-362.
- Silk, P. J., L. P. S. Kuenen and G. C. Lonergan, 1985(a). Abiologically active analogue of the primary sex-pheromone components of spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **117**: 257-260.

- Silk, P. J., L. P. S. Kuenen, S. H. Tan, W. L. Roelofs, C. J. Sanders and A. R. Alford, 1985(b). Identification of sex pheromone components of jack pine budworm, *Choristoneura pinus pinus* Freeman. *J. Chem. Ecol.*, **11**: 159-167.
- Silk, P. J., E. W. Butterworth, L. P. S. Kuenen, C. J. Northcott and E. G. Kettela, 1988(a). Sex pheromone of purple striped shootworm, *Zeiraphera unfortunana* Powell. *J. Chem. Ecol.*, **14**: 1417-1425.
- Silk, P. J. and L. P. S. Kuenen, 1988(b). Sex pheromones and behavioral biology of the coniferophagous *Choristoneura*. *Ann. Rev. Entomol.*, **33**: 83-101.
- Silk, P. J., E. W. Butterworth, L. P. S. Kuenen, C. J. Northcott, E. Dunkelblum and E. G. Kettela, 1989. Identification of sex pheromone component of spruce budmoth *Zeiraphera canadensis*. *J. Chem. Ecol.*, **15**: 2435-2444.
- Silk, P. J., L. P. S. Kuenen, A. R. Alford, G. C. Lonergan and G. G. Grant, 1997. Sex pheromone components of the oak leaf shredder, *Croesia semipurpurana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **129**: 1001-1008.
- Silk, P. J., G. C. Lonergan, D. C. Allen and J. Spear-O'Mara, 2000. Potential sex pheromone components of the saddled prominent (Lepidoptera: Notodontidae). *Can. Entomol.*, **132**: 681-684.
- Silk, P. J., E. Eveleigh, L. Roscoe, K. Burgess, S. Weatherby, G. Leclair, P. Mayo, M. Brophy, 2017. Unsaturated cuticular hydrocarbons enhance responses to sex pheromone in spruce budworm, *Choristoneura fumiferana*. *J. Chem. Ecol.*, **43**: 753-762.
- Silva, E. C. A. D., P. J. Silk, P. Mayo, N. K. Hillier, D. Magee and G. C. Cutler, 2013. Identification of sex pheromone components of blueberry spanworm *Itame argillacearia* (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **39**: 1169-1181.
- Silva, M. R. da, A. M. P. Cortes, G. P. Svensson, C. Löfstedt, E. R. Limac and P. H. G. Zarbin, 2021. Identification of two additional behaviorally active gland constituents of female *Diatraea saccharalis* (Fabricius) (Lepidoptera: Crambidae). *J. Braz. Chem. Soc.*, **32**: 225-230.
- Simmons, R. B., W. E. Conner and R. B. Davidson, 1998. Identification of major components of the female pheromone glands of *Euchaetes egle* Drury (Arctiidae). *J. Lepid. Soc.*, **52**: 356-363.
- Sinclair, E. R. and P. Sinclair, 1980. Trapping adult macadamia nut borer, *Cryptophlebia ombrodelta* (Lower) (Lepidoptera:

- Tortricidae). *J. Aust. Entomol. Soc.*, **19**: 211-216.
- Sisay, B., A. Tamiru, S. Subramanian, C. W. Weldon, F. Khamis, K. K. Green, P. Anderson and B. Torto, 2024. Pheromonal variation and mating between two mitotypes of fall armyworm (*Spodoptera frugiperda*) in Africa. *Sci. Rep.*, **14**: 3848.
- Slessor, K. N., J. Raine, G. G. S. King, S. J. Clements and S. A. Allan, 1987. Sex pheromone of blackheaded fireworm, *Rhopobota naevana* (Lepidoptera: Tortricidae), a pest of cranberry. *J. Chem. Ecol.*, **13**: 1163-1170.
- Smith, R. G., G. E. Daterman, G. D. Daves, Jr., K. D. McMurtrey and W. L. Roelofs, 1974. Sex pheromone of the European pine shoot moth: chemical identification and field tests. *J. Insect Physiol.*, **20**: 661-668.
- Smith, R. G., G. E. Daterman and G. D. Daves, Jr., 1975. Douglas-fir tussock moth: sex pheromone identification and synthesis. *Science*, **188**: 63-64.
- Smith, L. M., R. G. Smith, T. M. Loehr and G. D. Daves, Jr., 1978. Douglas fir tussock moth pheromone: identification of a diene analogue of the principal attractant and synthesis of stereochemically defined 1,6-, 2,6-, and 3,6-heneicosadien-11-ones. *J. Org. Chem.*, **43**: 2361-2366.
- Smith, M. T., L. M. McDonough, S. Voerman, H. G. Davis, R. D. Eikenbary and C. L. Smithhisler, 1987. Hickory shuckworm *Cydia caryana*: electroantennogram and flight tunnel studies of potential sex pheromone components. *Entomol. Exp. Appl.*, **44**: 23-30.
- Snow, J. W., T. D. Eichlin and J. H. Tumlinson, 1985. Seasonal captures of clearwing moths (Sesiidae) in traps baited with various formulations of 3,13-octadecadienyl acetate and alcohol. *J. Agric. Entomol.*, **2**: 73-84.
- Snow, J. W., M. Schwarz and J. A. Klun, 1987. The attraction of the grape root borer, *Vitacea polistiformis* (Harris) (Lepidoptera: Sesiidae) to (*E,Z*)-2,13-octadecadienyl acetate and the effects of related isomers on attraction. *J. Entomol. Sci.*, **22**: 371-374.
- Snow, J. W., M. Schwarz and T. D. Eichlin, 1989. Captures of clearwing moths (Lepidoptera: Sesiidae) with various octadecadienyl acetates and alcohols in central Georgia during 1983-1985. *Environ. Entomol.*, **18**: 216-222.
- Snow, W. J., D. T. Johnson and J. R. Meyer, 1991. The seasonal occurrence of the grape root borer, (Lepidoptera: Sesiidae) in the eastern United States. *J. Entomol. Sci.*, **26**: 157-168.

- Soares, A. M. L., P. H. B. França, M. F. Triana, J. M. D. Santos, N. S. Dias-Pini, H. F. Goulart, J. X. Araújo-Júnior and A. E. G. Santana, 2020. Identification of $\Delta 6$ -unsaturated, monoenyl type I pheromone compounds from the cashew stem borer *Anthistarcha binocularis* (Lepidoptera: Gelechiidae). *Pest Manag. Sci.*, **76**: 1435-1442.
- Soderstrom, E. L., D. G. Brandl, J. Myerson, R. G. Buttery and B. E. Mackey, 1985. Sex pheromone for attracting western grapeleaf skeletonizer (Lepidoptera: Zygaenidae). *J. Econ. Entomol.*, **78**: 799-801.
- Soldán, T., and Spitzer, K. 1983. Some moths recorded at sex pheromone traps in Mitidja, Algeria (Lepidoptera: Tortricidae, Pyralidae, Noctuidae). *Acta Entomol. Bohemoslov.*, **80**: 395-398.
- Solomon, J. D., M. E. Dix and R. E. Doolittle, 1978. Attractiveness of the synthetic carpenterworm sex attractant increased by isomeric mixtures and prolonged by preservatives. *Environ. Entomol.*, **7**: 39-41.
- Solomon, J. D., F. L. Oliveria, J. H. Tumlinson and R. E. Doolittle, 1982. Occurrence of clearwing borers (Sesiidae) in west central Mississippi. *J. Georgia Entomol. Soc.*, **17**: 4-12.
- Song, Y. H., H. Y. Song, H. K. Kim, Y. D. Chang and P. C. Lippold, 1978. Studies on the synthetic pheromones of striped rice borer and tortricid insect pests. *Kor. J. Plant Prot.*, **17**: 41-47.
- Sower, L. L., G. E. Daterman, C. Sartwell and H. T. Cory, 1979. Attractants for the Western pine shoot borer, *Eucosma sonomana*, and *Rhyacionia zozana* determined by field screening. *Environ. Entomol.*, **8**: 265-267.
- Spitzer, K., M. Rejmánek, T. Soldán and J. Zeleny', 1983. Flight activity of some moths (Lepidoptera) monitored by sex pheromones in Vietnam: dry season. *Acta Entomol. Bohemoslov.*, **80**: 413-418.
- Sreng, I., B. Maume and B. Frérot, 1985. Analysis of the pheromonal secretion of virgin females of *Sesamia nonagrioides* (Lef.) (Lepidoptera, Noctuidae). *C. R. Acad. Sci., Ser. 3*, **301**: 439-442.
- Stan, G. and L. Pop, 1992. Attractiveness of pheromone variants with (Z) -11-hexadecenyl acetate females of some lepidopteran species. *Soc. Lepid. Rom. Bull. Inf.*, **10**: 10-15.
- Starratt, A. N., K. H. Dahm, N. Allen, J. G. Hildebrand, T. L. Payne and H. Röllner, 1979. Bombykal, a sex pheromone of the sphinx moth *Manduca sexta*. *Z. Naturforsch.*, **34 c**, 9-12.

- Steck, W. F., B. K. Bailey, E. W. Underhill and M. D. Chisholm, 1976. A sex attractant for the great dart, *Eurois occulta*: a mixture of (Z)-9-tetradecen-1-ol acetate and (Z)-11-hexadecen-1-ol acetate. *Environ. Entomol.*, **5**: 523-526.
- Steck, W., E. W. Underhill, M. D. Chisholm, B. K. Bailey, J. Loeffler and C. G. Devlin, 1977(a). Sex attractants for males of 12 moth species found in Western Canada. *Can. Entomol.*, **109**: 157-160.
- Steck, W., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1977(b). A sex attractant for male moths of the glassy cutworm *Crymodes devastator* (Brace): a mixture of Z-11-hexadecen-1-yl acetate, Z-11-hexadecenal and Z-7-dodecen-1-yl acetate. *Environ. Entomol.*, **6**: 270-273.
- Steck, W., B. K. Bailey, M. D. Chisholm and E. W. Underhill, 1978. A sex attractant for males of the cutworm *Euxoa pleuritica* (Lepidoptera: Noctuidae). *Can. Entomol.*, **110**: 775-777.
- Steck, W. F., D. L. Struble, C. E. Lilly, M. D. Chisholm, E. W. Underhill and G. E. Swailes, 1979(a). A sex attractant for males of the early cutworm, *Euxoa tristicula* (Lepidoptera: Noctuidae). *Can. Entomol.*, **111**: 337-341.
- Steck, W. F., E. W. Underhill, M. D. Chisholm and H. S. Gerber, 1979(b). Sex attractant for male alfalfa looper moths, *Autographa californica*. *Environ. Entomol.*, **8**: 373-375.
- Steck, W. F., M. D. Chisholm, B. K. Bailey and E. W. Underhill, 1979(c). Moth sex attractants found by systematic field testing of 3-component acetate-aldehyde candidate lures. *Can. Entomol.*, **111**: 1263-1269.
- Steck, W. F., E. W. Underhill, M. D. Chisholm and J. R. Byers, 1979(d). Sex attractants for *Agrotis venerabilis* and *Euxoa albipennis* based on (Z)-5-decenyl acetate and (Z)-7-dodecenyl acetate. *Environ. Entomol.*, **8**: 1126-1128.
- Steck, W., E. W. Underhill and M. D. Chisholm, 1980(a). Trace components in lepidopterous sex attractants. *Environ. Entomol.*, **9**: 583-585.
- Steck, W., M. D. Chisholm, E. W. Underhill and C. C. Peters, 1980(b). Optimized conditions for sex attractant trapping male redbacked cutworm moths, *Euxoa ochrogaster* (Guenée). *J. Chem. Ecol.*, **6**: 585-591.
- Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1980(c). Improved sex attractant blend for adult males of the glassy cutworm, *Crymodes devastator* (Lepidoptera: Noctuidae). *Can. Entomol.*,

112: 751-752.

- Steck, W., E. W. Underhill and M. D. Chisholm, 1982(a). Structure-activity relationships in sex attractants for North American noctuid moth. *J. Chem. Ecol.*, **8**: 731-754.
- Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1982(b). Trace co-attractants in synthetic sex lures for 22 noctuid moths. *Experientia*, **38**: 94-96.
- Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1982(c). (Z)-7-Tetradecenal, a seasonally dependent sex pheromone of the w-marked cutworm, *Spaelotis clandestina* (Harris) (Lepidoptera: Noctuidae). *Environ. Entomol.*, **11**: 1119-1122.
- Steck, W., E. W. Underhill, M. D. Chisholm and B. K. Bailey, 1982(d). (Z)-5-Tetradecenyl acetate, a trapping synergist for the major sex pheromone of the spotted cutworm, *Amathes c-nigrum*. *Entomol. Exp. Appl.*, **31**: 328-330.
- Steck, W. F., E. W. Underhill, B. K. Bailey and M. D. Chisholm, 1982(e). A 4-component sex attractant for male moths of the armyworm, *Pseudaletia unipuncta*. *Entomol. Exp. Appl.*, **32**: 302-304.
- Steck, W. F., E. W. Underhill and P. Palaniswamy, 1986. Sex pheromone of the cutworm *Feltia jaculifera* (Lepidoptera: Noctuidae). *Environ. Entomol.*, **15**: 40-43.
- Steinbauer, M. J., F. Östrand, T. E. Bellas, A. Nilsson, F. Andersson, E. Hedenström, M. J. Lacey and F. P. Schiestl, 2004. Identification, synthesis and activity of sex pheromone gland components of the autumn gum moth (Lepidoptera: Geometridae), a defoliator of *Eucalyptus*. *Chemoecology*, **14**: 217-223.
- Sternlicht, M., S. Goldenberg, B. F. Nesbitt, D. R. Hall and R. Lester, 1978. Field evaluation of the synthetic female sex pheromone of the citrus flower moth, *Prays citri* (Mill.) (Lepidoptera: Yponomeutidae), and related compounds. *Phytoparasitica*, **6**: 101-113.
- Stevens, R. E., C. Sartwell, T. W. Koerber, G. E. Daterman, L. L. Sower and J. A. Powell, 1980. Western *Rhyacionia* (Lepidoptera: Tortricidae, Olethreutinae) pine tip moths trapped using synthetic sex attractants. *Can. Entomol.*, **112**: 591-603.
- Stevens, R. E., C. Sartwell, T. W. Koerber, J. A. Powell, G. E. Daterman and L. L. Sower, 1985. Forest tortricids trapped using *Eucosma* and *Rhyacionia* synthetic sex attractants. *J. Lepidop. Soc.*, **39**: 26-32.

- Stockel, J. and P. Anglade, 1977. Influence de la concentration en isomères cis et trans dans la phéromone sexuelle d'*Ostrinia nubilalis* Hb. sur le comportement d'orientation de cette espèce et sur celui de *Pyrausta aurata* Sc. *C. R. Hebd. Seances Acad. Sci., Ser. D*, **285**: 61-64.
- Stockel, J., 1981. An isomer mixture *Z/E* of 11-tetradecenyl acetate, sex attractant for small ermine moths (Lepidoptera, Hyponomeutinae). *Agronomie*, **1**: 355-358.
- Strong, W. B., J. G. Millar, G. G. Grant, J. A. Moreira, J. M. Chong and C. Rudolph, 2008. Optimization of pheromone lure and trap design for monitoring the fir coneworm, *Dioryctria abietivorella*. *Entomol. Exp. Appl.*, **126**: 67-77.
- Struble, D. L. and G. E. Swailes, 1975(a). A sex attractant for the clover cutworm, *Scotogramma trifolii* (Rottenberg), a mixture of *Z*-11-hexa-decen-1-ol acetate and *Z*-11-hexadecen-1-ol. *Environ. Entomol.*, **4**: 632-636.
- Struble, D. L., M. Jacobson, N. Green and J. D. Warthen, 1975(b). Bertha armyworm (Lepidoptera: Noctuidae): detection of a sex pheromone and the stimulatory effect of some synthetic chemicals. *Can. Entomol.*, **107**: 355-359.
- Struble, D. L., G. E. Swailes, W. F. Steck, E. W. Underhill and M. D. Chisholm, 1976. A sex attractant for the adult males of variegated cutworm, *Peridroma saucia*. *Environ. Entomol.*, **5**: 988-990.
- Struble, D. L. and C. E. Lilly, 1977(a). An attractant for the beet webworm, *Loxostege sticticalis* (Lepidoptera: Pyralidae). *Can. Entomol.*, **109**: 261-266.
- Struble, D. L. and G. E. Swailes, 1977(b). A sex attractant for the adult males of the army cutworm, *Euxoa auxiliaris*: a mixture of *Z*-5-tetradecen-1-yl acetate and *E*-7-tetradecen-1-yl acetate. *Environ. Entomol.*, **6**: 719-724.
- Struble, D. L., G. E. Swailes and G. L. Ayre, 1977(c). A sex attractant for males of the darksided cutworm, *Euxoa messoria* (Lepidoptera: Noctuidae). *Can. Entomol.*, **109**: 975-980.
- Struble, D. L., G. E. Swailes, W. F. Steck, E. W. Underhill and M. D. Chisholm, 1977(d). A sex attractant for *Leucania commoides*: a mixture of *Z*-9-tetradecen-1-yl acetate, *Z*-11-hexadecen-1-yl acetate, and *Z*-11-hexadecen-1-ol. *Can. Entomol.*, **109**: 1393-1398.
- Struble, D. L. and G. E. Swailes, 1978(a). A sex attractant for adult

- males of the pale western cutworm, *Agrotis orthogonia* (Lepidoptera: Noctuidae). *Can. Entomol.*, **110**: 769-773.
- Struble, D. L. and G. E. Swailes, 1978(b). Sex attractant for adult males of the redbacked cutworm, *Euxoa ochrogaster*. *Environ. Entomol.*, **7**: 384-386.
- Struble, D. L., H. R. Buser, H. Arn and G. E. Swailes, 1980(a). Identification of sex pheromone components of redbacked cutworm, *Euxoa ochrogaster*, and modification of sex attractant blend for adult males. *J. Chem. Ecol.*, **6**: 573-584.
- Struble, D. L., H. Arn, H. R. Buser, E. Städler and J. Freuler, 1980(b). Identification of 4 sex pheromone components isolated from calling females of *Mamestra brassicae*. *Z. Naturforsch.*, **35 c**: 45-48.
- Struble, D. L., 1981(a). Modification of the attractant blend for adult males of the army cutworm, *Euxoa auxiliaris* (Grote), and the development of an alternate 3-component attractant blend for this species. *Environ. Entomol.*, **10**: 167-170.
- Struble, D. L., 1981(b). A four-component pheromone blend for optimum attraction of redbacked cutworm males, *Euxoa ochrogaster* (Guenée). *J. Chem. Ecol.*, **7**: 615-625.
- Struble, D. L., W. F. Steck, G. E. Swailes, M. D. Chisholm, E. W. Underhill and C. E. Lilly, 1982. Two two-component sex attractant blends for adult males of the striped cutworm, *Euxoa tessellata* (Lepidoptera: Noctuidae). *Can. Entomol.*, **114**: 359-362.
- Struble, D. L., 1983(a). Sex pheromone components of *Euxoa drewseni*: chemical identification, electrophysiological evaluation, and field attractancy tests. *J. Chem. Ecol.*, **9**: 335-346.
- Struble, D. L. and K. W. Richards, 1983(b). Identification of sex pheromone components of the female driedfruit moth, *Vitula edmandsae serratilineella*, and a blend for attraction of male moths. *J. Chem. Ecol.*, **9**: 785-801.
- Struble, D. L., G. L. Ayre and J. R. Byers, 1984. Minor sex-pheromone components of *Mamestra configurata* (Lepidoptera: Noctuidae) and improved blends for attraction of male moths. *Can. Entomol.*, **116**: 103-105.
- Struble D. L. and J. R. Byers, 1985. Identification of sex-pheromone components of the sibling species *Euxoa ridingsiana* (Grt.) and *Euxoa maimes* (Sm.) (Lepidoptera: Noctuidae), and blends for their specific attraction. *Can. Entomol.*, **117**: 495-504.
- Struble, D. L., J. R. Byers, D. G. R. McLeod and G. L. Ayre, 1987(a).

- Sex pheromone components of an Alberta population of European corn borer, *Ostrinia nubilalis* (Hbn.) (Lepidoptera: Pyralidae). *Can. Entomol.*, **119**: 291-299.
- Struble, D. L. and J. R. Byers, 1987(b). Identification of sex pheromone components of dark sided cutworm, *Euxoa messoria*, and modification of sex attractant blend for adult males. *J. Chem. Ecol.*, **13**: 1187-1199.
- Struble, D. L., G. L. Ayre and J. R. Byers, 1987(c). Sex attractant blends for strawberry cutworm, *Amphipoea interoceanica* (Smith), and a closely related species, *Amphipoea americana* (Speyer) (Lepidoptera: Noctuidae). *Can. Entomol.*, **119**: 301-303.
- Struble, D. L., J. R. Byers, R. F. Shepherd and T. G. Gray, 1989. Identification of sex pheromone components of the black army cutworm, *Actebia fennica* (Tauscher) (Lepidoptera: Noctuidae), and a sex attractant blend for adult males. *Can. Entomol.*, **121**: 557-563.
- Su, M.-W., Y.-L. Fang, W.-Q. Tao, G.-Z. Yan, W.-E. Ma and Z.-N. Zhang, 2008. Identification and field evaluation of the sex pheromone of an invasive pest, the fall webworm *Hyphantria cunea* in China. *Chinese Sci. Bull.*, **53**: 555-560.
- Subchev, M. A. and N. V. Kuznetsov, 1974. Preliminary identification of the sex pheromone of *Rhynchopacha* sp. (Lepidoptera: Gelechiidae) by field screening tests. *Dokl. Bolg. Akad. Nauk. (C. R. Acad. Bulg. Sci.)*, **27**: 993-994.
- Subchev, M. A., 1981. Z-11-Hexadecenyl acetate: sexual attractant for males of eight species of moths (Insecta, Lepidoptera). *Ekologiya*, **9**: 59-62.
- Subchev, M. A. and I. L. Stoilov, 1984. Co-attractants for *Mamestra brassicae* L. and *Mamestra suasa* Denis & Schiff. (Lepidoptera: Noctuidae). *Dokl. Bolg. Akad. Nauk. (C. R. Acad. Bulg. Sci.)*, **37**: 353-354.
- Subchev, M. A., J. A. Ganey, O. Vostrowsky and H. J. Bestmann, 1986(a). Screening and use of sex attractants in monitoring of Geometrid moths in Bulgaria. *Z. Naturforsch.*, **41 c**: 1082-1086.
- Subchev, M. A., J. A. Ganey and I. L. Stoilov, 1986(b). Noctuid male sex attractants - cis-11-hexadecenyl compounds alone and in mixtures. *Ekologiya*, **18**: 33.
- Subchev, M. A., L. S. Stanimirova and T. S. Milkova, 1987. The effect of compounds related to cis-11-hexadecenyl acetates on its attractiveness to the males of *Mamestra brassicae* L. (Lepidoptera:

- Noctuidae) and some other noctuid species. *Folia Biol.*, **35**: 143-150.
- Subchev, M. A., I. A. Krusteva, J. A. Ganev and T. S. Milkova, 1990. Some new lepidopteran sex attractants and attractant inhibitors in Bulgaria. *J. Appl. Entomol.*, **109**: 189-193.
- Subchev, M., 1992. Sex attractant for *Trachea atriplicis* L. (Lepidoptera: Noctuidae) males. *Dokl. Bulg. Akad. Nauk. (C. R. Acad. Bulg. Sci.)*, **45**: 121-123.
- Subchev, M. and L. Pop, 1994. Aldehyde sex attractants for *Agrochola* (Lepidoptera: Noctuidae) species. *Comptes-rendus de l'Académie bulgare des sciences*, **47**(4): 101-104.
- Subchev, M., A. Botar and S. Voerman, 1995. Sex attractant and attractant inhibitors for *Atethmia ambusta* F. (Lep., Noctuidae). *J. Appl. Entomol.*, **119**: 207-209.
- Subchev, M. A., M. Tóth, G. Szöcs, G. Stan and A. Botar, 1996. Evidence for geographical differences in pheromonal responses of male *Amathes c-nigrum* L. (Lep., Noctuidae). *J. Appl. Entomol.*, **120**: 615-617.
- Subchev, M., A. Harizanov, W. Francke, S. Franke, E. Plass, A. Reckziegel, F. Schröder, J. A. Pickett, L. J. Wadhams and C. M. Woodcock 1998. Sex pheromone of female vine bud moth, *Theresimima ampellophaga* comprises (2S)-butyl (7Z)-tetradecenoate. *J. Chem. Ecol.*, **24**(7): 1141-1151. [Erratum: (2S)-butyl → (2R)-butyl, *J. Chem. Ecol.*, **25**: 1203]
- Subchev, M. A., M. Toth, D. Wu and Z. Imrei, 1999. (Z)-8-Tridecenyl acetate - A sex attractant for three microlepidopteran species (Lepidoptera: Cosmopterigidae, Symmocidae). *Dokl. Bulg. Akad. Nauk. (C. R. Acad. Bulg. Sci.)*, **52**(7-8): 113-114.
- Subchev, M., T. Toshova, L. Stanimirova, G. Stan, G. Embacher, W. Francke, A. Reckziegel, J. T. Ferreira and E. Priesner, 2000(a). 1-Methylethyl octanoate, a new lepidopteran sex pheromone from the bagworm, *Megalophanes viciella*. *J. Chem. Ecol.*, **26**(2): 487-495.
- Subchev, M., M. Toth, D. Wu, L. Stanimirova, T. Toshova and Z. Karpati, 2000(b). Sex attractant for *Diloba caeruleocephala* (L.), (Lep., Dilobidae) (Z)-8-tridecenyl acetate. *J. Appl. Entomol.*, **124**: 197-199.
- Subchev, M., A. Mircheva, J. Pickett, L. Wadhams, C. Woodcock, A. D. Santos, S. Franke and W. Francke, 2003. Sex pheromone of the leaf-miner *Phyllonorycter platani*: (Z10)-tetradecenyl acetate.

J. Chem. Ecol., **29**(10): 2391-2396.

- Subchev, M., T. Toshova, M. Tóth, E. Voigt, J. Mikulás and W. Francke, **2004**. Catches of vine bud moth *Theresimima ampellophaga* (Lepidoptera, Zygaenidae: Procridinae) males in pheromone traps: effect of purity and age of baits, design, colour and height of the traps, and daily sexual activity of males. *J. Appl. Entomol.*, **128**: 44-50.
- Subchev, M., T. Toshova, C. Koshio, S. Franke, A. Tröger, R. Twele, W. Francke, J. A. Pickett, L. J. Wadhams and C. M. Woodcock, **2009**. Identification and biological activity of sex pheromone components from females of the plum moth *Illiberis rotundata* Jordan (Lepidoptera: Zygaenidae: Procridinae). *Chemoecol.*, **19**: 47-54.
- Subchev, M. A., K. A. Efetov, T. B. Toshova, E. V. Parshkova, W. Tóth and W. Francke, **2010**. New sex attractants for species of the zygaenid subfamily Procridinae (Lepidoptera: Zygaenidae). *Entomol. Gen.*, **32**: 243-250.
- Subchev, M., C. Koshio, T. Toshova, K. A. Efetov and W. Francke, **2013**. (2R)-butyl (7Z)-dodecenoate, a main sex pheromone component of *Illiberis* (*Primilliberis*) *pruni* Dyar (Lepidoptera: Zygaenidae: Procridinae)? *Acta Zool. Bulg.*, **65**: 391-396.
- Subchev, M., **2014**. Sex pheromone communication in the family Zygaenidae (Insecta: Lepidoptera): a review. *Acta Zool. Bulg.*, **66**: 147-157.
- Suckling, D. M., A. R. Gibb, H. Gurlay, P. Conant, C. Hirayama, R. Leen and G. Szocs, **2000**. Sex attractant for the gorse biocontrol agent *Agonopterix ulicetella* (Oecophoridae). *Proc. N. Z. Plant Prot. Conf.*, **53**: 66-70. [*Chem. Abst.*, **135**: 57326]
- Suckling, D. M., A. R. Gibb, J. M. Daly, D. J. Rogers and G. P. Walker, **2002**. Improving the pheromone lure for diamondback moth. *NZ Plant Prot.*, **55**: 182-187.
- Suckling, D. M., A. R. Gibb, G. M. Burnip, C. Snelling, J. De Rutter, G. Langford and A. M. El-Sayed, **2005**(a). Optimization of pheromone lure and trap characteristics for currant clearwing, *Synanthedon tipuliformis*. *J. Chem. Ecol.*, **31**(2): 393-406.
- Suckling, D. M., A. R. Gibb, P. R. Dentener, D. S. Seldon, G. K. Clare, L. Jamieson, D. Baird, D. J. Kriticos and A. M. El-Sayed, **2005**(b). *Uraba lugens* (Lepidoptera: Nolidae) in New Zealand: pheromone trapping for delimitation and phenology. *J. Econ. Entomol.*, **98**: 1187-1192.
- Sugie, H., K. Yaginuma, Y. Tamaki, **1977**. Sex pheromone of the Asiatic

- leafroller, *Archippus breviplicanus* Walsingham (Lepidoptera: Tortricidae): isolation and identification. *Appl. Entomol. Zool.*, **12**: 69-74.
- Sugie, H., Y. Tamaki, R. Sato and M. Kumakura, 1984(a). Sex pheromone of the peach leafminer moth, *Lyonetia clerkella* Linné: isolation and identification. *Appl. Entomol. Zool.*, **19**: 323-330.
- Sugie, H., Y. Tamaki, S. Shirasaki and T. Kitamura, 1984(b). Further studies on the sex pheromone of the summer fruit tortrix moth, *Adoxophyes orana fasciata*. *Jap. J. Appl. Entomol. Zool.*, **28**: 156-160.
- Sugie, H., Y. Tamaki, K. Kawasaki, M. Wakou, T. Oku, C. Hirano and M. Horiike, 1986. Sex pheromone of the apple leafminer moth, *Phyllonorycter ringoniella* (Matsumura) (Lepidoptera: Gracillariidae): activity of geometrical isomers of tetradecadienyl acetates. *Appl. Entomol. Zool.*, **21**: 578-581.
- Sugie, H., K. Kawasaki, H. Higuchi, C. Kitamura and Y. Tamaki, 1990. Identification of the sex pheromone of the three-spotted plusia, *Acanthopplusia agnata* Staudinger (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **25**: 467-473.
- Sugie, H., K. Kawasaki, S. Nakagaki and N. Iwata, 1991(a). Identification of sex pheromone of the semi-looper, *Autographa nigrisigna* Walker (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **26**: 71-76.
- Sugie, H., S. Tatsuki, S. Nakagaki, C. B. J. Rao and A. Yamamoto, 1991(b). Identification of the sex pheromone of the oriental tobacco budworm, *Heliothis assulta* (Guenée) (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **26**: 151-153.
- Sugie, H., H. Noguchi, K. Shimizu, T. Kawana, H. Fukuda, A. Yamamoto, T. Fukumoto and T. Ihara, 1993. Attractants containing sex pheromones for *Mamestra brassicae*. *Jpn. Kokai Tokkyo Koho* JP 05,255,009. [*Chem. Abst.* vol.120: No.7 p71571u]
- Sugie, H., H. Noguchi, K. Shimizu, T. Kawana, H. Fukuda, A. Yamamoto, T. Fukumoto and T. Ihara, 1994. Insect attractants containing Z-11-hexadecenal, Z-9-hexadecenal, and Z-11-hexadecenol for *Pediasia teterrellus*. *Jpn. Kokai Tokkyo Koho* JP 06 56,608. [*Chem. Abst.* vol.121: No.3 p29291y]
- Sugie, H., J. Yase, K. Fujita and Y. Shirai, 2003. A sex attractant of the cabbage webworm, *Hellula undalis* Fabricius (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **38**(1): 45-48.
- Svatoš, A., A. B. Attygalle, G. N. Jham, R. T. S. Frighetto, E. F.

- Vilela, D. Saman and J. Meinwald, 1996. Sex pheromone of tomato pest *Scrobipalpaloides absoluta* (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **22**: 787-800
- Svatoš, A., B. Kalinová, M. Oskovee, J. Kindl, O. Hovorka and I. Hrdý, 1999. Identification of a new Lepidopteran sex pheromone in picogram quantities using an antennal biodeceptor: (8E,10Z)-tetradeca-8,10-dienal from *Cameraria ohridella*. *Tetrahedron Lett.*, **40**: 7011-7014.
- Svatoš, A., B. Kalinová, J. Kindl, J. Kuldová, O. Hovorka, R. R. D. Nascimento and N. J. Oldham, 2001. Chemical characterization and synthesis of the major component of the sex pheromone of the sugarcane borer *Diatraea saccharalis*. *Collect. Czech. Chem. Comm.*, **66**: 1682-1690.
- Svensson, M. G. E., M. Bengtsson and J. Löfqvist, 1997. Individual variation and repeatability of sex pheromone emission of female turnip moths *Agrotis segetum*. *J. Chem. Ecol.* **23**: 1833-1850.
- Svensson, G. P., V. Tönnberg and L. Sigsgaard, 2019. Identification and field evaluation of (E)-11,13-tetradecadienal as sex pheromone of the strawberry tortrix (*Acleris comariana*). *J. Appl. Entomol.*, **143**: 535-541.
- Svensson, G. P., O. Anderbrant, E. Öberg, E. V. Jirle, S. Hellqvist and C. Löfstedt, 2023. Identification of (E)- and (Z)-11-tetradecenyl acetate as sex pheromone components of the currant pest *Euphyponomeutoides albithoracellus*. *J. Appl. Entomol.*, **147**: 313-319.
- Svensson, G. P., H. Huynh, A.-K. Isaksson, L. B. L. Myhre, C. Löfstedt, S. Mogan, E. Öberg, M. Rantanen, N. Trandem and O. Anderbrant, 2025. Geographic distribution, flight phenology and infestation level of the lepidopteran pests *Euphyponomeutoides albithoracellus*, *Lampronia capitella* and *Synanthedon tipuliformis* on black currants in Northern Europe. *J. Appl. Entomol.*, **149**: 1196-1206.
- Sweeney, J. D., J. A. McLean and L. M. Friskie, 1990. Roles of minor components in pheromone-mediated behavior of western spruce budworm moths. *J. Chem. Ecol.*, **16**: 1517-1530.
- Sziráki, G., 1978(a). Examinations on tortricid moths trapped by synthetic attractants (Lepidoptera). *Folia Entomol. Hung.*, **31**: 259-264.
- Sziráki, G., 1978(b). Specificity of sexual attraction traps for signalization of Oriental fruit moth (*Grapholita molesta* Busck). *Acta Phytopathol. Acad. Sci. Hung.*, **13**: 205-212.

- Sziráki, G., 1980. Notes on Coleophora and Cnephasia species trapped by synthetic attractants (Lepidoptera: Coleophoridae and Tortricidae). *Folia Entomol. Hung.*, **33**: 161-166.
- Szöcs, G., M. Tóth and L. Novák, 1981. Sex attractants for eight lepidopterous species. *Z. Angew. Entomol.*, **91**: 272-280.
- Szöcs, G., M. Tóth and L. Novák, 1983. Sex attractants and inhibitors for lepidopterous species found by field screening of olefinic compounds in Hungary. *Z. Angew. Entomol.*, **96**: 56-67.
- Szöcs, G., M. Tóth, H. J. Bestmann and O. Vostrowsky, 1984. A two-component sex attractant for males of the geometrid moth *Alsophila quadripunctata*. *Entomol. Exp. Appl.*, **36**: 287-291.
- Szöcs, G., M. Schwarz, G. Sziráki, M. Toth, J. A. Klun and B. A. Leonhardt, 1985. Sex pheromone of the female currant borer, *Synanthedon tipuliformis*: identification and field evaluation. *Entomol. Exp. Appl.*, **39**: 131-133.
- Szöcs, G., M. Tóth, H. J. Bestmann, O. Vostrowsky, R. R. Heath and J. H. Tumlinson, 1987. Polyenic hydrocarbons as sex attractants for geometrids and amatids (Lepidoptera) found by field screening in Hungary. *Z. Naturforsch.*, **42 c**: 165-168.
- Szöcs, G., M. Tóth, Gy. Sziráki and M. Schwarz, 1989. 2,13- and 3,13-Octadecadienyl compounds composing sex attractants for tineid and sesiid moths (Lepidoptera). *Biochem. System. Ecol.*, **17**: 417-422.
- Szöcs, G., L. A. Miller, W. Thomas, R. A. Vickers, G. H. L. Rothschild, M. Schwarz and M. Tóth, 1990. Compounds modifying male responsiveness to main female sex pheromone component of the currant borer, *Synanthedon tipuliformis* Clerk (Lepidoptera: Sesiidae) under field conditions. *J. Chem. Ecol.*, **16**: 1289-1305.
- Szöcs, G., V. Bu'da, P. Charmillot, P. Esbjerg, B. Freier, R. Gottwald, B. Kovalev, S. Maini, M. G. Solomon, O. Sorum, M. Subchev, M. Tóth and M. Van de Veire, 1991. Field tests of (E,Z)-3,13-octadecadien-1-olacetate: a sex attractant synergist for male currant borer, *Synanthedon tipuliformis*. *Entomol. Exp. Appl.*, **60**: 283-288.
- Szöcs, G., A. Raina, M. Tóth and B. A. Leonhardt, 1993(a). Sex pheromone components of *Heliothis maritima*: chemical identification, flight tunnel and field tests. *Entomol. Exp. Appl.*, **66**: 247-253.
- Szöcs, G., M. Tóth, W. Francke, F. Schmidt, P. Philipp, W. A. König, K. Mori, B. S. Hansson and C. Löfstedt, 1993(b). Species

- discrimination in five species of winter-flying geometrids (Lepidoptera) based on chirality of semiochemicals and flight season. *J. Chem. Ecol.*, **19**: 2721-2735.
- Szöcs, G., M. Tóth, Z. Kárpáti, J. Zhu, C. Löfstedt, E. Plass and W. Francke, **2004**. Identification of polyenic hydrocarbons from the northern winter moth, *Operophtera fagata*, and development of a species specific lure for pheromone traps. *Chemoecology*, **14**(1): 53-58.
- Szöcs, G., M. Tóth and K. Mori, **2005**. Absolute configuration of the major sex pheromone component of the satin moth, *Leucoma salicis*, verified by field trapping test in Hungary. *Chemoecology*, **15**(2): 127-128.
- Tabata, J., T. Takanashi and Y. Ishikawa, **2003**. Pheromone analysis of wild female moths with a PBAN C-terminal peptide injection for an estimation of assortative mating in adzuki bean borer, *Ostrinia scapulalis*. *J. Chem. Ecol.*, **29**(12): 2749-2759.
- Tabata, J., S. Hoshizaki, S. Tatsuki and Y. Ishikawa, **2006**. Heritable pheromone blend variation in a local population of the butterbur borer moth *Ostrinia zaguliaevi* (Lepidoptera: Crambidae). *Chemoecology* **16**: 123-128.
- Tabata, J., T. Yokosuka, M. Hattori, M. Ohashi, H. Noguchi and H. Sugie, **2008**(a). Sex attractant pheromone of the limabean pod borer, *Etiella zinckenella* (Treitschke) (Lepidoptera: Pyralidae), in Japan. *Appl. Entomol. Zool.*, **43**: 351-358.
- Tabata, J., Y. Huang, S. Ohno, Y. Yoshiyasu, H. Sugie, S. Tatsuki and Y. Ishikawa, **2008**(b). Sex pheromone of *Ostrinia* sp. newly found on the leopard plant *Farfugium japonicum*. *J. Appl. Entomol.*, **132**: 566-574.
- Tabata, J., M. Minamishima, H. Sugie, T. Fukumoto, F. Mochizuki and Y. Yoshiyasu, **2009**. Sex pheromone components of the pear fruit moth, *Acrobasis pyrivorella* (Matsumura). *J. Chem. Ecol.*, **35**: 243-249.
- Tabata, J. and S. Kubota, **2017**. Sex pheromone of the iris gelechiid moth *Monochroa divisella*. *Entomol. Exp. Appl.*, **164**: 113-119.
- Tabata, J., R. Nakano, H. Yasui, K. Nakamura, K. Takehara, H. Matsuda, Y. Ikenoue, Y. Kusuhata, K. Kinjo, K. Nakama, Y. Tsuha and Y. Ishikawa, **2023**. Sex pheromone of the fall armyworm, *Spodoptera frugiperda*: identification of a trace component that enhances attractiveness and specificity. *Entomol. Exp. Appl.*, **171**: 535-541.

- Taft, W. H., 2005. *Synanthedon bolteri* (Sesiidae) in Michigan. *J. Lepid. Soc.*, **59**: 235-236.
- Tagestad, A. D., 1975. Spruce budworm detection in North Dakota shelterbelts and nurseries with a synthetic sex attractant. *North Dakota Farm Res.*, **33**: 17-19.
- Taguri, T., K. Yaginuma, M. Yamamoto, T. Fujii and T. Ando, 2014. Enantiospecific synthesis and field evaluation of four stereoisomers of 10,14-dimethyloctadec-1-ene, a sex pheromone component secreted by female moths of the apple leafminer. *Biosci. Biotechnol. Biochem.*, **78**: 761-765.
- Takács, S., G. Gries and R. Gries, 2001. Communication ecology of webbing clothes moth: 4. Identification of male- and female-produced pheromones. *Chemoecology*, **11**: 153-159.
- Takahashi, S., M. Kawaradani, Y. Sato and M. Sakai, 1979. Sex pheromone components of *Leucania separata* Walker and *Leucania loreyi* Duponchel. *Jap. J. Appl. Entomol. Zool.*, **23**: 78-81.
- Takahashi, S., M. Kawaradani, Y. Sato and M. Sakai, 1980. The components of the female sex pheromone of the lorey leafworm, *Leucania loreyi* Duponchel. *Appl. Entomol. Zool.*, **15**: 499-501.
- Takahashi, S., Y. Sato and T. Kodama, 1983. Attractiveness of synthetic sex pheromone to males of the lorey leafworm, *Acantholeucania loreyi* (Duponchel). *Appl. Entomol. Zool.*, **18**: 435-437.
- Takanashi, T., S. Ohno, Y. Huang, S. Tatsuki, H. Honda and Y. Ishikawa 2000. A sex pheromone component novel to *Ostrinia* identified from *Ostrinia latipennis* (Lepidoptera: Crambidae). *Chemoecology*, **10**: 143-147.
- Tamaki, Y., H. Noguchi, T. Yushima, C. Hirano, K. Honma and H. Sugawara, 1971(a). Sex pheromone of the summerfruit tortrix: isolation and identification. *Kontyu*, **39**: 338-340.
- Tamaki, Y., H. Noguchi, T. Yushima and C. Hirano, 1971(b). Two sex pheromones of the smaller tea tortrix: isolation, identification, and synthesis. *Appl. Entomol. Zool.*, **6**: 139-141.
- Tamaki, Y., H. Noguchi and T. Yushima, 1973. Sex pheromone of *Spodoptera litura* (F.) (Lepidoptera: Noctuidae): isolation, identification, and synthesis. *Appl. Entomol. Zool.*, **8**: 200-203.
- Tamaki, Y. and T. Yushima, 1974(a). Sex pheromone of the cotton leafworm, *Spodoptera littoralis*. *J. Insect Physiol.*, **20**: 1005-1014.
- Tamaki, Y. and T. Yushima, 1974(b). Biological activity of the

- synthesized sex pheromone and its geometrical isomers of *Spodoptera litura* (F.) (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **9**: 73-79.
- Tamaki, Y., K. Honma and K. Kawasaki, 1977(a). Sex pheromone of the peach fruit moth, *Carposina niponensis* Walsingham (Lepidoptera: Carposinidae): isolation, identification and synthesis. *Appl. Entomol. Zool.*, **12**: 60-68.
- Tamaki, Y., K. Kawasaki, H. Yamada, T. Koshihara, N. Osaki, T. Ando, S. Yoshida and H. Kakinohana, 1977(b). (*Z*)-11-Hexadecenal and (*Z*)-11-hexadecenyl acetates: sex-pheromone components of the diamondback moth (Lepidoptera: Plutellidae). *Appl. Entomol. Zool.*, **12**: 208-210.
- Tamaki, Y., T. Yushima, M. Oda, K. Kida, K. Kitamura, S. Yabuki and J.H. Tumlinson, 1977(c). Attractiveness of 3,13-octadecadienyl acetates for males of clearwing moths. *Jap. J. Appl. Entomol. Zool.*, **21**: 106-107.
- Tamaki, Y., H. Noguchi, H. Sugie, R. Sato and A. Kariya, 1979. Minor components of the female sex-attractant pheromone of the smaller tea tortrix moth (Lepidoptera: Tortricidae): isolation and identification. *Appl. Entomol. Zool.*, **14**: 101-113.
- Tamaki, Y., H. Noguchi, H. Sugie, A. Kariya, S. Arai, M. Ohba, T. Terada, T. Suguro and K. Mori, 1980. Four-component synthetic sex pheromone of the smaller tea tortrix moth: field evaluation of its potency as an attractant for male moth. *Jap. J. Appl. Entomol. Zool.*, **24**: 221-228.
- Tamaki, Y., H. Sugie, M. Osakabe and P. Sonnet, 1983. Biological activities of *R*- and *S*-10-methyldodecyl acetates, the chiral component of the sex pheromone of the smaller tea tortrix moth (*Adoxophyes* sp., Lepidoptera: Tortricidae). *Appl. Entomol. Zool.*, **18**: 292-294.
- Tamaki, Y., H. Sugie, Y. Ando, A. Yamashita, S. Ooya, H. Kamiwada, H. Suzuki and T. Fukumoto, 1996. Insect attractants for *Aedia leucomelas*. *Jpn. Kokai Tokkyo Koho*, JP 08,295 602 [96,295,602].
- Tamhankar, A. J., T. P. Rajendran, M. W. Bhamburkar and M. R. Harwalkar, 1993. Optimum pheromone blend for pink bollworm males: evaluation with two experimental designs. *Int. J. Pest Manage.*, **39**: 111-112.
- Tan, Z.-X., Y.-W. Wu, G.-Q. Lin, B.-C. Wu, H.-Q. Liu, X.-Y. Xu, W.-S. Zhou, G.-Q. Pu and M.-S. Zhang, 1984. Study on identification and synthesis of insect pheromone XVII. The sex pheromone of

- Euproctis similis xanthocampa*. *Huaxue Xuebao* (*Acta Chimica Sinica*), **42**: 1178-1182.
- Tan, Z.-X., G.-Q. Lin, H.-Q. Liu, G.-Q. Pu, H.-M. Du, J.-P. Mao, L.-Z. Meng and C.-H. Wu, **1992**. Identification and synthesis of insect pheromone XXX. Sex pheromone of mulberry clearwing moth *Paradoxecia pieli* Lieu. *J. Chem. Ecol.*, **18**: 419-425.
- Tan, Z.-X., R. Gries, G. Gries, G.-Q. Lin, G.-Q. Pu, K. N. Slessor and J.-X. Li, **1996**. Sexpheromone components of mulberry looper, *Hemerophila atrilineata* Butler (Lepidoptera: Geometridae). *J. Chem. Ecol.*, **22**: 2263-2271.
- Tanaka, A., K. Yasuda and S. Wakamura **2001**. Sex pheromone of *Acrolepiopsis nagaimo* Yasuda (Lepidoptera: Acrolepiidae), a predominant leafminer on Chinese yam in Tottori Prefecture: field evaluation and activity of the third component. *Appl. Entomol. Zool.*, **36**(2): 203-207.
- Tanaka, M., K. Abe, T. Ando and L. V. Vang, **2007**. Identification of the sex pheromone secreted by females of *Grapholita inopinata* Heinrich (Lepidoptera: Tortricidae). *Res. Bull. Plant Prot. Japan*, **43**: 17-22.
- Taschenberg, E. F., R. T. Cardé, A. Hill, J. P. Tette and W. L. Roelofs, **1974**. Sex pheromone trapping of the grape berry moth. *Environ. Entomol.*, **3**: 192-194.
- Tatsuki, S., M. Kurihara, K. Usui, Y. Ohguchi, K. Uchiumi and J. Fukami, **1983**. Sex pheromone of the rice stem borer, *Chilo suppressalis* (Walker) (Lepidoptera: Pyralidae): the third component, Z-9-hexadecenal. *Appl. Entomol. Zool.*, **18**: 443-446.
- Tatsuki, S., H. Sugie, K. Usui, J. Fukami, M. H. Sumartaputra and A. N. Kuswadi, **1985**. Identification of possible sex pheromone of the yellow stem borer moth, *Scirpophaga incertulas* (Walker) (Lepidoptera: Pyralidae). *Appl. Entomol. Zool.*, **20**: 357-359.
- Teal, P. E. A., R. R. Heath, J. H. Tumlinson and J. R. McLaughlin, **1981**. Identification of a sex pheromone of *Heliothis subflexa* (Gn.) (Lepidoptera: Noctuidae) and field trapping studies using different blends of components. *J. Chem. Ecol.*, **7**: 1011-1022.
- Teal, P. E. A., J. R. McLaughlin, J. H. Tumlinson and R. R. Rush, **1982**. A sex attractant for *Platysenta videns* (Gn.) similar to the sex pheromone of *Heliothis virescens* (F.). *Florida Entomol.*, **65**: 583-584.
- Teal, P. E. A., R. J. West and J. E. Laing, **1983**. Identification of a blend of sex pheromone components of the potato stem borer

- (Lepidoptera: Noctuidae) for monitoring adults. *Proc. Entomol. Soc. Ont.*, **114**: 15-19.
- Teal, P. E. A., J. H. Tumlinson, J. R. McLaughlin, R. Heath and R. A. Rush, 1984. (Z)-11-Hexadecen-1-ol: a behavioral modifying chemical present in the pheromone gland of female *Heliothis zea* (Lepidoptera: Noctuidae). *Can. Entomol.*, **116**: 777-779.
- Teal, P. E. A., E. R. Mitchell, J. H. Tumlinson, R. R. heath, and H. Sugie, 1985. Identification of volatile sex pheromone components released by the southern armyworm, *Spodoptera eridania* (Cramer). *J. Chem. Ecol.*, **11**: 717-725.
- Teal, P. E. A., J. H. Tumlinson and R. R. Heath, 1986. Chemical and behavioral analyses of volatile sex pheromone components released by calling *Heliothis virescens* (F.) females (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **12**: 107-126.
- Teal, P. E. A., R. R. Heath, B. D. Dueben, J. A. Coffelt and K. W. Vick, 1995. Production and release of (Z,E)-9,12-tetradecadienal by sex pheromone glands of females of *Plodia interpunctella* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **21**: 787-799.
- Teasdale, C., G. J. R. Judd, R. Gries and G. Gries, 2013. Evaluation of synthetic sex pheromone for monitoring and management of raspberry crown borer *Pennisetia marginata* (Lepidoptera: Sesiidae). *Agric. Forest Entomol.*, **15**: 285-293.
- Teixeira, A. L., J. S. Oliveira, E. F. Vilela, E. R. Lima and H. O. Reis, 1989. Sex pheromone components of *Spodoptera latifascia* Walker (Lepidoptera: Noctuidae). *Revista Ceres*, **36**: 373-381.
- Terytze, K., H. Adam and B. Kovaljev, 1987. Use of pheromone traps for monitoring harmful Lepidoptera species in cabbage. *Arch Phytopathol. Pflanzenschutz, Berlin*, **6**: 465-473.
- Thomson, D. R., N. P. D. Angerilli, C. Vincent and A. P. Gaunce, 1991. Evidence for regional differences in the response of obliquebanded leafroller (Lepidoptera: Tortricidae) to sex pheromone blends. *Environ. Entomol.*, **20**: 935-938.
- Tiberi, R. and A. Niccoli, 1984. Observations over several years on the use of traps with the sex pheromone of *Thaumetopoea pityocampa* (Den. et Schiff.) (Lepidoptera, Thaumetopoeidae). *Redia*, **67**: 129-144.
- Tonini, C., G. Cassani, P. Piccardi, S. Maini, P. L. Castellari and E. Pasqualini, 1982. Sex pheromone components of the leafroller moth *Pandemis cerasana*. *J. Insect Physiol.*, **28**: 443-446.

- Tonini, C., G. Cassani, P. Massardo, G. Guglielmetti and P. L. Castellari, 1986. Study of female sex pheromone of leopard moth, *Zeuzera pyrina* L. Isolation and identification of three components. *J. Chem. Ecol.*, **12**: 1545-1558.
- Toshova, T., M. Subchev, E. Plass and W. Francke, 2003. *Scoliopteryx libatrix* (L.) (Lep., Noctidae) male reaction to the synthetic main sex pheromone component and its isomers - wind tunnel and field investigations. *J. Appl. Entomol.*, **127**: 195-199.
- Toshova, T. B., B. Zlatkov, M. Subchev and M. Tóth, 2017. Monitoring the seasonal flight activity of three tortricid pests in Bulgaria with a single sex pheromone-baited trap. *Acta Zool. Bulg.*, **69**: 283-292.
- Tóth, M., G. Szöcs, L. Novák and Cs. Szántay, 1979. Post season field screening of lepidopterous sex attractant candidates in Hungary. *Acta Phytopathol. Acad. Sci. Hung.*, **14**: 195-199.
- Tóth, M., J. Jakab and L. Novák, 1980. Identification of two components from the sex pheromone system of the white-line dart moth, *Scotia segetum* (Schiff.) (Lep., Noctuidae). *Z. Angew. Entomol.*, **90**: 505-510.
- Tóth, M., G. Szöcs, B. Majoros, T. E. Bellas and L. Novák, 1983. Experiments with a two-component sex attractant of the silver Y moth (*Autographa gamma* L), and some evidence for the presence of both components in natural female sex pheromone. *J. Chem. Ecol.*, **9**: 1317-1325.
- Tóth, M., T. E. Bellas and G. H. L. Rothschild, 1984. Role of pheromone components in evoking behavioral responses from male potato tuberworm moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae). *J. Chem. Ecol.*, **10**: 271-280.
- Tóth, M., P. M. Guerin, H.-R. Buser, H. Müller, G. Szöcs, G. Sziráki and H. Arn, 1985. Z-11-Tetradecenyl acetate: sex attractant of *Agapeta zoegana* (Lepidoptera: Tortricidae), a potential species for the biological control of knapweed. *Can. Entomol.*, **117**: 1163-1165.
- Tóth, M., G. Sziráki and G. Szöcs, 1986(a). Sex attractants for *Pexicopia malvella* and *Platyedra subcinerea*, pests of malvaceous ornamentals and cotton. *Entomol. Exp. Appl.*, **40**: 147-150.
- Tóth, M., G. Szöcs, C. Löfstedt, B. S. Hansson and M. Subchev, 1986(b). Sex pheromone components of *Mamestra suasa*: chemical analysis, electrophysiological activity, wind tunnel activity and field tests in two European countries. *Entomol. Exp. Appl.*, **42**:

291-299.

- Tóth, M., G. Szöcs, W. Francke, P. M. Guerin, H. Arn and A. Schmid, 1987. Field activity of sex pheromone components of *Peribatodes rhomboidaria*. *Entomol. Exp. Appl.*, **44**: 199-204.
- Tóth, M., G. Szöcs, J. Molnár and I. Szarukán, 1988. Field tests with sex attractants of *Diachrysa chrysitis* and *D. tutti* (Lepidoptera: Noctuidae) at several sites in Hungary. *Z. Naturforsch.*, **43 c**: 463-466.
- Tóth, M., C. Löfstedt, B. S. Hansson, G. Szöcs and A. I. Farag, 1989(a). Identification of four components from the female sex pheromone of the lima-bean pod borer, *Etiella zinckenella*. *Entomol. Exp. Appl.*, **51**: 107-112.
- Tóth, M., H. R. Buser, A. Peña, H. Arn, K. Mori, T. Takeuchi, L. N. Nikolaeva and B. G. Kovalev, 1989(b). Identification of (3Z,6Z)-1,3,6-9,10-epoxyheneicosatriene and (3Z,6Z)-1,3,6-9,10-epoxyeicosatriene in the sex pheromone of *Hyphantria cunea*. *Tetrahedron Lett.*, **30**: 3405-3408.
- Tóth, M., G. Helmchen, U. Leikauf, GY. Sziráki and G. Szöcs, 1989(c). Behavioral activity of optical isomers of 5,9-dimethylheptadecane, the sex pheromone of *Leucoptera scitella* L. (Lepidoptera: Lyonetidae). *J. Chem. Ecol.*, **15**: 1535-1543.
- Tóth, M., G. Szöcs, C. Löfstedt, B. S. Hansson, F. Schmidt and W. Francke, 1991(a). Epoxyheptadecadienes identified as sex pheromone components of *Tephрина arenacearia* Hbn. (Lepidoptera: Geometridae). *Z. Naturforsch.*, **46 c**: 257-263.
- Tóth, M. and G. Szöcs, 1991(b). A sex attractant for male *Swammerdamia pyrella*. *Entomol. Exp. Appl.*, **61**: 95-96.
- Tóth, M. and G. Szöcs, 1991(c). (Z)-7-Dodecenol attractive to male *Agrotis segetum* Den. & Schiff. (Lep., Noctuidae) in the field. *J. Appl. Entomol.*, **112**: 202-206.
- Tóth, M., H. R. Buser, P. M. Guerin, H. Arn, F. Schmidt, W. Francke and G. Szöcs, 1992(a). *Abraxas grossulariata* L. (Lepidoptera: Geometridae): identification of (3Z,6Z,9Z)-3,6,9-heptadecatriene and (6Z,9Z)-6,9-cis-3,4-epoxyheptadecadiene in the female sex pheromone. *J. Chem. Ecol.*, **18**: 13-25.
- Tóth, M. and R. E. Doolittle, 1992(b). Sex attractants for the white-barred groundling moth (*Recurvaria leucatella* Clerck) and the lesser bud moth (*R. nanella* Hübner) (Lepidoptera: Gelechiidae).

- J. Chem. Ecol.*, **18**: 1093-1105.
- Tóth, M., C. Löfstedt, B. W. Blair, T. Cabello, A. I. Farag, B. S. Hansson, B. G. Kovalev, S. Maini, E. A. Nesterov, I. Pajor, A. P. Sazonov, I. V. Shamshev, M. Subchev and G. Szöcs, 1992(c). Attraction of male turnip moths *Agrotis segetum* (Lepidoptera: Noctuidae) to sex pheromone components and their mixtures at 11 sites in Europe, Asia, and Africa. *J. Chem. Ecol.*, **18**: 1337-1347.
- Tóth, M., G. Szöcs, G. Sziráki and W. Sauter, 1992(d). Sex attractants for male Microlepidoptera found in field trapping tests in Hungary. *J. Appl. Entomol.*, **113**: 342-355.
- Tóth, M., R. Gottwald, G. Szöcs and B. Freier, 1992(e). (Z)-11-Hexadecenyl acetate attracting male *Oriamusculosa* Hbn. (Lep., Noctuidae). *J. Appl. Entomol.*, **113**: 509-512.
- Tóth, M., G. Szöcs and M. Bengtsson, 1992(f). A sex attractant for male *Orthosia gothica* L. (Lepidoptera: Noctuidae). *Z. Naturforsch.*, **47c**: 613-617.
- Tóth, M., G. Szöcs and M. Bengtsson, 1993. Optimization of male sex attractants for *Orthosia cruda* Schiff., *O. munda* Schiff. and *O. stabilis* Schiff. (Lep., Noctuidae). *J. Appl. Entomol.*, **115**: 342-349.
- Tóth, M., G. Szöcs, W. Francke, F. Schmidt, P. Philipp, C. Löfstedt, B. S. Hansson and A. I. Farag, 1994(a). Pheromonal production of and response to optically active epoxydienes in some geometrid moths (Lepidoptera: Geometridae). *Z. Naturforsch.*, **49 c**: 516-521.
- Tóth, M., G. Szöcs and G. Sziráki, 1994(b). Male sex attractants for some Lepidopterous pests of the genera *Evergestis* and *Pyrausta* (Lepidoptera: Pyralidae). *Acta Phytopathol. Entomol. Hung.*, **29**: 155-159.
- Tóth, M., G. Szöcs, E. J. van Nieukerken, P. Philipp, F. Schmidt and W. Francke, 1995. Novel type of sex pheromone structure identified from *Stigmella malella* (Stainton) (Lepidoptera: Nepticulidae). *J. Chem. Ecol.*, **21**: 13-27.
- Tóth, M., N. S. Talekar and G. Szöcs, 1996. Optimization of blends of synthetic pheromone components for trapping male limabean pod borers (*Etiella zinckenella* Tr.) (Lepidoptera: Phycitidae): Preliminary evidence on geographical differences. *Bioorg. Medic. Chem.*, **4**: 495-497.
- Tsai, R.-S., E.-C. Yang, C.-Y. Wu, H.-K. Tseng and Y.-S. Chow 1999. A potent sex attractant of the male tea tussock moth, *Euproctis*

- pseudoconspersa* (Strand) (Lepidoptera: Lymantriidae) in Taiwan: Field and EAG responses. *Zool. Studies*, **38**(3): 301-306.
- Tumlinson, J. H., E. R. Mitchell, S. M. Browner and D. A. Lindquist, 1972. A sex pheromone for the soybean looper. *Environ. Entomol.*, **1**: 466-468.
- Tumlinson, J. H., C. E. Yonce, R. E. Doolittle, R. R. Heath, C. R. Gentry and E. R. Mitchell, 1974. Sex pheromones and reproductive isolation of the lesser peachtree borer and the peachtree borer. *Science*, **185**: 614-616.
- Tumlinson, J. H., D. E. Hendricks, E. R. Mitchell, R. E. Doolittle and M. M. Brennan, 1975. Isolation, identification, and synthesis of the sex pheromone of the tobacco budworm. *J. Chem. Ecol.*, **1**: 203-214.
- Tumlinson, J. H., E. R. Mitchell and P. E. Sonnet, 1981. Sex pheromone components of the beet armyworm, *Spodoptera exigua*. *J. Environ. Sci. Health*, **A 16**: 189-200.
- Tumlinson, J. H., E. R. Mitchell, P. E. A. Teal, R. R. Heath and L. J. Mengelkoch, 1986. Sex pheromone of fall armyworm, *Spodoptera frugiperda* (J. E. Smith). Identification of components critical to attraction in the field. *J. Chem. Ecol.*, **12**: 1909-1926.
- Tumlinson, J. H., M. M. Brennan, R. E. Doolittle, E. R. Mitchell, A. Brabham, B. E. Mazomenos, A. H. Baumhover and D. M. Jackson, 1989. Identification of a pheromone blend attractive to *Manduca sexta* (L.) males in a wind tunnel. *Arch. Insect Biochem. Physiol.*, **10**: 255-271.
- Tumlinson, J. H., E. R. Mitchell and H.-S. Yu, 1990. Analysis and field evaluation of volatile blend emitted by calling virgin females of beet armyworm moth, *Spodoptera exigua* (Hübner). *J. Chem. Ecol.*, **16**: 3411-3423.
- Tumlinson, J. H., E. R. Mitchell, R. E. Doolittle and D. M. Jackson, 1994. Field tests of synthetic *Manduca sexta* sex pheromone. *J. Chem. Ecol.*, **20**: 579-591.
- Turgeon, J. J. and G. G. Grant, 1988. Response of *Zeiraphera canadensis* (Lepidoptera: Tortricidae: Olethreutinae) to candidate sex attractants and factors affecting trap catches. *Environ. Entomol.*, **17**: 442-447.
- Twiddle, A. M., D. Barker and D. M. Suckling, 2020. (7Z)-Tricosene improves pheromone trap catch of raspberry bud moth, *Heterocrossa rubophaga*. *J. Chem. Ecol.*, **46**: 830-834.

- Uehara, T., H. Naka, S. Matsuyama, T. Ando and H. Honda, 2012. Identification and field evaluation of sex pheromones in two hawkmoths *Deilephila elpenor lewisii* and *Theretra oldenlandiae oldenlandiae* (Lepidoptera: Sphingidae). *Appl. Entomol. Zool.*, **47** (3): 227-232.
- Uehara, T., H. Naka, S. Matsuyama, L. V. Vang, T. Ando and H. Honda, 2013. Identification of conjugated pentadecadienals as sex pheromone components of the sphingid moth, *Dolbina tancrei*. *J. Chem. Ecol.*, **39**: 1441-1447.
- Uehara, T., H. Naka, S. Matsuyama, T. Ando and H. Honda, 2015. Identification of the sex pheromone of the diurnal hawk moth, *Hemaris affinis*. *J. Chem. Ecol.*, **41**: 9-14.
- Uehara, T., H. Kitahara, H. Naka, S. Matsuyama, T. Ando and H. Honda, 2016. Single-component pheromone consisting of bombykal in a diurnal hawk moth, *Neogurelca himachala sangaica*. *J. Chem. Ecol.*, **42**: 517-522.
- Ujiye, T., M. Wakou, T. Oku, K. Homma, K. Kawasaki, Y. Tamaki and H. Sugie, 1986. Sex pheromone of the apple leafminer moth (*Phyllonorycter ringoniella*): isolation and identification of the pheromone. *Jap. J. Appl. Entomol. Zool.*, **30**: 268-271.
- Ujváry, I., J. C. Dickens, J. A. Kamm and L. M. McDonough, 1993. Natural product analogs: stable mimics of aldehyde pheromones. *Arch. Insect Biochem. Physiol.*, **22**: 393-411
- Underhill, E. W., W. F. Steck and M. D. Chisholm, 1976. Sex pheromone of the clover cutworm moth, *Scotogramma trifolii*: isolation, identification and field studies. *Environ. Entomol.*, **5**: 307-310.
- Underhill, E. W., M. D. Chisholm and W. Steck, 1977(a). Olefinic aldehydes as constituents of sex attractants for noctuid moths. *Environ. Entomol.*, **6**: 333-337.
- Underhill, E. W., W. F. Steck and M. D. Chisholm, 1977(b). A sex pheromone mixture for the bertha armyworm moth, *Mamestra configurata*: (Z)-9-tetradecen-1-ol acetate and (Z)-11-hexadecen-1-ol acetate. *Can. Entomol.*, **109**: 1335-1340.
- Underhill, E. W., W. Steck, M. D. Chisholm, H. A. Worden and J. A. G. Howe, 1978. A sex attractant for the cottonwood crown borer, *Aegeria tibialis* (Lepidoptera: Sesiidae). *Can. Entomol.*, **110**: 495-498.
- Underhill, E. W., A. P. Arthur, M. D. Chisholm and W. F. Steck, 1979. Sex pheromone components of the sunflower moth, *Homoeosoma electellum*: Z-9, E-12-tetradecadienol and Z-9-tetradecenol.

Environ. Entomol., **8**: 740-743.

- Underhill, E. W., M. D. Chisholm and W. Steck, 1980.
(E)-5, (Z)-7-Dodecadienal, a sex pheromone component of the western tent caterpillar, *Malacosoma californicum* (Lepidoptera: Lasiocampidae). *Can. Entomol.*, **112**: 629-631.
- Underhill, E. W., W. F. Steck, J. R. Byers and M. D. Chisholm, 1981.
Biosystematics of the genus *Euxoa* (Lepidoptera: Noctuidae). XVI.
(Z)-5-Decenyl acetate a sex attractant for three closely related species, *Euxoa declarata*, *Euxoa campestris*, and *Euxoa rockburnei*.
Can. Entomol., **113**: 245-249.
- Underhill, E. W., C. E. Rogers, M. D. Chisholm and W. F. Steck, 1982.
Monitoring field populations of the sunflower moth, *Homoeosoma electellum* (Lepidoptera: Pyralidae), with its sex pheromone.
Environ. Entomol., **11**: 681-684.
- Underhill, E. W., P. Palaniswamy, S. R. Abrams, B. K. Bailey, W. F. Steck and M. D. Chisholm, 1983. Triunsaturated hydrocarbons, sex pheromone components of *Caenurgina erechtea*. *J. Chem. Ecol.*, **9**: 1413-1423.
- Underhill, E. W., A. P. Arthur and P. G. Mason, 1986. Sex pheromone of the banded sunflower moth, *Cochylis hospes* (Lepidoptera: Cochylidae): identification and field trapping. *Environ. Entomol.*, **15**: 1063-1066.
- Underhill, E. W., J. G. Millar, R. A. Ring, J. W. Wong, D. Barton and M. Giblin, 1987(a). Use of a sex attractant and an inhibitor for monitoring winter moth and Bruce spanworm populations. *J. Chem. Ecol.*, **13**: 1319-1330.
- Underhill, E. W., C. E. Rogers and L. R. Hogge, 1987(b). Sex attractants for two sunflower pests, *Eucosma womonana* (Lepidoptera: Tortricidae) and *Isophrictis similiella* (Lepidoptera: Gelechiidae). *Environ. Entomol.*, **16**: 463-466.
- Unelius, C. R., A. Eiras, P. Witzgall, M. Bengtsson, A. Kovaleski, E. F. Vilela and A.-K. Borg-Karlson, 1996. Identification and synthesis of the sex pheromone of *Phtheochroa cranaodes* (Lepidoptera: Tortricidae). *Tetrahedron Lett.*, **37**: 1505-1508.
- Usui, K., M. Kurihara, K. Uchiumi, J. Fukami and S. Tatsuki, 1987. Sex pheromone of the female large cabbage-heart caterpillar, *Crociodolomia binotalis* Zeller (Lepidoptera: Pyralidae). *Agric. Biol. Chem.*, **51**: 2191-2195.
- Vakenti, J. M., A. P. Gaunce, K. N. Slessor, G. G. S. King, S. A. Allan, H. F. Madsen and J. H. Borden, 1988. Sex pheromone components

- of the oblique-banded leafroller, *Choristoneura rosaceana* in the Okanagan Valley of British Columbia. *J. Chem. Ecol.*, **14**: 605-621.
- Van de Veire, M. and P. Dirinck, 1986. Sex pheromone components of the cabbage armyworm, *Mamestra brassicae*: isolation, identification and field experiments. *Entomol. Exp. Appl.*, **41**: 153-155.
- Vang, L. V., S. Inomata, M. Kinjo, F. Komai and T. Ando, 2005. Sex pheromones of five olethreutine species (Lepidoptera: Tortricidae) associated with the seedlings and fruits of mangrove plants in the Ryukyu Islands, Japan: identification and field evaluation. *J. Chem. Ecol.*, **31**(4): 859-878.
- Vang, L. V., M. Ishitani, F. Komai, M. Yamamoto and T. Ando, 2006. Sex pheromone of the soybean pod borer, *Leguminivora glycinivorella* (Lepidoptera: Tortricidae): Identification and field evaluation. *Appl. Entomol. Zool.*, **41** (3): 507-513.
- Vang, L. V., Md. A. Islam, N. D. Do, T. V. Hai, S. Koyano, Y. Okahana, N. Ohbayashi, M. Yamamoto and T. Ando, 2008. 7,11,13-Hexadecatrienal identified from female moths of the citrus leafminer as a new sex pheromone component: synthesis and field evaluation in Vietnam and Japan. *J. Pestic. Sci.*, **33**: 152-158.
- Vang, V. L., N. D. Do, L. K. An, P. K. Son and T. Ando, 2011. Sex pheromone components and control of the citrus pock caterpillar, *Prays endocarpa*, found in the Mekong Delta of Vietnam. *J. Chem. Ecol.*, **37**: 134-140.
- Vang, L. V., C. N. Q. Khanh, H. Shibasaki and T. Ando, 2012. Female sex pheromone secreted by *Carmentia mimosa* (Lepidoptera: Sesiidae), a biological control agent for an invasive weed in Vietnam. *Biosci. Biotechnol. Biochem.*, **76**: 2153-2155.
- Vang, L. V., H. N. Thuy, C. N. Q. Khanh, P. K. Son, Q. Yan, M. Yamamoto, U. Jinbo and T. Ando, 2013. Sex pheromones of three citrus leafrollers, *Archips atrolucens*, *Adoxophyes privata*, and *Homona* sp., inhabiting the Mekong Delta of Vietnam. *J. Chem. Ecol.*, **39**: 783-789.
- Vang, L. V., Q. Yan, N. T. N. Nghia, C. N. Q. Khanh and T. Ando, 2018. Unsaturated cuticular hydrocarbon components of the sex pheromone of eggplant fruit borer, *Leucinodes orbonalis* Guenée (Lepidoptera: Crambidae). *J. Chem. Ecol.*, **44**: 631-636.
- Vang, L. V., T. T. Thy, D. K. Hanh, T. P. Linh, M. Yamamoto and T. Ando, 2020. Sex pheromone analysis and effective attraction of males of the cabbage webworm, *Hellula undalis*, inhabiting the

- Mekong Delta of Vietnam. *J. Asia-Pac. Entomol.*, **23**: 935-941.
- Vasian, I., M. L. Duduman, E. Gal, A. Pîrnău and M. Gorgan, 2025. Green synthesis, formulation and test field of *Lymantria monacha* L. (Lepidoptera: Lymantriidae) sex pheromone in east European region. *Int. J. Mol. Sci.*, **26**: 568.
- Velcheva, N. and T. Milkova, 1990. Sex attractant for *Exapate congelatella* Cl (Lepidoptera, Tortricidae) - a new species for the Bulgarian fauna. *Ekologiya*, **23**: 75-78.
- Vetter, R. S. and T. C. Baker, 1983. Behavioral responses of male *Heliothis virescens* in a sustained-flight tunnel to combinations of seven compounds identified from female sex pheromone glands. *J. Chem. Ecol.*, **9**: 747-759.
- Vick, K. W., H. C. F. Su, L. L. Sower, P. G. Mahany and P. C. Drummond, 1974. (Z,E)-7,11-Hexadecadien-1-ol acetate: the sex pheromone of the Angoumois grain moth, *Sitotroga cerealella*. *Experientia*, **30**: 17-18.
- Vickers, N. J., 2002. Defining a synthetic pheromone blend attractive to male *Heliothis subflexa* under wind tunnel conditions. *J. Chem. Ecol.*, **28**(6): 1255-1267.
- Voerman, S. and W. M. Herrebout, 1978(a). A sex attractant for the leaf miner moth *Lithocolletis corylifoliella* and its influence on that of *L. blancardella* (Lep., Gracillariidae). *Entomol. Exp. Appl.*, **23**: 96-98.
- Voerman, S., A. K. Minks, G. Vanwetswinkel and J. H. Tumlinson, 1978(b). Attractivity of 3,13-octadecadien-1-ol acetates to the male clearwing moth *Synanthedon myopaeformis* (Borkhausen) (Lepidoptera, Sesiidae). *Entomol. Exp. Appl.*, **23**: 301-304.
- Voerman, S. and G. H. L. Rothschild, 1978(c). Synthesis of the two components of the sex pheromone system of the potato tuberworm moth, *Phthorimaea operculella* (Zeller) (Lepidoptera: Gelechiidae) and field experience with them. *J. Chem. Ecol.*, **4**: 531-542.
- Voerman, S., H. Audemard and E. Priesner, 1983. Sex attractants for clearing moths: *Synanthedon vespiformis* and *Chamaesphecia tenthrediniformis* (and/or *C. empiformis*). *Entomol. Exp. Appl.*, **34**: 203-205.
- Voerman, S., C. J. Persoons and E. Priesner, 1984. Sex attractant for currant clearwing moth, *Synanthedon tipuliformis* (Clerck) (Lepidoptera: Sesiidae). *J. Chem. Ecol.*, **10**: 1371-1376.

- Voerman, S., 1988. The pheromone bank: a collection of unsaturated compounds indispensable for discovery of sex attractants for Lepidoptera. *Agric. Ecosyst. Environ.*, **21**: 31-41.
- Voerman, S., 1991. A mono-unsaturated acetate as sex attractant for the leaf miner moth *Lithocolletis corylifoliella* (Hübner) (Lep., Gracillariidae). *J. Appl. Entomol.*, **112**: 207-210.
- Vrkoc, J., B. G. Kovalev and V. A. Starets, 1981. Isolation and identification of the main component of the female sexual pheromone of *Mamestra suasa* (Lepidoptera, Noctuidae). *Acta Entomol. Bohemoslov.*, **78**: 353-357.
- Vrkoc, J., V. P. Konyukhov and B. G. Kovalev, 1983. Isolation and identification of main components of the pheromone complex of the moth *Agrotis exclamationis* (Lepidoptera, Noctuidae). *Acta Entomol. Bohemoslov.*, **80**: 184-189.
- Vu, M. H., T. Ando, N. Takahashi, S. Tatsuki, A. Yamane, T. Ikeda and S. Yamazaki, 1980. Identification of the female sex pheromone of the pine moth. *Agric. Biol. Chem.*, **44**: 231-233.
- Wakamura, S., 1978. Sex attractant pheromone of the common cutworm moth, *Agrotis fucosa* Butler (Lepidoptera: Noctuidae): isolation and identification. *Appl. Entomol. Zool.*, **13**: 290-295.
- Wakamura, S., 1980. Sex attractant pheromone of the common cutworm moth *Agrotis fucosa* Butler: field evaluation. *Appl. Entomol. Zool.*, **15**: 167-174.
- Wakamura, S., 1985. Identification of sex-pheromone components of the podborer, *Matsumuraesia falcana* (Walshingham) (Lepidoptera: Tortricidae). *Appl. Entomol. Zool.*, **20**: 189-198.
- Wakamura, S., D. L. Struble, H. Matsuura, M. Sato and K. Kegasawa, 1986(a). Sex pheromone of the black cutworm moth, *Agrotis ipsilon* Hufnagel (Lepidoptera: Noctuidae): attractant synergist and improved formulation. *Appl. Entomol. Zool.*, **21**: 299-304.
- Wakamura, S. and K. Kegasawa, 1986(b). Sex pheromone of the podborer, *Matsumuraesia falcana* (Walshingham) (Lepidoptera: Tortricidae): activity of the third component, (E,Z)-7,9-dodecadienyl acetate, and 3-component formulation. *Appl. Entomol. Zool.*, **21**: 334-339.
- Wakamura, S., 1987. Sex pheromone of the beet armyworm, *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae): Field attraction of male moths in Japan to (Z,E)-9,12-tetradecadienyl acetate and (Z)-9-tetradecen-1-ol. *Appl. Entomol. Zool.*, **22**: 348-351.
- Wakamura, S., T. Yasuda, A. Ichikawa, T. Fukumoto and F. Mochizuki,

1994. Sex attractant pheromone of the tea tussock moth, *Euproctis pseudoconspersa* (Strand) (Lepidoptera: Lymantriidae): identification and field attraction. *Appl. Entomol. Zool.*, **29**: 403-411.
- Wakamura, S., T. Yasuda, M. Watanabe, K. Kiguchi, M. Shimoda and T. Ando, 1996(a). Sex pheromone of the sweetpotato hornworm, *Agrius convolvuli* (L.) (Lepidoptera: Sphingidae): identification of a major component and its activity in a wind tunnel. *Appl. Entomol. Zool.*, **31**: 171-174.
- Wakamura, S., A. Ichikawa, T. Yasuda, N. Arakaki and T. Fukumoto, 1996(b). EAG and field responses of the male tea tussock moth, *Euproctis pseudoconspersa* (Strand) (Lepidoptera: Lymantriidae) to (R)- and (S)-enantiomers and racemic mixture of 10,14-dimethylpentadecyl isobutyrate. *Appl. Entomol. Zool.*, **31**(4): 623-625.
- Wakamura, S., N. Arakaki, T. Yasuda and K. Kawasaki, 1997. Sex pheromone of *Statherotis discana* (Felder et Rogenhofer) (Lepidoptera: Tortricidae), a pest of litchi leaves: identification and field attraction. *Appl. Entomol. Zool.*, **32**(1): 267-269.
- Wakamura, S., N. Arakaki, K. Kinjo and T. Yasuda 1998. Sex pheromone of the cotton caterpillar, *Diaphania indica* (Saunders) (Lepidoptera: Pyralidae): Identification and field attraction. *Appl. Entomol. Zool.*, **33**(3): 429-434.
- Wakamura, S., M. Hattori, K. Igita, K. Yasuda and Tridjaka, 1999. Sex pheromone of *Etiella behrii*, a pod borer of soybean in Indonesia: identification and field attraction. *Entomol. Exp. Appl.*, **91**: 413-420.
- Wakamura, S., N. Arakaki, M. Yamamoto, S. Hiradate, H. Yasui, T. Yasuda, and T. Ando, 2001(a). Posticlude: a Novel *trans*-Epoxide as a sex Pheromone Component of the Tussock Moth, *Orgyia postica* (Walker). *Tetrahedron Lett.*, **42**(4): 687-689.
- Wakamura, S., A. Tanaka, K. Yasuda and T. Yasuda, 2001(b). Sex pheromone of *Acrolepiopsis nagaimo* Yasuda (Lepidoptera: Acrolepiidae), a predominant leafminer on Chinese yam in Tottori Prefecture: identification and field attraction. *Appl. Entomol. Zool.*, **36**(1): 71-76.
- Wakamura, S., N. Arakaki, H. Ono and H. Yasui, 2001(c). Identification of novel sex pheromone components from a tussock moth, *Euproctis pulverea*. *Entomol. Exp. Appl.*, **100**: 109-117.
- Wakamura, S., N. Arakaki, H. Yamazawa, N. Nakajima, M. Yamamoto and T. Ando, 2002. Identification of epoxyhenicoxadiene and novel diepoxy derivatives as sex pheromone components of the

- clear-winged tussock moth *Perina nuda*. *J. Chem. Ecol.*, **28**(3): 449-467.
- Wakamura, S. and N. Arakaki, 2004. Sex pheromone components of pyralid moths *Terastia subjectalis* and *Agathodes ostentalis* feeding on the coral tree, *Erithrina variegata*: two sympatric species share common components in different ratios. *Chemoecology*, **14**: 181-185.
- Wakamura, S., N. Arakaki, M. Yamamoto, S. Hiradate, H. Yasui, K. Kinjo, T. Yasuda, H. Yamazawa and Tetsu Ando, 2005(a). Sex pheromone and related compounds in the Ishigaki and Okinawa strains of the tussock moth *Orgyia postica* (Walker) (Lepidoptera: Lymantriidae). *Biosci. Biotechnol. Biochem.*, **69** (5): 957-965.
- Wakamura, S., N. Arakaki and K. Kinjo, 2005(b). Sex pheromone components of an olethreutid moth, *Strepsicrates semicanella* (Walker) (Lepidoptera: Tortricidae), a pest of guava and eucalyptus in Okinawa. *Appl. Entomol. Zool.*, **40**(4): 637-642.
- Wakamura, S., H. Tanaka, Y. Masumoto, H. Sawada and N. Toyohara, 2007(a). Sex pheromone of the blue-striped nettle grub moth *Parasa lepida* (Cramer) (Lepidoptera: Limacodidae): identification and field attraction. *Appl. Entomol. Zool.*, **42** (3): 347-352.
- Wakamura, S., T. Yasuda, Y. Hirai, H. Tanaka, T. Doki, Y. Nasu, M. Shibao, A. Yunotani and K. Kadono, 2007(b). Sex pheromone of the oriental tussock moth *Artaxa subflava* (Bremer) (Lepidoptera: Lymantriidae): identification and field attraction. *Appl. Entomol. Zool.*, **42** (3): 375-382.
- Wakamura, S., S. Ohno, N. Arakaki, T. Kohama, D. Haraguchi and H. Yasui, 2010. Identification and field activity of the sex pheromone component of the sweetpotato vine borer moth *Omphisa anastomosalis* (Lepidoptera: Crambidae). *Appl. Entomol. Zool.*, **45** (4): 635-640.
- Wakamura, S., H. Yasui, M. Shimatani, T. Tokashiki, T. Okuhira, N. Fujiwara-Tsujii, M. Ishikawa, T. Uesato, H. Zukeyama and H. Tobaru, 2011. Sex pheromone of the African armyworm *Spodoptera exempta* (Lepidoptera: Noctuidae): identification of components of the Okinawan population and formulation for population monitoring. *Appl. Entomol. Zool.*, **46**: 415-420.
- Wakamura, S., N. Arakaki, S. Yoshimatsu, 2021. Sex pheromone of the fall armyworm *Spodoptera frugiperda* (Lepidoptera: Noctuidae) of a "Far East" population from Okinawa, Japan. *Appl. Entomol. Zool.*, **56**: 19-25.

- Walker, P. W., G. R. Allen, N. W. Davies, J. A. Smith, P. P. Molesworth, A. Nilsson, F. Andersson and E. Hedenström, 2009. Identification, synthesis and field testing of (3Z,6Z,9Z)-3,6,9-henicosatriene, a second bioactive component of the sex pheromone of the autumn gum moth, *Mnesampela privata*. *J. Chem. Ecol.*, **35**: 1411-1422.
- Wall, C., A. G. Greenway and P. E. Burt, 1976. Electroantennographic and field responses of the pea moth, *Cydia nigricana*, to sex attractants and related compounds. *Physiol. Entomol.*, **1**: 151-157.
- Wall, C., R. Bourneville, A. R. Greenway and J. Einhorn, 1987. Efficacy and stability of sex-attractant lures for *Cydia nigricana* (F.) and *Cydia medicaginis* (Kuzn.). *Crop Prot.*, **6**: 7-12.
- Wallin, E. A., B. Kalinová, J. Kindl, E. Hedenström and I. Valterová, 2020. Stereochemistry of two pheromonal components of the bumblebee wax moth, *Aphomia sociella*. *Sci. Rep.*, **10**: 2094.
- Wallner, W. E., R. T. Cardé, C. Xu, R. M. Weseloh, S. Xilin, J. Yan and P. W. Schaefer, 1984. Gypsy moth (*Lymantria dispar* L.) attraction to disparlure enantiomers and the olefin precursor in the People's Republic of China. *J. Chem. Ecol.*, **10**: 753-757.
- Wang, X.-W. and M.-Y. Liu 1997. Extract components of single sex pheromone gland of oriental armyworm *Mythimna separata* Walker with trace analytical techniques. *Acta Entomol. Sinica (Kunchong Xuebao)*, **40**: 158-165.
- Wang, H.-L., G. P. Svensson, O. Rosenberg, M. Bengtsson, E. V. Jirle and C. Löfstedt, 2010. Identification of the sex pheromone of the spruce seed moth, *Cydia strobilella* L. *J. Chem. Ecol.*, **36**: 305-313.
- Wang, H. L., G. P. Svensson, J. Jakobsson, E. V. Jirle, O. Rosenberg, W. Francke, O. Anderbrant, J. G. Millar and C. Löfstedt, 2015(a). Sex pheromone of the cloaked pug moth, *Eupithecia abietaria* (Lepidoptera: Geometridae), a pest of spruce cones. *J. Appl. Entomol.*, **139**: 352-360.
- Wang, H.-L., H. Geertsema, E. J. van Nieuwerkerken and C. Löfstedt, 2015(b). Identification of the female-produced sex pheromone of the leafminer *Holocacista capensis* infesting grapevine in South Africa. *J. Chem. Ecol.*, **41**: 724-731.
- Wang, H.-L., M. Baldessari, G. Anfora, E. J. van Nieuwerkerken and C. Löfstedt, 2019. Sex pheromones of two leafminer species, *Antispila oinophylla* and *Holocacista rivillei* (Lepidoptera: Heliozelidae) infesting grapevine in Italy *J. Chem. Ecol.*, **45**: 1-8.

- Wang, C., S. Zhang, M.-B. Guo, X.-Q. Hou, B. Yang and G.-R. Wang, 2022. Optimization of a pheromone lure by analyzing the peripheral coding of sex pheromones of *Spodoptera frugiperda* in China. *Pest Manag. Sci.*, **78**: 2995-3004.
- Wang, C., Y.-H. Li, L. Wang, B. Yang and G.-R. Wang, 2023. Development of a new sex attractant via the peripheral coding of pheromones in *Mythimna loreyi*. *J. Agric. Food Chem.*, **71**: 2795-2803.
- Wang, L., C. Yang, X. Mei, B. Guo, X. Yang, T. Zhang and J. Ning, 2024. Identification of sex pheromone in *Macdunnoughia crassisigna* Warren (Lepidoptera: Noctuidae) and field optimization of the sex attractant. *Pest Manag. Sci.*, **80**: 577-585.
- Warthen, D. and M. Jacobson, 1967. Insect sex attractants. VII. 5,9-Tridecadien-1-ol acetates. *J. Med. Chem.*, **10**: 1190-1191.
- Warthen, J. D., J. A. Klun and E. D. DeVilbiss, 1996. Structure-activity relationship observations for the bagworm moth pheromone. *J. Chem. Ecol.*, **22**: 1315-1324.
- Weatherston, J., W. Roelofs, A. Comeau and C. J. Sanders, 1971. Studies of physiologically active arthropod secretions. X. Sex pheromone of the eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **103**: 1741-1747.
- Weatherston, J., L. M. Davidson and D. Simonini, 1974(a). Attractants for several male forest Lepidoptera. *Can. Entomol.*, **106**: 781-782.
- Weatherston, J. and W. Maclean, 1974(b). The occurrence of (E)-11-tetradecen-1-ol, a known sex attractant inhibitor, in the abdominal tips of virgin female eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **106**: 281-284.
- Weatherston, J., J. E. Percy and L. M. MacDonald, 1976. Field testing of *cis*-11-tetradecenal as attractant or synergist in Tortricinae. *Experientia*, **32**: 178-179.
- Weatherston, J., A. F. Hedlin, D. S. Ruth, L. M. MacDonald, C. C. Leznoff and T. M. Fyles, 1977. Chemical and field studies on the sex pheromones of the cone and seed moths *Barbara colfaxiana* and *Laspeyresia youngana*. *Experientia*, **33**: 723-725.
- Weatherston, J., G. G. Grant, L. M. MacDonald, D. Frech, R. A. Werner, C. C. Leznoff and T. M. Fyles, 1978. Attraction of various tortricine moths to blends containing *cis*-11-tetradecenal. *J.*

- Chem. Ecol.*, **4**: 543-549.
- Wei, H.-Y. and J.-W. Du, 2003. Identification of active components of sex pheromone for *Sidemna depravata* and field trapping. *Yingyong Shengtai Xuebao*, **14**: 730-732.
- Wei, Z.-H. and F.-M. Pan, 1985. Preliminary report on the sex pheromone of the armyworm *Mythimna separata*. *Kunchong Xuebao (Acta Entomol. Sinica)*, **28**: 348-350.
- Wen, X.-J., B. Kalinová, M.-G. Liu and M. Hoskovec, 2009. Candidate sex pheromone components of persimmon bark borer, *Euzophera batangensis*. *Linze Kexue*, **45**: 83-89. [*SciFinder*. AN2010:149071]
- Wen, X., T. Ma, Z. Liu, C. Sun and W. Cai, 2014. Sex attractant for controlling *Diaphania glauculalis*. CN 103875661. [CAN 161:180686]
- Whittle, C. P., T. E. Bellas, M. Horak and B. Pinese, 1987. The sex pheromone and taxonomic status of *Homona spargotis* Meyrick sp. rev., an Australian pest species of the *Coffearia* group (Lepidoptera: Tortricidae: Tortricinae). *J. Aust. Entomol. Soc.*, **26**: 169-179.
- Whittle, C. P., T. E. Bellas and A. L. Bishop, 1991. Sex pheromone of lucerne leafroller, *Merophyas divulsana* (Walker) (Lepidoptera: Tortricidae): evidence for two distinct populations. *J. Chem. Ecol.*, **17**: 1883-1895.
- Whittle, C. P., R. A. Vickers, L. S. Kuniata, T. E. Bellas and E. R. Rumbo, 1995. Identification of an attractant for the cane borer *Sesamia grisescens* Walker (Lepidoptera: Noctuidae). *J. Chem. Ecol.*, **21**: 1409-1420.
- Wiesner, C. J., P. J. Silk, S.-H. Tan, P. Palaniswamy and J. O. Schmidt, 1979. Components of the sex pheromone gland of the eastern spruce budworm, *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Can. Entomol.*, **111**: 1311.
- Willemse, L. P. M., C. J. H. Booij and S. Voerman, 1987. New sex attractants for male Lepidoptera (Coleophoridae, Gelechiidae, Momphidae, Oecophoridae, and Yponomeutidae) found by field screening in the Netherlands. *J. Appl. Entomol.*, **103**: 508-515.
- Williams, R. N., D. T. Johnson and E. Priesner 1999. *Synanthedon rileyana* (H. Edwards) response to selected clearwing pheromone blends. *J. Entomol. Sci.*, **34** (2): 219-224.
- Witjaksono, K. Ohtani, M. Yamamoto, T. Miyamoto and T. Ando 1999. Responses of Japanese giant looper male moth to synthetic sex pheromone and related compounds. *J. Chem. Ecol.*, **25**(7):

1633-1642.

- Witzgall, P. and B. Frérot, 1989(a). Pheromone emission by individual females of carnation tortrix, *Cacoecimorpha pronubana*. *J. Chem. Ecol.*, **15**: 707-717.
- Witzgall, P., W. Sauter, H.-R. Buser, S. Rauscher, H. Arn, P. J. Charmillot and Th. Wildbolz, 1989(b). Use of pheromone chemistry to identify *Grapholita lobarzewskii* as an occasional pest of apple and plum. *Entomol. Exp. Appl.*, **53**: 133-136.
- Witzgall, P., 1990. Attraction of *Cacoecimorpha pronubana* male moths to synthetic sex pheromone blends in the wind tunnel. *J. Chem. Ecol.*, **16**: 1507-1515.
- Witzgall, P., M. Bengtsson, H.-R. Buser, P.-J. Chambon, E. Priesner, T. Wildbolz and H. Arn, 1991(a). Sex pheromone of *Spilonota ocellana* and *Spilonota laricana*. *Entomol. Exp. Appl.*, **60**: 219-223.
- Witzgall, P. P., B. Frérot and C. Malosse, 1991(b). Identification of the major component of the pheromone secretion of *Notocelia uddmanniana* L. (Lep., Tortricidae). *J. Appl. Entomol.*, **112**: 71-75.
- Witzgall, P., M. Bengtsson, C. R. Unelius and J. Löfqvist, 1993. Attraction of pea moth *Cydia nigricana* F. (Lepidoptera: Tortricidae) to female sex pheromone (*E,E*)-8,10-dodecadien-1-yl acetate, is inhibited by geometric isomers *E,Z*, *Z,E*, and *Z,Z*. *J. Chem. Ecol.*, **19**: 1917-1928.
- Witzgall, P., J.-P. Chambon, M. Bengtsson, C. R. Unelius, M. Appelgren, G. Makranczy, N. Muraleedharan, D. W. Reed, K. Hellrigl, H.-R. Buser, E. Hallberg, G. Bergström, M. Tóth, C. Löfstedt and J. Löfqvist, 1996. Sex pheromone and attractants in the Eucosmini and Grapholitini (Lepidoptera, Tortricidae). *Chemoecology*, **7**: 13-23.
- Witzgall, P., M. Bengtsson and R. M. Trimble, 2000. Sex pheromone of grape berry moth (Lepidoptera: Tortricidae). *Environ. Entomol.*, **29**(3): 433-436.
- Witzgall, P., M. Tasin, H.-R. Buser, G. Wegner-Kiß, V. S. M. Mancebón, C. Ioriatti, A.-C. Bäckman, M. Bengtsson, L. Lehmann and W. Francke, 2005. New pheromone components of the grapevine moth *Lobesia botrana*. *J. Chem. Ecol.* **31**(12): 2923-2932.
- Wong, J. W., P. Palaniswamy, E. W. Underhill, W. F. Steck and M. D. Chisholm, 1984(a). Novel sex pheromone components from the fall cankerworm moth, *Alsophila pometaria*. *J. Chem. Ecol.*, **10**:

463-473.

- Wong, J. W., P. Palaniswamy, E. W. Underhill, W. F. Steck and M. D. Chisholm, 1984(b). Sex pheromone components of fall cankerworm moth, *Alsophila pometaria*. Synthesis and field trapping. *J. Chem. Ecol.*, **10**: 1579-1596.
- Wong, H. R., J. A. Drouin and C. L. Rentz, 1985(a). *Petrova albicapitana* (Busck) and *P. metallica* (Busck) (Lepidoptera: Tortricidae) in *Pinus contorta* Dougl. stands of Alberta. *Can. Entomol.*, **117**: 1463-1470.
- Wong, J. W., E. W. Underhill, S. L. MacKenzie and M. D. Chisholm, 1985(b). Sex attractants for geometrid and noctuid moths. Field trapping and electroantennographic responses to their hydrocarbons and monoepoxydiene derivatives. *J. Chem. Ecol.*, **11**: 727-756.
- Wu, D.-M. and G.-L. Li, 1982. EAG analysis and field attractant effect of sex pheromone of the striped sugarcane borer. *Kunchong Xuebao (Acta Entomol. Sinica)*, **25**: 358-362.
- Wu, D.-M., Y.-H. Yan, B.-H. Wang and H.-L. Huang, 1984. The chemical structure and field attractant effect of the sex pheromone of sugarcane stem borer *Chilo infuscatellus* (Snellen). *Kunchong Xuebao (Acta Entomol. Sinica)*, **27**: 368-374.
- Wu, D.-M. and J.-R. Cui, 1986(a). Electrophysiological responses of the purple stem borer to sex pheromone and related compounds. *Kunchong Xuebao (Acta Entomol. Sinica)*, **29**: 239-245.
- Wu, D.-M., Z.-R. Ding, J.-R. Cui and Y.-H. Yan, 1986(b). The study of sex attractant of the pinetip moth *Dioryctria rubella* Hampson. *Scientia Silvae Sinicae*, **22**: 368-372.
- Wu, D., Y. Yan, F. Kong, W. Li, S. Yu, J. Li and W. Zeng, 1990(a). Sex pheromone of the sugarcane striped borer -- identification of three components. *Youji Huaxue*, **10**: 47-49. [*Chem. Abst.*, **112**: 174029n]
- Wu, D.-M., Y.-H. Yan, J.-R. Cui, M.-Y. Liu, D.-F. Ren, A. Chen and Q. Zhang, 1990(b). Sex pheromone of sugarcane tip borer *Scirpophaga excerptalis* Walker. *Chin. Sci. Bull.*, **35**: 1469-1473.
- Wu, D.-M., J.-R. Cui and Y.-H. Yan, 1993. Electrophysiological analysis and field trapping results of sex pheromone of the reed stalk borer. *Acta Phytophylacica Sinica*, **20**: 229-232.
- Wu, W.-Q., B. S. Hansson and C. Löfstedt, 1995. Electrophysiological and behavioural evidence for a fourth sex pheromone component in the turnip moth, *Agrotis segetum*. *Physiol. Entomol.*, **20**:

81-92.

- Wu, W.-Q, C. B. Cottrell, B. S. Hansson and C. Löfstedt 1999. Comparative study of pheromone production and response in Swedish and Zimbabwean populations of turnip moth, *Agrotis segetum*. *J. Chem. Ecol.*, **25**: 177-196.
- Wu, J., X. Wu, H. Chen, L. Xu, G. Liu, B. Mao, R. Quo and Y. Du, 2013. Optimization of the sex pheromone of the rice leaf folder moth *Cnaphalocrocis medinalis* as a monitoring tool in China. *J. Appl. Entomol.*, **137**: 509-518.
- Wunderer, H., K. Hansen, T. W. Bell, D. Schneider and J. Meinwald, 1986. Sex pheromones of two Asian moths (*Cretonotos transiens*, *C. gangis*; Lepidoptera - Arctiidae): behavior, morphology, chemistry and electrophysiology. *Exp. Biol.*, **46**: 11-27.
- Xiao, W. S. Matsuyama, T. Ando, J. G. Millar and H. Honda, 2012. Body wax unsaturated hydrocarbons as synergist of sex pheromone in the yellow peach moth, *Conogethes punctiferalis* (Crambidae: Lepidoptera). *J. Chem. Ecol.*, **38**: 1143-1150.
- Yadav, J. S., M. Y. Valli and A. R. Prasad, 2001. Isolation, identification, synthesis, and bioefficacy of female *Diacrisia oblique* (Arctiidae) sex pheromone blend. An Indian agricultural pest. *Pure Appl. Chem.*, **73**: 1157-1162.
- Yaginuma, K., M. Kumakura, Y. Tamaki, T. Yushima and J. H. Tumlinson, 1976. Sex attractant for the cherry tree borer, *Synanthedon hector* Butler (Lepidoptera: Sesiidae). *Appl. Entomol. Zool.*, **11**: 266-268.
- Yamada, H. and T. Koshihara, 1980. Interspecific attractiveness of female sex pheromone components of the diamondback moth, *Plutella xylostella* (L.), and their analogue to the stone leek miner, *Acrolepiopsis sapporensis* Matsumura. *Kontyû*, **48**: 104-110.
- Yamakawa, R., N. D. Do, Y. Adachi, M. Kinjo and T. Ando, 2009. (6Z,9Z,12Z)-6,9,12-Octadecatriene and (3Z,6Z,9Z,12Z)-3,6,9,12-icosatetraene, the novel sex pheromones produced by emerald moths. *Tetrahedron Lett.*, **50**: 4738-4740.
- Yamakawa, R., N. D. Do, K. Kinjo, Y. Terashima and T. Ando, 2011(a). Novel components of the sex pheromones produced by emerald moths: Identification, synthesis, and field evaluation. *J. Chem. Ecol.*, **37**: 105-113.
- Yamakawa, R., R. Kiyota, T. Taguri and T. Ando, 2011(b). (5R,7R)-5-Methylheptadecan-7-ol: a novel sex pheromone

- component produced by a female lichen moth, *Miltochrista calamine*, in the family Arctiidae. *Tetrahedron Lett.*, **52**: 5808-5811.
- Yamakawa, R., Y. Takubo, K. Ohbayashi, H. Naka and T. Ando, **2012** (a). Female sex pheromone of *Cystidia couaggaria couaggaria* (Lepidoptera: Geometridae): Identification and field attraction. *Biosci. Biotechnol. Biochem.*, **76**: 1303-1307.
- Yamakawa, R., Y. Takubo, H. Shibasaki, Y. Murakami, M. Yamamoto and T. Ando, **2012** (b). Characterization of epoxytrienes derived from (3Z,6Z,9Z)-1,3,6,9-tetraenes, sex pheromone components of arctiid moths and related compounds. *J. Chem. Ecol.*, **38**: 1042-1049.
- Yamamoto, M., Y. Takeuchi, Y. Ohmasa, H. Yamazawa and T. Ando, **1999**. Chiral HPLC resolution of monoepoxides derived from 6,9-dienes and its application to stereochemistry assignment of fruit-piercing noctuid pheromone. *Biomed. Chromatogr.*, **13**: 410-417.
- Yamamoto, M., M. Kiso, H. Yamazawa, J. Takeuchi and T. Ando, **2000**. Identification of chiral sex pheromone secreted by giant geometrid moth, *Biston robustum* Butler. *J. Chem. Ecol.*, **26** (11): 2579-2590.
- Yamamoto, M., T. Kamata, N. D. Do, Y. Adachi, M. Kinjo and T. Ando, **2007**. A novel lepidopteran sex pheromone produced by females of a Lithosiinae species, *Lyclene dharma dharma*, in the family of Arctiidae. *Biosci. Biotechnol. Biochem.*, **71** (11): 2860-2863.
- Yamamoto, M., R. Yamakawa, T. Oga, Y. Takei, M. Kinjo and T. Ando, **2008**. Synthesis and chemical characterization of hydrocarbons with a 6,9,11-, 3,6,9,11-, or 1,3,6,9-polyene system, pheromone candidates in Lepidoptera. *J. Chem. Ecol.*, **34** (8): 1057-1064.
- Yamaoka, R., H. Fukami and S. Ishii, **1976**. Isolation and identification of the female sex pheromone of the potato tuberworm moth, *Phthorimaea operculella* (Zeller). *Agric. Biol. Chem.*, **40**: 1971-1977.
- Yamaoka, R., Y. Shiraishi, T. Ueno, Y. Kuwahara and H. Fukami, **1985**. Structure elucidation of koiganal I and II, the sex pheromones of the webbing clothes moth, using capillary GC/MS. *Mass Spectro.*, **33**: 189-195.
- Yamaoka, R., Y. Kuwabara, M. Ogawa, S. Nakamura and R. Muneyuki, **1990**. Preparation of *trans*-2-octadecenal as insect attractant, for *Tinea pellionella*. *Jpn. Kokai Tokkyo Koho* JP 02,243,605.

- Yamazawa, H., N. Nakajima, S. Wakamura, N. Arakaki, M. Yamamoto and T. Ando, 2001. Synthesis and characterization of diepoxyalkenes derived from (3Z,6Z,9Z)-trienes: lymantriid sex pheromones and their candidates. *J. Chem. Ecol.*, **27**(11): 2153-2167.
- Yan, Q., L. V. Vang, C. N. Q. Khanh, H. Naka and T. Ando, 2014. Reexamination of the female sex pheromone of the sweet potato vine borer moth: identification and field evaluation of a tricosatriene. *J. Chem. Ecol.*, **40**: 590-598.
- Yan, Q., A. Kanegae, T. Miyachi, H. Naka, H. Tatsuta and T. Ando, 2015(a). Female sex pheromones of two Japanese saturniid species, *Rhodinia fugax* and *Loepa sakaei*: identification, synthesis, and field evaluation. *J. Chem. Ecol.*, **41**: 1-8.
- Yan, Q., K. Kuriyama, K. Nishikawa, S. Tominaga, H. Tatsuta, T. Ando and H. Naka, 2015(b). (Z)-13-Hexadecenylacetate: a novel moth sex pheromone component from *Herpetogramma submarginale* (Lepidoptera: Crambidae). *J. Chem. Ecol.*, **41**: 441-445.
- Yan, Q., M.-Y. Zheng, J.-W. Xu, J.-F. Ma, Y. Chen, Z.-P. Dong, L. Liu, S.-L. Dong and Y.-N. Zhang, 2018(a). Female sex pheromone of *Athetis lepigone* (Lepidoptera: Noctuidae): Identification and field evaluation. *J. Appl. Entomol.*, **142**: 125-130.
- Yan, Q., H.-D. Li, Y. Chen, Z.-F. Ye, X.-Y. You, J. Zhou, L.-F. Mu, S.-J. Liu, X.-B. Kong, S. A. Khuhro and S.-L. Dong, 2018(b). Identification and field evaluation of the sex pheromone of *Orthaga achatina* (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **44**: 886-893.
- Yan, Q., A. Fujino, H. Naka, S.-L. Dong and T. Ando, 2018(c). Chemical analysis of the female sex pheromone in *Palpita nigropunctalis* (Lepidoptera: Crambidae). *J. Asia-Pac. Entomol.*, **21**: 1283-1288.
- Yan, J., M. Zhang, A. Ali, X. Du, X. Mei and Y. Gao, 2022. Optimization and field evaluation of sex-pheromone of potato tuber moth, *Phthorimaea operculella* Zeller (Lepidoptera: Gelechiidae). *Pest Manag. Sci.*, **78**: 3903-3911.
- Yang, C.-Y., S. Lee, K.-S. Choi, H.-Y. Jeon and K.-S. Boo, 2007. Sex pheromone production and response in Korean populations of the diamondback moth, *Plutella xylostella*. *Entomol. Exp. Appl.*, **124**: 293-298.
- Yang, C. Y., J. R. Cho, T. J. Kang and H. Y. Jeon, 2008. Identification and field testing of sex pheromone components of a Korean population of the allium leaf miner, *Acrolepiopsis sapporensis*. *Entomol. Exp. Appl.*, **129**: 216-222.
- Yang, C. Y., K. S. Han and K. S. Boo, 2009(a). Sex pheromones and reproductive isolation of three species in genus *Adoxophyes*. *J.*

Chem. Ecol., **35**: 342-348.

- Yang, C. Y., J. Kim, T. J. Kang and H. Y. Jeon, 2009(b). Identification and field bioassays of the sex pheromone of *Synanthedon haitangvora*. *J. Chem. Ecol.*, **35**: 1197-1201.
- Yang, C. Y., S. J. Kim and S. G. Lee, 2011(a). Identification and field evaluation of the sex pheromone of *Synanthedon bicingulata* (Staudinger). *J. Chem. Ecol.*, **37**: 398-402.
- Yang, C. Y., M. R. Cho and D. R. Choi, 2011(b). Identification and field evaluation of the sex pheromone components of a Korean population of *Glossosphecia romanovi*. *Entomol. Exp. Appl.*, **138**: 244-248.
- Yang, C. Y., C. H. Paik, G. H. Lee and J. Y. Park, 2011(c). Female sex pheromone of the gelechiid moth *Scrobipalpa salinella* (Zeller). *J. Chem. Ecol.*, **37**: 1216-1221.
- Yang, C. Y., H. S. Lee and C. G. Park, 2012. Sex pheromone of the smaller clearwing moth *Synanthedon tenuis* (Butler). *J. Chem. Ecol.*, **38**: 1159-1162.
- Yang, C. Y., K. S. Choi and M. R. Cho, 2013. (*E*)-5-Hexadecenyl acetate: a novel moth sex pheromone component from *Stathmopoda auriferella*. *J. Chem. Ecol.*, **39**: 555-558.
- Yang, M.-H., H.-X. Liu, J.-L. Liu, X.-Y. Jing, J.-T. Zhang, L.-H. Fan and S.-F. Wang, 2015. Sex pheromone components of the carpenterworm moth, *Holcocerus vicarious*. *Entomol. Exp. Appl.*, **154**: 199-205.
- Yang, Y., L. Zhang, F. Guo, Y. Long, Y. Wand and X. Wan, 2016. Reidentification of sex pheromones of tea geometrid *Ectropis oblique* Prout (Lepidoptera: Geometriidae). *J. Econ. Entomol.*, **109**: 167-175.
- Yao, E.-Y., Z.-M. Li, Z.-Q. Luo, Z.-Z. Shang, K.-S. Yin and B.-B. Hong, 1991. Report on structural elucidation of sex pheromone components of a tea pest (*Ectropis obliqua* Prout). *Prog. Nat. Sci.*, **1**: 566-569.
- Yasuda, T., S. Yoshii and S. Wakamura, 1994. Identification of the sex attractant pheromone of the browntail moth, *Euproctis similis* (Fuessly) (Lepidoptera: Lymantriidae). *Appl. Entomol. Zool.*, **29**: 21-30.
- Yasuda, T., S. Wakamura and N. Arakaki, 1995. Identification of sex attractant pheromone components of the tussock moth, *Euproctis taiwana* (Shiraki) (Lepidoptera: Lymantriidae). *J. Chem. Ecol.*,

21: 1813-1822.

- Yasui, H., S. Wakamura, N. Arakaki, H. Irei, C. Kiyuna, H. Ono, H. Yamazawa and T. Ando, 2005. Identification of a sex pheromone component of the geometrid moth *Milionia basalis pryeri*. *J. Chem. Ecol.*, **31**(3): 647-656.
- Yazdani, M., E. Howse, W. T. Tay, H. Spafford and R. D. V. Klinken, 2024. Lure specificity, phenology, and damage caused by *Epiphyas* moths (Lepidoptera: Tortricidae) in Western Australian apple orchards. *J. Econ. Entomol.*, **117**: 1959-1967.
- Yeh, S.-F., K.-C. Lee, K.-T. Chang, F.-C. Yen and J.-S. Hwang, 1989. Sex pheromone components from Asian corn borer, *Ostrinia furnacalis* (Hubner) (Lepidoptera: Pyralidae) in Taiwan. *J. Chem. Ecol.*, **15**: 497-505.
- Youm, O. and P. S. Beevor, 1995. Field evaluation of pheromone-baited traps for *Coniesta ignefusalis* (Lepidoptera: Pyralidae) in Niger. *J. Econ. Entomol.*, **88**(1): 65-69.
- Young, H., R. A. Galbreath, M. H. Benn, V. A. Holt and D. L. Struble, 1985. Sex pheromone components in the New Zealand brown headed leafroller *Ctenopseustis obliquana* (Lepidoptera: Tortricidae). *Z. Naturforsch.*, **40c**: 262-265.
- Yushima, T., Y. Tamaki, S. Kamano and M. Oyama, 1974. Field evaluation of a synthetic sex pheromone, "litlure", as an attractant for males of *Spodoptera litura* (F.) (Lepidoptera: Noctuidae). *Appl. Entomol. Zool.*, **9**: 147-152.
- Zagatti, P., B. Lalanne-Cassou, C. Descoins and M. Gallois, 1983(a). New results on the sex pheromone of the false codling moth, *Cryptophlebia leucotreta* Meyr. (Lepidoptera, Tortricidae). *Agronomie*, **3**: 75-80.
- Zagatti, P., G. A. Bosson, J. Etienne, J. Brenière, C. Descoins and M. Gallois, 1983(b). Sex pheromone of an African rice borer moth: *Chilo zacconius* Blesz. (Lepidoptera, Pyralidae). *C. R. Acad. Sci., Ser. 3*, **296**: 85-88.
- Zagatti, P., A. Hamadoun, M. Lettere, D. Bordat, M. Gallois and C. Malosse, 1988. Sex pheromone of *Sesamia calamistis* (Hampson) (Lepidoptera: Noctuidae). *C. R. Acad. Sci., Ser. 3*, **307**: 837-840.
- Zagatti, P., M. Renou, C. Malosse, B. Frérot, C. Pavis, M. Lettere, C. Descoins, A. Permana, Y. Pivot and F. Leclant, 1991. Sex pheromone of the European sunflower moth, *Homoeosoma nebulellum* (Den. & Schiff.) (Lepidoptera: Pyralidae). *J. Chem. Ecol.*, **17**: 1399-1414.

- Zagatti, P., P. Lucas, P. Genty, S. Arango, C. Malosse and F. Tellier, 1996. Sex pheromone of *Stenoma cecropia* Meyrick (Lepidoptera: Elachistidae). *J. Chem. Ecol.*, **22**: 1103-1121.
- Zarbin, P. H. G., L. M. Lorini, B. G. Ambrogi, D. M. Vidal and E. R. Lima, 2007. Sex pheromone of *Lonomia obliqua*: daily rhythm of production, identification, and synthesis. *J. Chem. Ecol.*, **33** (3): 555-565.
- Zeleny, J., K. Spitzer and J. Ondracek, 1982. Tropical moths (Lepidoptera) attracted by synthetic sex pheromones in Vietnamese agro-ecosystems. *Acta Entomol. Bohemoslov.*, **80**: 413-418.
- Zhan, Y.-D., Y.-J. Liu, J.-H. Liu and Y. Liu, 2023. Pheromones emitted by both female and male moths regulate coordination between the sexes for *Agriphila aeneociliella* (Lepidoptera: Crambidae). *Insect Sci.*, **30**: 1481-1492.
- Zhang, X.-H., G.-Z. Guo, G.-Q. Lin, Y.-W. Wu, P.-H. Wu, Z.-Y. Li and K.-N. Wei, 1986. Studies on identification and syntheses of insect pheromones. XXII Sex pheromone of poplar twig clearwing moth *Paranthrene tabaniformis* Rott -- structure and synthesis. *J. Chem. Ecol.*, **12**: 1263-1271.
- Zhang, Q.-H., E. Priesner, H.-X. Wei and Q.-C. Chen, 1993. Sex-attractant and attraction-inhibitors in Chinese larch casebearer, *Coleophora sinensis* Yang (Lep., Coleophoridae). *J. Appl. Entomol.*, **116**: 516-522.
- Zhang, Q.-H., F. Schlyter, D. Chu, X.-Y. Ma and Y. Ninomiya, 1996. Female calling behaviour and response of males to synthetic pheromone and virgin females in spring and summer generations of *Hyphantria cunea* (Lep., Arctiidae) (Drury) in Northeastern China. *J. Appl. Entomol.*, **120**: 467-476.
- Zhang, J.-T., Han Y. and Meng X.-Z., 2001. Sex pheromone of the carpenterworm, *Holcocerus insularis* (Lepidoptera, Cossidae). *Z. Naturforsch.*, **56c**: 423-429.
- Zhang, A.-B., Z.-J. Wang, S.-J. Tan and D.-M. Li, 2003(a). Monitoring the masson pine moth, *Dendrolimus punctatus* (Walker) (Lepidoptera: Lasiocampidae) with synthetic sex pheromone-baited traps in Qianshan County, China. *Appl. Entomol. Zool.*, **38** (2): 177-186.
- Zhang, A. and S. Polavarapu, 2003(b). Sex pheromone of the cranberry blossomworm, *Epiglaea apiata*. *J. Chem. Ecol.*, **29** (9): 2153-2164.
- Zhang, A. and S. Polavarapu, 2004. Identification of a sex pheromone

- component for the blueberry leafminer, *Caloptilia porphyretica*. *J. Chem. Ecol.* **30**: 1531-1545.
- Zhang, A., T. C. Leskey, J. C. Bergh and J. F. Walgenbach, 2005. Sex pheromone of the dogwood borer, *Synanthedon scitula*. *J. Chem. Ecol.*, **31** (10): 2463-2479.
- Zhang, J.-T., X.-Y. Jing, Y.-Q. Luo, Z.-W. Li, S.-X. Zong and M.-H. Yang, 2009. The sex pheromone of the sand sagebrush carpenterworm, *Holcocerus artemisiae* (Lepidoptera, Cossidae). *Z. Naturforsch.*, **64c**: 590-596.
- Zhang, J., H. Liu, W. Zhao, J. Liu and S. Zong, 2011. Identification of the sex pheromone of *Isoceras sibirica* Alpheraky (Lepidoptera, Cossidae). *Z. Naturforsch.*, **66c**: 527-533.
- Zhang, J.-P., C. Salcedo, Y.-L. Fang, R.-J. Zhang and Z.-N. Zhang, 2012. An overlooked component: (Z)-9-tetradecenal as a sex pheromone in *Helicoverpa armigera*. *J. Insect Physiol.*, **58**: 1209-1216.
- Zhao, C.-H., Q. Li, X.-Y. Guo and X.-Y. Wang, 1993. New components of sex pheromone in the pine caterpillar moth, *Dendrolimus punctatus*: identification of chemical structures and field tests. *Kunchong Xuebao (Acta Entomol. Sinica)*, **36**(2): 247-250.
- Zhao, C.-H., J. G. Millar, K.-H. Pan and C.-S. Xu 1998. Responses of tea tussock moth, *Euproctis pseudoconspersa*, to its pheromone, (R)-10,14-dimethylpentadecyl isobutyrate, and to the S-enantiomer of its pheromone. *J. Chem. Ecol.*, **24** (8): 1347-1353.
- Zhou, Z. L., 1995. Preliminary study on the mass sex pheromone trapping against the grape clearwing moth. *Acta Phytophylacica Sinica*, **22**: 17-21.
- Zhu, Y.-X., Y.-W. Wu, G.-Z. Guo, Z.-X. Tan, G.-Q. Lin, W.-S. Zhou and W.-G. Li, 1981. Determination of the isomeric ratio of the sex pheromone of the pink bollworm moth. *Kexue Tongbao*, **26**: 1073-1077.
- Zhu, P.-C., F.-L. Kong, S.-D. Yu, Y.-Q. Yu, S.-P. Jin, X.-H. Hu and J.-W. Xu, 1987(a). Identification of the sex pheromone of eggplant borer *Leucinodes orbonalis* Guenée (Lepidoptera: Pyralidae). *Z. Naturforsch.*, **42 c**: 1347-1348.
- Zhu, P.-C., F.-L. Kong and Z.-H. Wang, 1987(b). Sex pheromone components of purple stem borer *Sesamia inferens* (Walker). *J. Chem. Ecol.*, **13**: 983-989.

- Zhu, P.-C., F.-L. Kong and Y.-Q. Yu, 1987(c). Sex pheromone of Oriental armyworm *Mythimna separata* Walker. *J. Chem. Ecol.*, **13**: 977-981.
- Zhu, J.-W., M. V. Kozlov, P. Philipp, W. Francke and C. Löfstedt, 1995(a). Identification of a novel moth sex pheromone in *Eriocrania cicatricella* (Zett.) (Lepidoptera: Eriocraniidae) and its phylogenetic implications. *J. Chem. Ecol.*, **21**: 29-43.
- Zhu, J.-W., C. Löfstedt, P. Philipp, W. Francke, T. Tammaru and E. Haukioja, 1995(b). A sex pheromone component to the Geometridae identified from *Epirrita autumnata*. *Entomol. Exp. Appl.*, **75**: 159-164.
- Zhu, J., N. Ryrholm, H. Ljungberg, B. S. Hansson, D. Hall, D. Reed and C. Löfstedt, 1996. Olefinic acetates, $\square$ -9,11-14:OAc and $\square$ -7,9-12:OAc used as sex pheromone components in three geometrid moths, *Idaea aversata*, *I. straminata*, and *I. biselata* (Geometridae, Lepidoptera). *J. Chem. Ecol.*, **22**: 1505-1526.
- Zhu, J., C. Ryne, C. R. Unelius, P. G. Valeur and C. Löfstedt, 1999. Reidentification of the female sex pheromone of the Indian meal moth, *Plodia interpunctella*: evidence for a four-component pheromone blend. *Entomol. Exp. Appl.*, **92**: 137-146.
- Zhu, J. and K. F. Haynes, 2004. Sex pheromone components of the bronzed cutworm, *Nehelodes minians*, a prey species of a bolas spider, *Mastophora hutchinsoni*. *J. Chem. Ecol.* **30**: 2047-2056.
- Zhu, J.-W., S. Polavarapu, K-C. Park, C. Garvey, D. Mahr, S. Nojima, W. Roelofs and T. Baker, 2009. Reidentification of pheromone composition of *Sparganothis sulfureana* (Clemens) and evidence of geographic variation in male responses from two US states. *J. Asia Pacific Entomol.*, **12**: 247-252.
- Zilahi-Balogh, G. M. G., N. P. D. Angerilli, J. H. Borden, M. Meray, M. Tulung and D. Sembel, 1995. Regional differences in pheromone responses of diamondback moth in Indonesia. *Int. J. Pest Manage.*, **41**(4): 201-204.